

# click to campus

## **GPAT 2016 Question Paper with Answer Key**

#### Graduate Pharmacy Aptitude Test conducted by NBEMS

Download GPAT Previous Year Question Papers & Syllabus - Click Here



## **GPAT QUESTION PAPER 2016 WITH ANSWER KEY**

1.	What is the Carr's index o	f good flow powder prope	erty			
	(a) 5-15	(b) 12-16		18-21	(d)	28-35
2.	Which microorganism is i	nvolved in the assay of Ri	famp	icin		
	(a) Bacillus subtilis	(b) Micrococcus luteus	(c)	Bacillus pumilus	(d)	Bacillus cereus
3.	Which percentage of dextr	ose is isotonic with the b	lood	plasma		
	(a) 5%	(b) 10%	(c)	15%	(d)	20%
4.	What is the biological sou		. ,			
	(a) Cassia aungustifolia		(b)	Cassia acutifolia		
	(c) Cassia Bravifolia		(d)	Cassia Nerifolia		
5.	HLB of SPAN and TWEEN	surfactants may be obtain	ned fi	om which of the follow	wing	equations
	(a) $HLB = E/5$	-			_	-
	(b) $HLB = (E+P)/5$					
	(c) $HLB = 20 [(1 - S)/A)]$					
	(d) HLB = (hydrophilic gi	roup numbers)- (lipophili	c gro	up numbers)+7		
6.	If the given drug is absorb	ed by passive what will b	e its a	absorption kinetics		
	(a) Zero Order		(b)	First Order		
	(c) Second order		(d)	Pseudo-zero Order		
7.	Which of the following is o	delayed type of her sensitiv	vity r	reaction		
	(a) Arthus reaction		(b)	Penicillin sensitivity		
	(c) Tubercullin sensitivity	y	(d)	ABO incompatibity		
8.	Which of the following is l	NOT suitable as a post-coit	al co	ntraceptive		
	(a) Levonorgestrel 1.5mg	g	(b)	Ulipristal acetate 30 r	ng	
	(c) Mifepristone 10-25 m	ng	(d)	Mestranol 1.5 mg		
9.	Which of the following pro-	operties are characteristic	cs of	tannis		
	(P) They give a precipitat	e with alkaloids				
	(Q) They give a yellow of	bluish red color with iron	(III)	chloride		
	(R) They transform hide i	nto leather				
	(S) They give a pale-pink	precipipate with iodine				
	(a) P, Q, S	(b) P and Q	(c)	P and R	(d)	Only Q
10.	Adverse drug Event report	ting in the responsibility o	of all	of the following EXCE	PT	
	(a) Pharmacist and physi	cian	(b)	Manufacturer		
	(c) Consumer		(d)	Regulatory authoritie	S	
11.	Time dependent dilatant be	ehavior is knows as				
	(a) Thixotrophy	(b) Rheopexy	(c)	Rheomalaxis		(d) Plastic
12.	Chairman of DTAB is:					
	(a) Health minister of Ind	lia	(b)	Director general of H	ealth	i services
	(c) Drug controller of Ind	lia	(d)	President of AICTE		
13.	Plasmodesmata is:					
	(a) Lignified element		(b)	Vascular element		
	(c) Very fine protoplasmi	c thread	(d)	None		



<ol><li>Efficiency of drug is checked in modest population</li></ol>	n in	
--	------	--

(a) Clinical trail-phase1

. . . . .

- (c) Clinical trail-phase3 (d) Clinical trail-phase4
- 15. Which of the following statements is correct for gram positive becteria
  - (a) Cell wall has a thin peptidoglycan layer
  - (b) Cell wall lipid content is very low and smaller volume of periplasm
  - (c) Lipopolysaccharide layer is present
  - (d) Teichoic acid is present
- 16. The terms upper consulate temperature and lower consulate temperature are related to which phenomenon
  - (a) Cloud point
  - (c) Kraft point
- 17. Match the alkaloids with their synthesis precursors.
  - (1) Pilocarpine
  - (2) Connine
  - (3) Caffine
  - (4) Yohimbine
  - (a) 1-S, 2-R, 3-P, 4-Q
  - (c) 1-P, 2-R, 3-S, 4-Q

(d) Phase inversion

(b) Critical solution temperature

(b) Clinical trail-phase2

- (P) Necleotide
- (Q) Tryptophan
- (R) Histadine
- (S) Acetate derived
- (b) 1-S, 2-Q, 3-P, 4-R
- (d) 1-R, 2-S, 3-P, 4-Q
- 18. Which one of the following is a solid dosage form excipient which can play the role of a diluent, a disintegrant, a glidant, a lubricant and a pore/ channel former.
  - (a) Lactose
  - (c) Ethyl cellulose
- 19. What is the required floor area for running a pharmacy for wholesale or distribution
  - (a) 6 sq meters
  - (c) 15 sq meters

- (d) 30 sq meters
- 20. Bioavailability differences among drug's oral formulations are most likely to occur if it
  - (a) Is freely water soluble
  - (c) Is completely absorbed
- Match the drug with their receptor profiles
  - (1) Ergotamine
  - (2) Ondansetron
  - (3) Sumartriptan
  - (4) Ketanserin
  - (a) 1-R, 2-S, 3-Q, 4-P
  - (c) 1-R, 2-S, 3-P, 4-Q
- 22. What strategy of drug design is frequently used on complex lead compounds derived from natural products
  - (a) Extension
  - (c) Rigidication

- (b) Simplication
- (d) Conformational block

23. Which type of photon detector is commonly microfabricated into arrays of 500 or More individual detectors

- (a) Photocell
- (c) Photumultiplier tube

- (b) Phototube
- (d) Photodiode

- (b) Microcrystalline cellulose
- (d) Eudragit RL 100
- (b) 10 sq meters
- (b) Is incompletely absorbed
- (d) Undergoes little first-pass metabolism
- (Q) 5-HT, partial agonist /antagonist
- (R) 5-HT<sub>3</sub> antagonist
- (S) 5-HT<sub>1D</sub> agonist
- (d) 1-S, 2-R, 3-P, 4-Q

(P) 5-HT<sub>2A</sub> antagonist

- (b) 1-Q, 2-R, 3-S, 4-P



24.	Which of the following is a physe II drug metabolism	reaction associated with genetic polymorphism
	(a) Glucuroinidation	(b) Acytylation
	(c) Reduction	(d) Glutathione conjugation
25.	A gram-negative diplococcus associated with urina	rry tract infection , pelvic inflammatory disease and
	conjunctivitis, meningitis is	
	(a) Neissria gonorrhoeae	(b) Chlamdia Trachomatis
	(c) Hemophilus influenza	(d) Streptococcus pneumoniae
26.	Pregnancy test kits are designed to detect	
	(a) Estrogen	(b) Human chorionic gonadotropin
	(c) Follick-stimulating hormone	(d) Luteinizing hormone
27.	Drug Z is a depolarizing neuromuscular blocking age	ent effective for the treatment of pinworm.
	Identify drug Z	
	(a) Phyrantel	(b) Paramomycin
	(c) Integrase	(d) Protease
28.	Metheneamine used for UTI is a prodrug. How and	to what is is converted into
	(a) At low pH of Urine, to formaldehye	(b) At high pH of urine, to aminosalicylic acid
	(c) At low pH of Urine, to amonosalicylic acid	(d) At high pH of urine, to formalde hye
29.	The correct order for the basic features of a mass sp	pectrometer is
	(a) Acceleration, deflection, detection, ionization	(b) Ionization ,Acceleration, deflection, detection
	(c) Acceleration, ionization, deflection, detection	(d) Acceleration, deflection, ionization, detection
30.	Match the following enzymes/protein with specific for	unctions in DNA replication
	(1) Helicases	(P) Processive unwinding of DNA
	(2) DNA Primses	(Q) Seals the single strand
	(3) DNA ligases	(R) Relieves torsional strain
	(4) Topiosomerases	(S) Initates synthesis of RNA Primers
	(a) 1-P, 2-Q, 3-R, 4-S	(b) 1-P, 2-S, 3-Q, 4-R
	(c) 1-S, 2-Q, 3-P, 4-R	(d) 1-P, 2-Q, 3-R, 4-S
31.	Which is the first line drug for the treatment of gene	ralized seizures
	(a) Valproic acid	(b) Anhydortetracycline
	(c) Carbamazepine	(d) Doxycycline
32	Tetracyline in basic solution is usstable and forms when	hich product
	(a) Epithtracycline	(b) Anhydrotetracycline
	(c) Isotetracycline	(d) Doxycycline
33.	The location of the blood-brain barrier is considered	to be
	(a) At the level of the brain capillaries	(b) At the level of gila
	(c) At the level of neorons	(d) At the level of dendrites
34.	The following drug metabolizing reaction in entirely	non-microsomal:
	(a) Glucuronide conjugation	(b) Acetylation
	(c) Oxidation	(d) Reduction
35.	Which of the following methods is used to determine use	whether a process functions properly for its intended
	(a) Capacity	(b) Inspection
	(c) Validation	(d) Design Review



- 36. Match product, source and plant part form which they areobtained
  - (1) Bacosides (P) Aciacia catechu
  - (2) Cutch

(3) Henna

- (P) Rubiatictorium
- (R) Bacopamonnieri
- (4) Alizarm (S) Lawsoniainternis
- (a) 1-R-ii, 2-S-I, 3Q-Q-iii, 4-P-iv
- (c) 1-Q-ii, 2-P-iii, 3-S-iv, 4-R-I
- 37. What is the significance of term overfill
  - (a) It is similar to overage
  - (b) It is the excess volume to be field in containers as vials, ampoules to avoid loss by degradation
  - (c) It is the excess filing of container as vials, ampoules to avoid the loss during case
  - (d) It is violation of packaging regulation as per GMP

#### 38. Identity the false statements

- (a) A characteristics of drugs eliminated by zero order kinetic process is that the half -life is not constrains
- (b) The plasma drug concertation versus time curve for a drug eliminated by zero order kinetics linear
- (c) A fundamental characteristics of all first order pharmacogenetics processes is that the rate of the process is proportional to drug concentration
- (d) A characteristics of absorption by lipid diffusion is its saturability at high drug concentrations
- 39. 2', 3'-Didehydro-3'-deoxy thymidine is the chemical name of which of following antiviral agents
  - (a) Didenosine (b) Zidovudine (c) Stavudine (d) Zakitabine

40. Oseltamivir is antiviral drug. It produces its action by inhibiting which enzymes

- (a) DNA polymerase (b) Neuraminidase
- (c) Praziquantel (d) Ivermectin

41. In NMR spectrum, a signal is observed as triplet. What will be the ratio of relative peak areas in this signal

(c) 1:3:1 (a) 1:1:1 (b) 1:2:1 (d) 1:4:1

42. Which problem can arise if the material to be compressed into tablet tends to adhere to die walls

- (a) Picking (b) Sticking (c) Capping What is the half life of Tc-99m
- (a) 66 years (b) 66 hours

(a) Phthalate polymers

44. Eudragits are

- (c) 6 hours
  - (d) 60 minutes

(d) Marbling

- (b) Cellulose polymers
- (c) Acrylate polymers (d) Amide polymers 45. Drugs (price control) order 1995 and related orders form time to time are enforeced by (a) NPPA (b) CSIR (c) DBT (d) ICMR
- 46. Match the drugs with plant from which they are isolated and their families (1) Artemisinin (P) Periwinkle (i) Dioscoreaceae (2) Diosgenin (Q) May apple (ii) Apocynaceae
  - (3) Etoposide (R) Sweet wormwood (iii) Berberidaceae (4) Vinblastine and Vicnristine (S) Maxican wild yam (iv) Asteraceae (a) 1-R-iv, 2-S-i, 3-Q-iii, 4-P-ii
  - (c) 1-Q-iii, 2-R-ii, 3-Q-i, 4-P-iv
- (b) 1-s-iv, 2-R-i, 3-Q-iii, 4-P-ii
- (d) 1-R-iv, 2-S-iii, 3-Q-i, 4-P-ii

- (b) 1-R-I, 2-P-iv, 3-S-ii, 4-Q-i
- (d) 1-S-ii, 2-R-iv, 3-P-I, 4-Q-i
- (iii) Root (iv) Stem

(i) Herb

(ii) Leaf



47.	Which of the following	is an irreversible phenom	enonralated to	stability of emulsi	ion
	(a) Craking		(b) Crean	ning	
	(c) Coalescence		(d) Floccu	ulation	
48.	If a drug is highly boun	d to plasma proteins, wha	t might be its i	reason or consequ	ience
	(a) It most likely carri	ed by $\alpha$ -glycorprotien	(b) It has	a high renal clear	ance
	(c) It has a large Vd		(d) It is a	likely candidate fo	or drug interactions
49.	In order to make a gene	ric substitution ; a pharma	icist must do a	also act as a hydro	gen bond acceptor
	(a) Notify the patient of	f the substitution		ž	
	(b) Charge the same or	lower price for the gener	ic		
	(c) Place the brand nan	ne on the label and write "	substitute for"	,	
	(d) Obtain the physician	n's consent to substitute tl	ne product		
50.	Which of the following g	groups can form ionic inte	ractions and a	also act as a hydrog	gen bond acceptor
	(a) Hydroxyl group (Oł	4)	(b) Carbox	xylate group (RCO	0)
	(c) Aminononium grou	p (RNH <sup>3+</sup> )	(d) Ketone	e (C=O)	
51.	Which of the following of	drug does not give pink col	our with ruth	enium red	
	(a) Agar	(b) Guar gum	(c) Pectin	(d)	) Isabgol
52.	The IUPAC name, 4-Amin	10-N(5,6-dimethoxy-4-pyr	imidinly) benz	zenseulfonamide b	elong to which generic
	drug				
	(a) Sulfadimidine	(b) Sulfadoxine	(c) Sulfale	ne (d)	) Sufamerazine
53.	Method of inspections u	sed to determine the abso	rption rate co	nstants. It assumes	s that
	(P) Ka is at least five tin	ne grater activities			
	(Q) Absorption in comp	lete (i.e. > 95% complete)	at the time of	peak concentratio	n
	(R) Both Absorption an	nd elimination are first or	der processes		
	(a) P and Q	(b) Q and R	(c) Q and	R (d)	) P and R
54.	The clinical trial registry	in India is maintained by			
	(a) World health organi				
		edical research , New Delł	i		
		Research, New Delhi			
		ard control Organization, I			
55.		r for unsaturation in follo	0		
	(1) Palmitoleic Acid		(2) Linoler		
	(3) Linoleic acid		(4) Arachi		
	(a) 1>2>3>4	(b) 3>4>2>1	(c) 4>3>2	.>1 (d	) 4>2>3>1
56.		s an example of Diaz oniu			
	(a) $CH_3^+N_3^-$	(b) $CH_{3}N_{2}^{+}$	(c) $H_2$ N-N	-	) None of these
57.		The degree to which a set	of inherent pi	roperties of a proc	luct system , of rocess
	fulfils requirements, the				
50	(a) Standard	(b) Quality	(c) Quality		) State of control
58.		e of lidocaine if the admir	istered dose i	is 0.2 g and the to	Dial body clearance is
	45 L/h	(b) $0.0044$ b m $\pi^{4}$	(a) 0.00 b	am /I ca	0.000 h mat
	(a) 4.44 h.mg/L	(b) 0.0044 h.mg/L	(c) 9.00 h.	ւցույս (a	) 9000 h. mg/L



59.	Toluene is converted of which compound in presence	e of	CrO <sub>3</sub> with acetic anhydride
	(a) Benzyl alcohol	(b)	Benzaldehyde
	(c) Benzoic acid	(d)	Benzoin
<del>60</del> .	Match the antimalarial drugs with their modes of act	ion	
	(1) Artemisinin	(P)	Inhibition of parasite mitochondrial electron
			Tranport
	(2) Pyremethamine	(Q)	Inhibition of heme polymerase
	(3) Quinine	(R)	Generation of oxygen and carbon- centered
			redicals
	(4) Atovaquone/Proguanil	(S)	Inhibition of dihydrofolate reductase
	(a) 1-P, 2-S, 3-Q, 4-R	(b)	1-Q, 2-S, 3-P, 4-R
	(c) 1-S, 2-R, 3-Q, 4-P	(d)	1-R, 2-S, 3-Q, 4-P
61.	Methyl ether of erythromycin is		
	(a) Carithromycin	(b)	Dirithromycina
	(c) Azithromycin	(d)	Mithramycin
62.	Ebullioscopicmethod is based on which of the follow	ving	observation
	(a) Freezing point depression	(b)	Boiling point elevation
	(c) Osmotic pressure change	(d)	None of the above
63.	How many fundamental svibrations can be expected	l for	C <sub>2</sub> H <sub>5</sub> CL
	(a) 7 (b) 11	(c)	14 (d) 18
<u>64</u> .	Which of the antibodies provide passive immunity to	o nev	vborn baby
	(a) lgG (b) lgM	(c)	lgA (d) lgE
65.	Increased number of number mitosis may be presen	nt in	the following tissue EXCEPT
	(a) Bone marrow cells (b) Nails	(c)	Hepatocytes (d) Intestinal Cells
66.	To which chemical class the vinca alkaloids belong		
	(a) Tropane (b) Indole		Tryptopha (d) Purine
67.	An antibiotic thatn resembles the 3' end of a charged	d tRN	IA molecule is
	(a) Streptomycin (b) Vincamycin	` '	Puromycin (d) Tetracycline
68.	If the cohesive forces in between similar molecules a		-
	molecules, a deviation in Raolt's law is observed . her		
	(a) Positive	(b)	Negative
	(c) Absent		Either positive or negative
69.	Phenylalanine, a precursor of most of the phenolic	s is	higher plants is a product of which one of the
	following pathways		
	(a) Shikimic acid pathway	. ,	Malonic acid pathway
	(c) Mevalonic acid pathway	(d)	Methylerhtritol pathway



70.	Within how many days a pharmacist should dispe	ense diluted ac	ueous mixtures	
	(a) 7 days (b) 14 days	(c) 21 day	/s (d) 30	) days
71.	What molecular feature is penicillin G is said o min	nic		
	(a) Disaccharide of N-acetylmuranicanidN- actylg	ukosamine		
	(b) N-acetylneuraminic acid			
	(c) The pentapeptide moiety of five glycine units			
	(d) The dipeptide moiety D-ala-D-Ala			
72.	If a drug is known to be distributed into total bod	ly water, how	many milligrams are n	eeded to obtain an
	initial plasma level of 5mg/L in a patient weighting	g 70 kg		
	(a) 210 (b) 150	(c) 50	(d) 35	5
73.	What does 'pharmacokinetical compartment' mea	in		
	(a) Part of the body water which is located is the	e vascular syste	em	
	(b) Total body water			
	(c) Plasma, intracellular fluid, together , anatomic	al water comp	artments where drug i	s absorbed
	(d) Part of the body water in which the change o	f a drug conce	ntration has the same	kinetics
74.	The resistance to macrolide antibiotics by of gran	n positive orga	nism is deve <b>loped due</b>	to
	(a) Decreases uptake of antibiotics			
	(b) Synthesis of esterase enzyme that hydrolyzes	s lactone ring o	of macrolide	
	(c) Methylation of 50S subunit at the antibiotic b	inding site		
	(d) Increased metabolism of antibiotic			
75.	The ethanolic solution contaminated with benzele	showed abso	rbance of 0.69 at 260 r	nm in a 2 cm cell if
	the molar absorptivity of benzene in thanolis 2	30 M <sup>-1</sup> cm <sup>-1</sup> , w	hat is the concentrati	on of benzene in
	the solution			
	(a) 0.003 M (b) 0.0015 M	(c) 0.001	M (d) 0.4	015M
76.	The most effective agent for treating psychosis w	ould be		
	(a) Buspirone	(b) Sertal	ne	
	(c) Dextroamphetamine	(d) Olanza	pine	
77.	TGA is regulatory body of which country			
	(a) Europe (b) Australia	(c) Canad	a (d) UI	K
78.	Which prostaglandins have a keto function at C-	9 and a α- Hy	droxyl group at C-11	in prostanoic acid
	backbone			
	(a) PGA (b) PGI	(c) PGE	(d) PC	GF
79.	When morphine is heated at 140 $^{\circ}\mathrm{C}$ under pressure $% \mathrm{C}$	re, with strong	HCL, it converts pnto:	
	(a) Morphinone	(b) Apom	orphine	
	(c) Codeine	(d) Oxym	orphine	



80.	All of the following are gram-negative rods EXC	EPT
	(a) Clostridium (b) Escherichia	(c) Salmanella (d) Shigella
81.	The cells which secrets male sexhormone testos	sterone are:
	(a) Crypts of lieberkuhn	(b) Escherichia
	(c) Salmonella	(d) Shigella
82.	If QA and QC are compared	
	(a) Both are literally the same	
	(b) QA is a higher activity in the management h	nierarchy
	(c) QA is a higher activity in the management h	nierarchy
	(d) QA is a done by the production person and	QC is done by analyst
83.	Bio availability of drug refers to	
	(P) The ratio of drug excreted unchanged in un	rine to that excreted as metabolites
	(Q) Fraction of the drug reaching the target to $\label{eq:Q}$	produce the action
	(R) The length of time an administered drug is	available for action
	(S) Percentages of administered dose that reac	hes systemic circulation in the unchanged form
	(a) Only P	(b) Q and R
	(c) R and S	(d) Only S
84.	Following intravenous administration, drugs are	e distributed fastest to
	(a) The skin, kidney, and brain	(b) The liver, kidney, and brain
	(c) The liver, adipose, and brain	(d) The liver, kidney, and adipose
85.	Which of the following agents act as hypoglycen	-
	(a) Mitiglinide	(b) Pioglitazone
	(c) Liraglutide	(d) Sitagliptin
86.	Carabilide shows stongs IR absorption in which	
	(a) 3200-3600	(b) 1640-1690
	(c) 1000-1300	(d) 2210-2260
87.	Match the drugs with their adverse effects	
	(1) Cyclophosphamide	(P) Pulmonary fibrosis
	(2) Doxorubicin	(Q) Nephrotoxicity, ototoxicity
	(3) Bleomycin	(R) Acute hemorrhage cystitis
	(4) Cisplatin	(S) Cardiotoxicity
	(a) 1-S, 2-R, 3-P, 4-Q	(b) 1-P, 2-Q, 3-S, 4-R
	(c) 1-P, 2-S, 3-Q, 4-R	(d) 1-R, 2-S, 3-P, 4-Q
88.	Which one of the following technique is used to	· ·
	(a) X-ray diffractometry	(b) Raman spectroscopy
	(c) Differential scanning calorimetry	(d) Atomic forced microscopy



89.	Mat	ch the schedules with	the particulars they describ	oe 🛛				
	(1)	Schedule T		(P)	Standards for patent of	or pr	oprietary medicines	
	(2)	Schedule U		(Q)	Requirements/ guid manufactures new Dr		nes to import &/or	
	(3)	Schedule V		(R)	GMP practices for Ay medicines	urv	edic, siddha & Unani	
	(4)	Schedule Y		(S)	Particulars to be sho records	wn i	n the manufacturing	
	(a)	1-R, 2-S, 3-Q, 4-P		(b)	1-S, 2-Q, 3-P, 4-R			
	(c)	1-R, 2-S, 3-P, 4-Q		(d)	1-S, 2-R, 3-P, 4-Q			
90.	Wh	ich of the following UV	<sup>7</sup> rays cause cancer					
	(a)	UVA	(b) UVB	(c)	UVC	(d)	All of the above	
91.	Wh	ich are the types of an	tibodies involved in hypers	sensi	tivity reactions			
	(a)	LgG and LgD	(b) LgG and LgM	(c)	LgD and LgA	(d)	LgM and LgD	
92.	The	term used to describe	unequal distribution of co	olour	on a tablet is			
	(a)	Chipping	(b) Mottling	(c)	Lamination	(d)	Double impress	
93.	Why	y acetyl chloride under	goes nucleophilic substitu	tion	at a faster <mark>rate than m</mark>	ethy	l acetate	
	(a)	The ester is more ste	rically hindered than the a	cid c	hloride			
	(b) The chloride ion is a better leaving group than methoxide							
	(c)	The acid chloride is m	ore sterically hindered that	n th	e ester			
	(d)	The methoxide ion is	a better leaving group that	n chle	oride			
94.	The	key concept of Total Q	uality Managements (TQM	) is				
	(a)	Total control of all qua	ality related activities					
	(b)	Commitment of all em	ployees to quality improv	eme	nt and having team me	etin	gs	
	(c)	Top management's di	rect involvement					
	(d)	The Introduction ot th	ne ISO 9000 Series					
95.	A dı	rug of low water solubi	lity when given orally is ab	sorb	ed up to 90% of the ad	mini	stered does. The drug	
	belo	ongs to which class acc	ording to BCS classification	n				
	(a)	Class IV	(b) Class III	(c)	Class II	(d)	Class I	

96. Which of the following is NOTa component of evaporator (c) Condenser (a) Heat exchange (b) Vacuum separator (d) Cyclone separator 97. In parkinson's disease, there is a predominant loss of dopaminergic neurons (a) Substantia (b) Cerebellar (c) Cerebral cortext (d) Locus ceruleus

- 98. At equilibrium the receptor occupancy is related to drug concentration by
  - (a) Henderson-Hasselbach equation (b) Hill- Langmuir equation
  - (c) Lineweaver-Burk equation (d) Langmuir adsorption isotherm



99.	Which method is not suitable to calculate area under	the	curve	
	(a) Least square method	(b)	Weighing and platometer	ery
	(c) Trapezoid rule	(d)	Integration of curve	
100.	OROS is a technology developed for/as			
	(a) Oral release rapid onset system	(b)	Orally rapid disintegra	ting tablets
	(c) Osmotic controlled oral drug delivery system	(d)	Transdermal drug deliv	very system
101.	Match the events in tablet manufacturing process with	th th	e effects found in tables	
	(1) Rapid drying of coated tablets after coaching	(P)	Increased distegration	time
	(2) Use to highly viscous solution	(Q)	Weight variation	
	(3) Improper feed rate form hopper	(R)	Orange peel	
	(4) Excessive compression force	(S)	Blistering	
	(a) 1-R, 2-S, 3-Q, 4-P	(b)	1-R, 2-S, 3-P, 4-Q	
	(c) 1-S, 2-R, 3-Q, 4-P	(d)	1-R, 2-P, 3-S, 4-Q	
102.	At pH 5, the ratio of the protonated to unprotanated	t fori	ns of morphine pKa 7 v	would be
	(a) 1:100 (b) 1:10	(c)	10:1 (	(d) 100:1
103.	What structure is formed if the acyl side chain of per	nicill	in is hydro <b>lysed</b>	
	(a) Penicellienic acids	(b)	Penillic acids	
	(c) 7-Aminopenicillanic acid	(d)	6-Aminopenicillanic ac	id
104.	The clinical trial is being conducted with 1500 volu	nteei	rs which m <b>ay spans on</b>	period of 2 years as per
	protocol. The clinical trial is in which phase			
	(a) Phase 1 (b) Phase 2	(c)	Phase 3	(d) Phase 4
105.	Which of the following drug has not undergone a clin	nical	trial for its use to decrea	ase intracranial pressure
	(a) Dideoxyinosine (b) Zidovudine	(c)	Acetazolamide (	(d) Nicotine
106.	Colloidal depression have which type of rheology			
	(a) Newtonian (b) Pesudoplastic			(d) Dilatant
107.	Which of the reactive oxygen species is most danger			
	(a) Singlet oxygen (b) Hydroxyl radical	(c)	Superoxide (	(d) Peroxide
108.	The Gibb's Phase rule			
	(a) Holds only for systems with more than compon			
	(b) Predicts that a maximum of three phase can exi			n
	(c) Does not count phase compositions as intensive			
	(d) Does not count pressure and temperature as in			
109.	Nerve impulse from the cochlea arrive first in which			
	(a) Auditory cortex	• •	Thalamus	
	(c) Medulla oblongata	(d)	Inferior colliculus	



110. The p	product of a Michael re	eactior	n of a ketone enolate	to ai	na, b-unsaturated keto	ne is	And addition
react	tion occurs in a/an						
(a) 1	1,5- deketone; 1,4-fas	hion		(b)	$\alpha$ -substituted acetate;	; 1,2	fashion
(c) (	3 - hydorxy keto; 1,3	– fashi	on	(d)	$\alpha,\beta$ - keto ester; 1,5 $\cdot$	-fasł	nion
111. Whic	ch of the following is a	a drug	considered as potas:	sium	sparing diuretic		
(a) 1	Trimatrine	(b) C	hlorthiazide	(c)	Mannitol	(d)	Furosemide
112. Whic	ch of the reagent from	n the gi	iven can be used to	prot	ect ketone group		
(a) A	Acidic methanol	(b) B	asic methanol	(c)	Methanol + KCN	(d)	Phenobarbitone
113. Whic	ch of the following dru	igs cau	ises less inhibition o	r RE	M sleep		
(a) Z	Zolpidem	(b) E	thanol	(c)	Lorazepam	(d)	Phenobarbitone
	starting materials for						
(a) 4	4- Aminobenzene-1-s	ulfonyl	l amide + 3-chloro-5	-met	hyl isoxazole		
	4- Aminobenzene-1-s	-			•		
	4- Aminobenzene-1-s	-			-		
	4- Aminobenzene-1-s	-			•		
	h the following plant p	produc	t with their chemica				
	b-amyrin				Alkaloid secondary al	coho	1
	Sqaulene				Alkaloid, phenol		
	Morphine				Triterpene, secondary	·	
	Ephedrine				Asyclic triterpene, po	lyen	e
	1-R, 2-S, 3-Q, 4-P			` `	1-S, 2-Q, 3-P, 4-R		
	1-P, 2-S, 3-Q, 4-R			` ´	1-R, 2-S, 3-P, 4-Q		
	••••	one ai	re subject to therap		drug monitoring, Whi	ich o	ne
	Phenytoin				Lithium		
	Gentamicin				Losartan		
	mine deficiency cause			ction	because		
• • •	It is required for the	•					
	It is a co-factor in oxid			,	1		
	It is a co-enzyme for t			-			
	-	••		•	ha ketoglutarate dehyd	-	
		gs are o	often found in both pi		iption and over-the-co	unte	r nasal decongestands
. ,	Alpha 2 agonists				Alpha 1 agonists		
	Alpha 1 antagonists	n of	aton of at 2500	(a)	Beta 2 agonists		
	t is the surface tensio	II OI W	ater of at 25°C	ക	68 duna /cm		
	58 dyne/cm 72 dyne/cm				68 dyne/cm 82 dyne/cm		
	- a ayne/ cm			(u)	on ayne/en		



120. Acridine and xanthene rings are related to each o	other in	that
(a) Xanthene is oxygen isoster of acridine	(b)	Acridine is
(c) Xanthene is nitrogen isoster of aciridine	(d)	Xanthene is

- s oxygen isoster of xanthene
- (d) Xanthene is sulfur isoster os acridine
- 121. Colligative properties depend on.....
  - (a) Structural arrangement of atoms within the molecules of solute and solvent
  - (b) The number of solute particles is solution
  - (c) The physical properties of the solute particles dissolved is solution
    - (d) Sum of the corresponding properties of individual atoms or functional group within the molecules
- 122. In polarography.....current must be blocked
  - (a) Residual (b) Migration (c) Diffusion (d) None
- 123. The propellant commonly used in topical aerosols is
  - (a) Trichloromonfluoromethane
  - (c) Dichlordifluromthane

- 124. Which of the following increase systolic and diastolic pressure in normal patient
  - (a) Epineherine
  - (c) Tyramine
- 125. A large Reynold number is indication of which type of flow
  - (a) Smooth and stream line flow
  - (c) Steady flow

- (b) Laminar flow
- (d) Highly turbulent flow

#### ANSWER KEY GPAT 2016

			<u> </u>						
1-b	2-a	3-a	4-b	5-c	6-b	7-c	8-d	9-c	10-d
11-b	12-b	13-с	14-b	15-b	16-b	17-d	18-b	19-с	20-b
21 <b>-</b> b	22-b	23-d	24-b	25-a	26-b	27-a	28-a	29-b	30-b
31-a	32-c	33-a	34-b	35-c	36-b	37-с	38-d	39-с	40-b
41-b	42-b	43-c	44-a	45-a	46-a	47-b	48-d	49-a	50-d
51-b	52-b	53-a	54-b	55-d	56-b	57-b	58-a	59-b	60-d
61-a	62-b	63-d	64-c	65-b	66-b	67-c	68-b	69-a	70-b
71-d	72-a	73-d	74-b	75-b	76-d	77-b	78-с	79-b	80-a
81-b	82-b	83-d	84-b	85-a	86-b	87-d	88-c	89-c	90-b
91-b	92-b	93-b	94-c	95-c	96-d	97-a	98-b	99-a	100-с
101-с	102-c	103-d	104-с	105-c	106-с	107-b	108-b	109-с	110-a
111-а	112-a	113-a	114-b	115-a	116-a	117-d	118-b	119-с	120-a
121-b	122-b	123-b	124-c	125-d				19	

- (b) Trifluromonfluroethane
- (d) Isopropyl alcohol

(b) Norepinephrine

(d) Phenylephrine