

click to campus

JEE Main 2019 Question Paper with Answer

January & Other (All Shifts)

JEE Main 2019 – 9th Jan Shift 1 Question Paper	Page No. 2 to 25
JEE Main 2019 – 9th Jan Shift 2 Question Paper	Page No. 26 to 48
JEE Main 2019 – 10th Jan Shift 1 Question Paper	Page No. 49 to 71
JEE Main 2019 – 10th Jan Shift 2 Question Paper	Page No. 72 to 93
JEE Main 2019 – 11th Jan Shift 1 Question Paper	Page No. 94 to 115
JEE Main 2019 – 11th Jan Shift 2 Question Paper	Page No. 116 to 135
JEE Main 2019 – 12th Jan Shift 1 Question Paper	Page No. 136 to 165
JEE Main 2019 – 12th Jan Shift 2 Question Paper	Page No. 166 to 179
JEE Main 2019 – Other Question Paper	Page No. 180 to 365

Download more JEE Main Previous Year Question Papers: Click Here

TEST PAPER OF JEE(MAIN) EXAMINATION - 2019 (Held On Wednesday 09th JANUARY, 2019) TIME : 9 : 30 AM To 12 : 30 PM MATHEMATICS

3.

1. The area (in sq. units) bounded by the parabola $y = x^2 - 1$, the tangent at the point (2, 3) to it and the y-axis is :

(1) $\frac{14}{3}$ (2) $\frac{56}{3}$ (3) $\frac{8}{3}$ (4) $\frac{32}{3}$

Ans. (3)

Sol.

K –axis v = (4x - 5)

Equation of tangent at (2,3) on $y = x^2 - 1$, is y = (4x - 5)....(i) :. Required shaded area

= ar (
$$\Delta ABC$$
) $-\int_{-1}^{1} \sqrt{y+1} dy$
= $\frac{1}{2} \cdot (8) \cdot (2) - \frac{2}{3} ((y+1)^{3/2})_{-1}^{3}$
= $8 - \frac{16}{3} = \frac{8}{3}$ (square units)

2. The maximum volume (in cu. m) of the right circular cone having slant height 3m is :

> (1) $3\sqrt{3} \pi$ (2) 6π (4) $\frac{4}{3}\pi$ (3) $2\sqrt{3} \pi$

Ans. (3)

Sol.
h
h

$$e^{h} = 3 (given)$$

 $\therefore h = 3 \cos \theta$
 $r = 3 \sin \theta$
Now,
 $V = \frac{1}{3}\pi r^{2}h = \frac{\pi}{3}(9\sin^{2}\theta).(3\cos\theta)$

 $\therefore \frac{dV}{d\theta} = 0 \implies \sin \theta = \sqrt{\frac{2}{2}}$ $\frac{d^2 V}{d\theta^2} \bigg]_{\sin\theta = \sqrt{\frac{2}{3}}} = \text{negative}$ Also, \Rightarrow Volume is maximum, $\sin\theta = \sqrt{\frac{2}{2}}$ when $\therefore V_{max}\left(\sin\theta = \sqrt{\frac{2}{3}}\right) = 2\sqrt{3}\pi$ (in cu. m) For $x^2 \neq n\pi + 1$, $n \in N$ (the set of natural numbers), the integral $\int x \sqrt{\frac{2\sin(x^2-1) - \sin 2(x^2-1)}{2\sin(x^2-1) + \sin 2(x^2-1)}} dx$ is equal to : (where c is a constant of integration) (1) $\log_{e} \left| \sec \left(\frac{x^2 - 1}{2} \right) \right| + c$ (2) $\log_{e} \left| \frac{1}{2} \sec^{2} (x^{2} - 1) \right| + c$ (3) $\frac{1}{2}\log_{e}\left|\sec^{2}\left(\frac{x^{2}-1}{2}\right)\right|+c$ (4) $\frac{1}{2}\log_{e}|\sec(x^{2}-1)|+c$ Ans. (1) **Sol.** Put $(x^2 - 1) = 1$ $\Rightarrow 2xdx = dt$ $\therefore I = \frac{1}{2} \int \sqrt{\frac{1 - \cos t}{1 + \cos t}} dt$

$$= \frac{1}{2} \int \tan\left(\frac{t}{2}\right) dt$$
$$= \ln\left|\sec\left(\frac{t}{2}\right)\right| + c$$
$$I = \ln\left|\sec\left(\frac{x^2 - 1}{2}\right)\right| + c$$

E

collegebatch.com

JEE (Main) Examination-2019/Morning Session/09-01-2019

4. Let α and β be two roots of the equation $x^{2} + 2x + 2 = 0$, then $\alpha^{15} + \beta^{15}$ is equal to : (2) - 512(3) - 256(1) 512(4) 256 Ans. (3) Sol. We have $(x + 1)^2 + 1 = 0$ \Rightarrow (x + 1)² - (i)² = 0 \Rightarrow (x + 1 + i) (x + 1 - i) = 0 $\therefore \qquad \mathbf{x} = -(1+\mathbf{i}) \quad -(1-\mathbf{i})$ So, $\alpha^{15} + \beta^{15} = (\alpha^2)^7 \alpha + (\beta^2)^7 \beta$ = -128 (-i + 1 + i + 1)= -2565. If y = y(x) is the solution of the differential equation, $x\frac{dy}{dx} + 2y = x^2$ satisfying y(1) = 1, then $y\left(\frac{1}{2}\right)$ is equal to : (1) $\frac{7}{64}$ (2) $\frac{13}{16}$ (3) $\frac{49}{16}$ (4) $\frac{1}{4}$ Ans. (3)**Sol.** $\frac{dy}{dx} + \left(\frac{2}{x}\right)y = x$ \Rightarrow I.F. = x^2 :. $yx^2 = \frac{x^4}{4} + \frac{3}{4}$ (As, y(1) = 1) $\therefore \quad y\left(x=\frac{1}{2}\right) = \frac{49}{16}$ Equation of a common tangent to the circle, 6. $x^2 + y^2 - 6x = 0$ and the parabola, $y^2 = 4x$, is: (1) $2\sqrt{3} y = 12 x + 1$ S (2) $2\sqrt{3}y = -x - 12$ (3) $\sqrt{3} y = x + 3$ (4) $\sqrt{3} y = 3x + 1$ Ans. (3) **Sol.** Let equation of tangent to the parabola $y^2 = 4x$ is $y = mx + \frac{1}{m}$, \Rightarrow m²x-ym+1 = 0 is tangent to x² + y² - 6x = 0 $\Rightarrow \frac{|3m^2+1|}{\sqrt{m^4+m^2}}=3$

$$m = \pm \frac{1}{\sqrt{3}}$$

$$\Rightarrow \text{ tangent are } x + \sqrt{3}y + 3 = 0$$

and $x - \sqrt{3}y + 3 = 0$

7. Consider a class of 5 girls and 7 boys. The number of different teams consisting of 2 girls and 3 boys that can be formed from this class, if there are two specific boys A and B, who refuse to be the members of the same team, is:

Ans. (2)

Sol. Required number of ways

= Total number of ways – When A and B are always included.

$$= {}^{5}C_{2} \cdot {}^{7}C_{3} - {}^{5}C_{1} \cdot {}^{5}C_{2} = 300$$

8. Three circles of radii a, b, c(a < b < c) touch each other externally. If they have x-axis as a common tangent, then :

(1)
$$\frac{1}{\sqrt{a}} = \frac{1}{\sqrt{b}} + \frac{1}{\sqrt{c}}$$

(2) a, b, c are in A. P.

(3)
$$\sqrt{a}, \sqrt{b}, \sqrt{c}$$
 are in A. P.

$$(4) \quad \frac{1}{\sqrt{b}} = \frac{1}{\sqrt{a}} + \frac{1}{\sqrt{c}}$$

Ans. (1)

ol.

$$\frac{b}{A} = \frac{c}{B} = axis$$

$$AB = AC + CB$$

$$\sqrt{(b+c)^{2} - (b-c)^{2}}$$

$$= \sqrt{(b+a)^{2} - (b-a)^{2}} + \sqrt{(a+c)^{2} - (a-c)^{2}}$$

$$\sqrt{bc} = \sqrt{ab} + \sqrt{ac}$$

$$\frac{1}{\sqrt{a}} = \frac{1}{\sqrt{c}} + \frac{1}{\sqrt{b}}$$

2



If the fractional part of the number $\frac{2^{403}}{15}$ is $\frac{k}{15}$, 9. then k is equal to : (1) 14(2) 6(4) 8(3) 4Ans. (4) **Sol.** $\frac{2^{403}}{15} = \frac{2^3 \cdot (2^4)^{100}}{15} = \frac{8}{15} (15+1)^{100}$ $=\frac{8}{15}(15\lambda+1)=8\lambda+\frac{8}{15}$ \therefore 8 λ is integer \Rightarrow fractional part of $\frac{2^{403}}{15}$ is $\frac{8}{15} \Rightarrow k = 8$ 10. Axis of a parabola lies along x-axis. If its vertex and focus are at distances 2 and 4 respectively from the origin, on the positive x-axis then which of the following points does not lie on it ? (1) (4, -4)(2) $(5, 2\sqrt{6})$ (4) 6. $4\sqrt{2}$ (3) (8, 6) Ans. (3) v-axis Sol. → x-axis A(2,0)equation of parabola is $y^2 = 8(x - 2)$ (8, 6) does not lie on parabola. 11. The plane through the intersection of the planes x + y + z = 1 and 2x + 3y - z + 4 = 0 and parallel to y-axis also passes through the point : (1) (-3, 0, -1)(2) (3, 3, -1)(3) (3, 2, 1)(4) (-3, 1, 1)Ans. (3) Sol. Equation of plane $(x + y + z - 1) + \lambda(2x + 3y - z + 4) = 0$ $\Rightarrow (1+2\lambda)x + (1+3\lambda)y + (1-\lambda)z - 1 + 4\lambda = 0$ dr's of normal of the plane are $1 + 2\lambda, 1 + 3\lambda, 1 - \lambda$ Since plane is parallel to y - axis, $1 + 3\lambda = 0$ \Rightarrow $\lambda = -1/3$ So the equation of plane is $\mathbf{x} + 4\mathbf{z} - 7 = 0$ Point (3, 2, 1) satisfies this equation Hence Answer is (3)

12. If a, b and c be three distinct real numbers in
G. P. and
$$a + b + c = xb$$
, then x cannot be :
(1) 4 (2) -3 (3) -2 (4) 2
Ans. (4)

Sol.
$$\frac{b}{r}$$
, b, br \rightarrow G.P. $(|r| \neq 1)$
given $a + b + c = xb$
 $\Rightarrow b/r + b + br = xb$
 $\Rightarrow b = 0$ (not possible)

or
$$1+r+\frac{1}{r}=x \implies x-1=r+\frac{1}{r}$$

 $\Rightarrow x-1 > 2 \text{ or } x-1 < -2$
 $\Rightarrow x > 3 \text{ or } x < -1$
So x can't be '2'

- 13. Consider the set of all lines px + qy + r = 0 such that 3p + 2q + 4r = 0. Which one of the following statements is true ?
 - (1) The lines are all parallel.
 - (2) Each line passes through the origin.
 - (3) The lines are not concurrent The lines are concurrent at the point

$$(4) \left(\frac{3}{4}, \frac{1}{2}\right)$$

Sol. Given set of lines px + qy + r = 0given condition 3p + 2q + 4r = 0

$$\Rightarrow \frac{3}{4}p + \frac{1}{2}q + r = 0$$

$$\Rightarrow$$
 All lines pass through a fixed point $\left(\frac{3}{4}, \frac{1}{2}\right)$.

14. The system of linear equations.

x + y + z = 2 2x + 3y + 2z = 5 $2x + 3y + (a^{2} - 1)z = a + 1$ (1) has infinitely many solutions for a = 4 (2) is inconsistent when $|a| = \sqrt{3}$ (3) is inconsistent when a = 4 (4) has a unique solution for $|a| = \sqrt{3}$

Ε



Ans. (2) **Sol.** $D = \begin{vmatrix} 1 & 1 & 1 \\ 2 & 3 & 2 \\ 2 & 3 & a^2 - 1 \end{vmatrix} = a^2 - 3$ $D_{1} = \begin{vmatrix} 2 & 1 & 1 \\ 5 & 3 & 2 \\ a+1 & 3 & a^{2}-1 \end{vmatrix} = a^{2} - a + 1$ $D_{2} = \begin{vmatrix} 1 & 2 & 1 \\ 2 & 5 & 2 \\ 2 & a+1 & a^{2}-1 \end{vmatrix} = a^{2} - 3$ $D_3 = \begin{vmatrix} 1 & 1 & 2 \\ 2 & 3 & 5 \\ 2 & 3 & a+1 \end{vmatrix} = a - 4$ D = 0 at $|a| = \sqrt{3}$ but D₃ = $\pm \sqrt{3} - 4 \neq 0$ So the system is Inconsistant for $|a| = \sqrt{3}$ Let $\vec{a} = \hat{i} - \hat{j}$, $\vec{b} = \hat{i} + \hat{j} + \hat{k}$ and \vec{c} be a vector 15. such that $\vec{a} \times \vec{c} + \vec{b} = \vec{0}$ and $\vec{a} \cdot \vec{c} = 4$, then $|\vec{c}|^2$ is equal to :-(1) $\frac{19}{2}$ (2) 8 (3) $\frac{17}{2}$ (4) 9 Ans. (1) Sol. $\vec{a} \times \vec{c} = -\vec{b}$ $(\vec{a} \times \vec{c}) \times \vec{a} = -\vec{b} \times \vec{a}$ \Rightarrow $(\vec{a} \times \vec{c}) \times \vec{a} = \vec{a} \times \vec{b}$ \Rightarrow $(\vec{a}.\vec{a})\vec{c}-(\vec{c}.\vec{a})\vec{a}=\vec{a}\times\vec{b}$ $\Rightarrow 2\vec{c} - 4\vec{a} = \vec{a} \times \vec{b}$ Now $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 0 \\ 1 & 1 & 1 \end{vmatrix} = -\hat{i} - \hat{j} + 2\hat{k}$ So, $2\vec{c} = 4\hat{i} - 4\hat{i} - \hat{i} - \hat{i} + 2\hat{k}$ $=3\hat{i}-5\hat{i}+2\hat{k}$ $\Rightarrow \vec{c} = \frac{3}{2}\hat{i} - \frac{5}{2}\hat{j} + \hat{k}$ $\left|\vec{c}\right| = \sqrt{\frac{9}{4} + \frac{25}{4} + 1} = \sqrt{\frac{38}{4}} = \sqrt{\frac{19}{2}}$ $|\vec{c}|^2 = \frac{19}{2}$

16. Let a_1, a_2, \dots, a_{30} be an A. P., $S = \sum_{i=1}^{30} a_i$ and $T = \sum_{i=1}^{15} a_{(2i-1)}$. If $a_5 = 27$ and S - 2T = 75, then a_{10} is equal to : (3) 42 (1) 57 (2) 47 (4) 52 Ans. (4) **Sol.** $S = a_1 + a_2 + \dots + a_{30}$ $S = \frac{30}{2} [a_1 + a_{30}]$ $S = \overline{15}(a_1 + a_{30}) = 15 (a_1 + a_1 + 29d)$ $T = a_1 + a_3 + \dots + a_{29}$ $= (a_1) + (a_1 + 2d) \dots + (a_1 + 28d)$ $= 15a_1 + 2d(1 + 2 + \dots + 14)$ $T = 15a_1 + 210 d$ Now use S - 2T = 75 \Rightarrow 15 (2a₁ + 29d) - 2 (15a₁ + 210 d) = 75 \Rightarrow d = 5 Given $a_5 = 27 = a_1 + 4d \Rightarrow a_1 = 7$ Now $a_{10} = a_1 + 9d = 7 + 9 \times 5 = 52$ 17. 5 students of a class have an average height 150 cm and variance 18 cm². A new student, whose height is 156 cm, joined them. The variance (in cm²) of the height of these six students is: (1) 22(2) 20(3) 16 (4) 18 Ans. (2) **Sol.** Given $\vec{x} = \frac{\Sigma x_i}{5} = 150$ $\Rightarrow \sum_{i=1}^{5} x_i = 750$(i) $\frac{\Sigma x_i^2}{5} - \left(\vec{x}\right)^2 = 18$ $\frac{\Sigma x_i^2}{5} - (150)^2 = 18$

$$\Sigma x_i^2 = 112590$$
(ii)

Given height of new student $x_6 = 156$

Now,
$$\vec{x}_{new} = \frac{\sum_{i=1}^{6} x_i}{6} = \frac{750 + 156}{6} = 151$$

Also, New variance
$$=\frac{\sum_{i=1}^{3} x_i^2}{6} - (\overline{x}_{new})^2$$

$$=\frac{112590 + (156)^2}{6} - (151)^2$$
$$= 22821 - 22801 = 20$$

4



18. Two cards are drawn successively with 20. Let replacement from a well-shuffled deck of 52 cards. Let X denote the random variable of number of aces obtained in the two drawn cards. Then P(X = 1) + P(X = 2) equals : (1) 52/169(2) 25/169(3) 49/169 (4) 24/169 Ans. (2) Sol. Two cards are drawn successively with replacement 4 Aces 48 Non Aces Ans. (2) $P(x=1) = \frac{{}^{4}C_{1}}{{}^{52}C_{1}} \times \frac{48C_{1}}{52C_{1}} + \frac{48C_{1}}{52C_{1}} \times \frac{4C_{1}}{52C_{1}} = \frac{24}{169}$ $P(x=2) = \frac{{}^{4}C_{1}}{{}^{52}C_{1}} \times \frac{{}^{4}C_{1}}{{}^{52}C_{2}} = \frac{1}{169}$ $P(x = 1) + P(x = 2) = \frac{25}{169}$ For $x \in R - \{0, 1\}$, let $f_1(x) = \frac{1}{x}$, 19. $f_2(x) = 1 - x$ and $f_3(x) = \frac{1}{1 - x}$ be three given functions. If a function, J(x) satisfies $(f_2 \circ J \circ f_1)(x) = f_3(x)$ then J(x) is equal to :-(1) $f_3(x)$ (2) $f_1(x)$ (4) $\frac{1}{x} f_3(x)$ (3) $f_2(x)$ Ans. (1) **Sol.** Given $f_1(x) = \frac{1}{x}$, $f_2(x) = 1 - x$ and $f_3(x) = \frac{1}{1 - x}$ $(f_2 \circ J \circ f_1)(x) = f_3(x)$ 21. $f_{2} \circ \left(J \left(f_{1} \left(x \right) \right) \right) = f_{3} \left(x \right)$ $f_2 \circ \left(J\left(\frac{1}{x}\right)\right) = \frac{1}{1-x}$ $1-J\left(\frac{1}{x}\right)=\frac{1}{1-x}$ Ans. (4) $J\left(\frac{1}{x}\right) = 1 - \frac{1}{1-x} = \frac{-x}{1-x} = \frac{x}{x-1}$ Now $x \to \frac{1}{x}$ $J(x) = \frac{\frac{1}{x}}{\frac{1}{x-1}} = \frac{1}{1-x} = f_3(x)$

$$A = \left\{ 0 \in \left(-\frac{\pi}{2}, \pi \right) : \frac{3 + 2i\sin\theta}{1 - 2i\sin\theta} \text{ is purely imaginary} \right\}$$

Then the sum of the elements in A is :

(1)
$$\frac{5\pi}{6}$$
 (2) $\frac{2\pi}{3}$
(3) $\frac{3\pi}{4}$ (4) π

Sol. Given
$$z = \frac{3 + 2i\sin\theta}{1 - 2i\sin\theta}$$
 is purely img

$$z = \left(\frac{3 + 2i\sin\theta}{1 - 2i\sin\theta}\right) \times \left(\frac{1 + 2i\sin\theta}{1 + 2i\sin\theta}\right)$$
$$z = \frac{(3 - 4\sin^2\theta) + i(8\sin\theta)}{i + 4\sin^2\theta}$$
Now Re(z) = 0
$$\frac{3 - 4\sin^2\theta}{1 + 4\sin^2\theta} = 0$$
$$\sin^2\theta = \frac{3}{4}$$
$$\sin\theta = \pm \frac{\sqrt{3}}{2} \implies \theta = -\frac{\pi}{3}, \frac{\pi}{3}, \frac{2\pi}{3}$$
$$\because \theta \in \left(-\frac{\pi}{2}, \pi\right)$$

then sum of the elements in A is

$$-\frac{\pi}{3} + \frac{\pi}{3} + \frac{2\pi}{3} = \frac{2\pi}{3}$$

21. If θ denotes the acute angle between the curves, y = 10 - x² and y = 2 + x² at a point of their intersection, then $|\tan \theta|$ is equal to :

Sol. Point of intersection is P(2,6).

Also,
$$m_1 = \left(\frac{dy}{dx}\right)_{P(2,6)} = -2x = -4$$

 $m_2 = \left(\frac{dy}{dx}\right)_{P(2,6)} = 2x = 4$
 $\therefore |\tan \theta| = \left|\frac{m_1 - m_2}{1 + m_1 m_2}\right| = \frac{8}{15}$

collegebatch.com

Ε



JEE (Main) Examination-2019/Morning Session/09-01-2019

22. If $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$, then the matrix A^{-50} when $\theta = \frac{\pi}{12}$, is equal to : (1) $\begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$ (2) $\begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}$ (2) $\begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}$ (3) $\begin{bmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}$ (4) $\begin{bmatrix} \frac{\sqrt{3}}{2} & -\frac{1}{2} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$ (5) Ans. (2) Sol. $e = \sqrt{1 + \tan^2 \theta} = \sec \theta$ As, $\sec \theta > 2 \Rightarrow \cos \theta$ $\Rightarrow \theta \in (60^\circ, 90^\circ)$ Now, $\ell(L \cdot R) = \frac{2b^2}{a} = 2(\sec \theta - \cos \theta)$ Which is strictly incr ℓ (L.R) $\in (3, \infty)$. 24. The equation of th (-4, 3, 1), parallel to th and intersecting the line (1) $\frac{x+4}{-1} = \frac{y-3}{1} = \frac{z-1}{12}$

Ans. (1)

Sol. Here, $AA^T = I$

$$\Rightarrow A^{-1} = A^{T} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$
Also, $A^{-n} = \begin{bmatrix} \cos(n\theta) & \sin(n\theta) \\ -\sin(n\theta) & \cos(n\theta) \end{bmatrix}$

$$\therefore A^{-50} = \begin{bmatrix} \cos(50)\theta & \sin(50)\theta \\ -\sin(50)\theta & \cos(50)\theta \end{bmatrix}$$

$$= \begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$$
Let $0 < \theta < \frac{\pi}{2}$. If the eccentric

23. Let $0 < \theta < \frac{\pi}{2}$. If the eccentricity of the hyperbola $\frac{x^2}{\cos^2 \theta} - \frac{y^2}{\sin^2 \theta} = 1$ is greater than 2, then the length of its latus rectum lies in the interval :

(1) (2, 3]	(2) (3, ∞)
(3) (3/2, 2]	(4) (1, 3/2]

Ans. (2) As, sec $\theta > 2 \Rightarrow \cos \theta < \frac{1}{2}$ $\Rightarrow \theta \in (60^{\circ}, 90^{\circ})$ Now, $\ell(L \cdot R) = \frac{2b^2}{a} = 2\frac{(1 - \cos^2 \theta)}{\cos^2 \theta}$ =2(sec θ - cos θ) Which is strictly increasing, so ℓ (L.R) $\in (3,\infty)$. The equation of the line passing through 24. (-4, 3, 1), parallel to the plane x + 2y - z - 5 = 0and intersecting the line $\frac{x+1}{-3} = \frac{y-3}{2} = \frac{z-2}{-1}$ is: (1) $\frac{x+4}{-1} = \frac{y-3}{1} = \frac{z-1}{1}$ (2) $\frac{x+4}{3} = \frac{y-3}{-1} = \frac{z-1}{1}$ (3) $\frac{x+4}{1} = \frac{y-3}{1} = \frac{z-1}{3}$ (4) $\frac{x-4}{2} = \frac{y+3}{1} = \frac{z+1}{4}$ Ans. (2) Sol.

Normal vector of plane containing two intersecting lines is parallel to vector.

$$\left(\vec{\mathbf{V}}_{1}\right) = \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ \mathbf{3} & \mathbf{0} & \mathbf{1} \\ -\mathbf{3} & \mathbf{2} & -\mathbf{1} \end{vmatrix}$$

 $= -2\hat{i} + 6\hat{k}$ ∴ Required line is parallel to vector

$$\left[\vec{V}_{2}\right) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & -1 \\ -2 & 0 & 6 \end{vmatrix} = 3\hat{i} - \hat{j} + \hat{k}$$

 \Rightarrow Required equation of line is

$$\frac{x+4}{3} = \frac{y-3}{-1} = \frac{z-1}{1}$$

6

collegebatch.com

For any $\theta \in \left(\frac{\pi}{4}, \frac{\pi}{2}\right)$, the expression 25. $3(\sin\theta - \cos\theta)^4 + 6(\sin\theta + \cos\theta)^2 + 4\sin^6\theta$ equals : (1) $13 - 4 \cos^6\theta$ (2) $13 - 4 \cos^4\theta + 2 \sin^2\theta\cos^2\theta$ (3) $13 - 4 \cos^2\theta + 6 \cos^4\theta$ (4) $13 - 4 \cos^2\theta + 6 \sin^2\theta\cos^2\theta$ Ans. (1) Sol. We have, $3(\sin \theta - \cos \theta)^4 + 6(\sin \theta + \cos \theta)^2 + 4 \sin^6 \theta$ $= 3(1 - \sin 2\theta)^2 + 6(1 + \sin 2\theta) + 4\sin^6\theta$ $= 3(1 - 2\sin 2\theta + \sin^2 2\theta) + 6 + 6\sin 2\theta + 4\sin^6\theta$ $= 9 + 12 \sin^2\theta \cdot \cos^2\theta + 4(1 - \cos^2\theta)^3$ $= 13 - 4 \cos^6\theta$ If $\cos^{-1}\left(\frac{2}{3x}\right) + \cos^{-1}\left(\frac{3}{4x}\right) = \frac{\pi}{2}\left(x > \frac{3}{4}\right)$ 26. then x is equal to (2) $\frac{\sqrt{145}}{10}$ (1) $\frac{\sqrt{145}}{12}$ (3) $\frac{\sqrt{146}}{12}$ (4) $\frac{\sqrt{145}}{11}$ Ans. (1) **Sol.** $\cos^{-1}\left(\frac{2}{3x}\right) + \cos^{-1}\left(\frac{3}{4x}\right) = \frac{\pi}{2} \left(x > \frac{3}{4}\right)$ $\cos^{-1}\left(\frac{3}{4x}\right) = \frac{\pi}{2} - \cos^{-1}\left(\frac{2}{3x}\right)$ $\cos^{-1}\left(\frac{3}{4\mathbf{x}}\right) = \sin^{-1}\left(\frac{2}{3\mathbf{x}}\right)$ $\cos\left(\cos^{-1}\left(\frac{3}{4x}\right)\right) = \cos\left(\sin^{-1}\frac{2}{3x}\right)$ $\frac{3}{4x} = \frac{\sqrt{9x^2 - 4}}{3x}$ $\frac{81}{16} + 4 = 9x^2$ $x^2 = \frac{145}{16 \times 9} \implies x = \frac{\sqrt{145}}{12}$

27. The value of
$$\int_{0}^{\pi} |\cos x|^{3} dx$$

(1) 2/3 (2) 0
(3) -4/3 (4) 4/3
Ans. (4)
Sol. $\int_{0}^{\pi} |\cos x|^{3} dx = \int_{0}^{\pi/2} \cos^{3} x dx - \int_{\pi/2}^{\pi} \cos^{3} x dx$
 $= \int_{0}^{\pi/2} \left(\frac{\cos 3x + 3\cos x}{4} \right) dx - \int_{\pi/2}^{\pi} \left(\frac{\cos 3x + 3\cos x}{4} \right) dx$
 $= \frac{1}{4} \left[\left(\frac{\sin 3x}{3} + 3\sin x \right)_{0}^{\pi/2} - \left(\frac{\sin 3x}{3} + 3\sin x \right)_{\pi/2}^{\pi} \right]$
 $= \frac{1}{4} \left[\left(\frac{-1}{3} + 3 \right) - (0 + 0) - \left\{ (0 + 0) - \left(\frac{-1}{3} + 3 \right) \right\} \right]$
 $= \frac{4}{3}$
28. If the Boolean expression
(p \oplus q) ^ (~p \odot q) is equivalent to p ^ q, where
 \oplus , $\odot \in \{\wedge, \vee\}$, then the ordered pair (\oplus , \odot) is

- (1) (\wedge, \vee) (2) (\vee, \vee)
- (3) (\wedge, \wedge)
- (4) (∨,∧)

Ans. (1)

Sol. $(p \oplus q) \land (\sim p \square q) \equiv p \land q$ (given)

р	q	$\sim p$	$p \land q$	$p \lor q$	$\sim p \lor q$	$\sim p \wedge q$	$(p \land q) \land (\sim p \lor q)$
Т	Т	F	Т	Т	Т	F	Т
Т	F	F	F	Т	F	F	F
F	Т	Т	F	Т	Т	Т	F
F	F	Т	F	F	Т	F	F

from truth table $(\oplus, \square) = (\land, \lor)$

7



+1

29.
$$\lim_{y \to 0} \frac{\sqrt{1 + \sqrt{1 + y^4}} - \sqrt{2}}{y^4}$$
(1) exists and equals $\frac{1}{4\sqrt{2}}$
(2) does not exist
(3) exists and equals $\frac{1}{2\sqrt{2}}$
(4) exists and equals $\frac{1}{2\sqrt{2}}(\sqrt{2})$

Ans. (1)

Sol.
$$\lim_{y \to 0} \frac{\sqrt{1 + \sqrt{1 + y^4}} - \sqrt{2}}{y^4}$$

$$= \lim_{y \to 0} \frac{1 + \sqrt{1 + y^4} - 2}{y^4 \left(\sqrt{1 + \sqrt{1 + y^4}} + \sqrt{2}\right)}$$
$$= \lim_{y \to 0} \frac{\left(\sqrt{1 + y^4} - 1\right)\left(\sqrt{1 + y^4} + 1\right)}{y^4 \left(\sqrt{1 + \sqrt{1 + y^4}} + \sqrt{2}\right)\left(\sqrt{1 + y^4} + 1\right)}$$

$$= \lim_{y \to 0} \frac{1 + y^4 - 1}{y^4 \left(\sqrt{1 + \sqrt{1 + y^4}} + \sqrt{2}\right) \left(\sqrt{1 + y^4} + 1\right)}$$
$$= \lim_{x \to 0} \frac{1}{\sqrt{1 + y^4}} = \frac{1}{\sqrt{1 + y^4}}$$

$$= \lim_{y \to 0} \frac{1}{\left(\sqrt{1 + \sqrt{1 + y^4}} + \sqrt{2}\right)\left(\sqrt{1 + y^4} + 1\right)} = \frac{1}{4\sqrt{2}}$$

30. Let $f : R \to R$ be a function defined as :

$$f(x) = \begin{cases} 5, & \text{if} \quad x \le 1 \\ a + bx, & \text{if} \quad 1 < x < 3 \\ b + 5x, & \text{if} \quad 3 \le x < 5 \\ 30, & \text{if} \quad x \ge 5 \end{cases}$$

Then, f is :

(1) continuous if a = 5 and b = 5

(2) continuous if a = -5 and b = 10

(3) continuous if a = 0 and b = 5

(4) not continuous for any values of a and b

Ans. (4)

Sol.
$$f(x) = \begin{cases} 5 & \text{if } x \le 1 \\ a + bx & \text{if } 1 < x < 3 \\ b + 5x & \text{if } 3 \le x < 5 \\ 30 & \text{if } x \ge 5 \end{cases}$$

$$f(1) = 5, \quad f(1^{-}) = 5, \quad f(1^{+}) = a + b$$

$$f(3^{-}) = a + 3b, \quad f(3) = b + 15, \quad f(3^{+}) = b + 15$$

$$f(5^{-}) = b + 25; \quad f(5) = 30 \qquad f(5^{+}) = 30$$

from above we concluded that f is not continuous for any values of a and b.





TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Wednesday 09th JANUARY, 2019) TIME : 9 : 30 AM To 12 : 30 PM PHYSICS

1. A current loop, having two circular arcs joined by two radial lines is shown in the figure. It carries a current of 10 A. The magnetic field at point O will be close to :



	(1) 1.0×10^{-5} T	(2) $1.5 \times 10^{-5} \text{ T}$
	(3) $1.0 \times 10^{-7} \text{ T}$	(4) $1.0 \times 10^{-7} \text{ T}$
Ans.	(1)	



$$\vec{B} = \frac{\mu_0 i}{4\pi} \theta \left[\frac{1}{r_1} - \frac{1}{r_2} \right] \hat{k}$$

$$r_1 = 3 \text{ cm} = 3 \times 10^{-2} \text{ m}$$

$$r_2 = 5 \text{ cm} = 5 \times 10^{-2} \text{ m}$$

$$\theta = \frac{\pi}{4}, \text{ i} = 10 \text{ A}$$

$$\Rightarrow \vec{B} = \frac{4\pi \times 10^{-7}}{16} \times 10 \left[\frac{1}{3 \times 10^{-2}} - \frac{1}{5 \times 10^{-2}} \right] \hat{k}$$

$$\Rightarrow \left| \vec{B} \right| = \frac{\pi}{3} \times 10^{-5} \text{ T}$$

$$\approx 1 \times 10^{-5} \text{ T}$$

2. A gas can be taken from A to B via two different processes ACB and ADB.



When path ACB is used 60 J of heat flows into the system and 30 J of work is done by the system. If path ADB is used work done by the system is 10 J. The heat Flow into the system in path ADB is : (1) 80 J (2) 20 J (3) 100 J (4) 40 J

Ans. (4) P C B C A D V $\Delta Q_{ACB} = \Delta W_{ACB} + \Delta U_{ACB}$ $\Rightarrow 60 \text{ J} = 30 \text{ J} + \Delta U_{ACB}$ $\Rightarrow \Delta U_{ACB} = 30 \text{ J}$ $\Rightarrow \Delta U_{ACB} = \Delta U_{ACB} = 30 \text{ J}$ $\Delta Q_{ACD} = \Delta U_{ACB} + \Delta W_{ADB}$ = 10 J + 30 J = 40 J

3. A plane electromagnetic wave of frequency 50 MHz travels in free space along the positive x-direction. At a particular point in space and time, $\vec{E} = 6.3\hat{j}V/m$. The corresponding magnetic field \vec{B} , at that point will be:

(1)
$$18.9 \times 10^{-8} \hat{k} T$$
 (2) $6.3 \times 10^{-8} \hat{k} T$

(3)
$$2.1 \times 10^{-8} \hat{k} T$$
 (4) $18.9 \times 10^{8} \hat{k} T$

Ans. (3)

Sol.
$$|B| = \frac{|E|}{C} = \frac{6.3}{3 \times 10^8} = 2.1 \times 10^{-8} \text{ T}$$

and $\hat{E} \times \hat{B} = \hat{C}$
 $\hat{j} \times \hat{B} = \hat{i}$
 $\hat{B} = \hat{k}$
 $\vec{B} = |B|\hat{B} = 2.1 \times 10^8 \hat{k} \text{ T}$

1



E

JEE (Main) Examination-2019/Morning Session/09-01-2019

- **4.** Two coherent sources produce waves of different intensities which interfere. After interference, the ratio of the maximum intensity to the minimum intensity is 16. The intensity of the waves are in the ratio:
 - (1) 4 : 1(2) 25 : 9(3) 16 : 9(4) 5 : 3
- Ans. (2)

Sol. $\frac{I_{max}}{I_{min}} = 16$

$$\Rightarrow \frac{A_{max}}{A_{min}} = 4$$

$$\Rightarrow \frac{\mathbf{A}_1 + \mathbf{A}_2}{\mathbf{A}_1 - \mathbf{A}_2} = \frac{4}{1}$$

Using componendo & dividendo.

$$\frac{A_1}{A_2} = \frac{5}{3} \Longrightarrow \frac{I_1}{I_2} = \left(\frac{5}{3}\right)^2 = \frac{25}{9}$$

5. An L-shaped object, made of thin rods of uniform mass density, is suspended with a string as shown in figure. If AB = BC, and the angle made by AB with downward vertical is θ , then :





(4)
$$\tan \theta = \frac{1}{2\sqrt{3}}$$

Ans. (2) Sol. $C_1 \qquad \theta \qquad P$ $M \qquad mg$

> Let mass of one rod is m. Balancing torque about hinge point. mg (C₁P) = mg (C₂N)

$$mg\left(\frac{L}{2}\sin\theta\right) = mg\left(\frac{L}{2}\cos\theta - L\sin\theta\right)$$

$$\Rightarrow \frac{3}{2} \text{mgL} \sin \theta = \frac{\text{mgL}}{2} \cos \theta$$
$$\Rightarrow \tan \theta = \frac{1}{3}$$

6. A mixture of 2 moles of helium gas (atomic mass = 4 u), and 1 mole of argon gas (atomic mass = 40 u) is kept at 300 K in a container. The ratio

of their rms speeds
$$\left[\frac{V_{rms}(helium)}{V_{rms}((argon)}\right]$$
, is close to

$$\begin{array}{c} \cdot \\ (1) \ 2.24 \\ (3) \ 0.32 \end{array} \qquad \begin{array}{c} (2) \ 0.45 \\ (4) \ 3.16 \end{array}$$

Ans. (4)

Sol.
$$\frac{V_{rms}(He)}{V_{rms}(Ar)} = \sqrt{\frac{M_{Ar}}{M_{He}}} = \sqrt{\frac{40}{4}} = 3.16$$

7. When the switch S, in the circuit shown, is closed, then the value of current *i* will be :

$$\begin{array}{c} 20V i_1 & C & \downarrow i_2 10V \\ A & 2\Omega & \downarrow i & 4\Omega & B \\ & & & & \\ & & & \\ & & & & & \\ & & & & \\ & & & & \\ & & & & & \\ & & & & \\ & & & & \\ & & & &$$

2



Ans. (2)

Sol.
$$20V \stackrel{i_1}{A} \xrightarrow{XV} \xrightarrow{V} 4\Omega \xrightarrow{i_2} 10V$$

$$2\Omega \xrightarrow{i_2} 2\Omega$$

$$2\Omega \xrightarrow{i_3} 0$$
Let voltage at C = xv
KCL : $i_1 + i_2 = i$

$$\frac{20 - x}{2} + \frac{10 - x}{4} = \frac{x - 0}{2}$$

$$\Rightarrow x = 10$$
and $i = 5$ amp.

8. A resistance is shown in the figure. Its value and tolerance are given respectively by:

- (1) 27 KΩ, 20%
 (2) 270 KΩ, 5%
 (3) 270 KΩ, 10%
 (4) 27 KΩ, 10%
- Ans. (4)
- Sol. Color code :

Red violet orange silver $R = 27 \times 10^3 \Omega \pm 10\%$ $= 27 K\Omega \pm 10\%$

9. A bar magnet is demagnetized by inserting it inside a solenoid of length 0.2 m, 100 turas, and carrying a current of 5.2 A. The coercivity of the bar magnet is :

(1) 1200 A/m	(2) 2600 A/m
(3) 520 A/jm	(4) 285 A/m

Ans. (2)

E

Sol. Coercivity =
$$H = \frac{B}{\mu_0}$$

= ni = $\frac{N}{\ell}i$ = $\frac{100}{0.2} \times 5.2$
= 2600 A/m

10. A rod, of length L at room temperature and uniform area of cross section A, is made of a metal having coefficient of linear expansion α/ °C. It is observed that an external compressive force F, is applied on

each of its ends, prevents any change in the length of the rod, when its temperature rises by ΔT K. Young's modulus, Y, for this metal is :

(1)
$$\frac{F}{2A\alpha\Delta T}$$
 (2) $\frac{F}{A\alpha(\Delta T - 273)}$

(3)
$$\frac{F}{A\alpha\Delta T}$$
 (4) $\frac{2F}{A\alpha\Delta T}$

Ans. (3)

Sol. Young's modulus
$$y = \frac{\text{Stress}}{\text{Strain}}$$

$$=\frac{F/A}{\left(\Delta\ell/\ell\right)}$$
$$=\frac{F}{A(\alpha\Delta T)}$$

11. A block of mass m, lying on a smooth horizontal surface, is attached to a spring (of negligible mass) of spring constant k. The other end of the spring is fixed, as shown in the figure. The block is initially at rest in its equilibrium position. If now the block is pulled with a constant force F, the maximum speed of the block is :





JEE (Main) Examination-2019/Morning Session/09-01-2019

Ans. (3)

Sol. Maximum speed is at mean position (equilibrium). F = kx

$$x = \frac{F}{k}$$

$$W_F + W_{sp} = \Delta KE$$

$$F(x) - \frac{1}{2}kx^2 = \frac{1}{2}mv^2 - 0$$

$$F\left(\frac{F}{k}\right) - \frac{1}{2}k\left(\frac{F}{k}\right)^2 = \frac{1}{2}mv^2$$

$$\Rightarrow v_{max} = \frac{F}{\sqrt{mk}}$$

12. Three charges +Q, q, +Q are placed respectively, at distance, 0, d/2 and d from the origin, on the x-axis. If the net force experienced by + Q, placed at x= 0, Ls zero, then value of q is :

(1) + Q/2(2) - Q/2(3) - Q/4(4) + O/4Ans. (3)

Sol.
$$F_{a} \xleftarrow{F_{b}}{+Q} \xrightarrow{F_{b}}{d/2} q \xrightarrow{d/2} +Q$$

For equilibrium,
 $\vec{F}_{a} + \vec{F}_{B} = 0$
 $\vec{F}_{a} = -\vec{F}_{B}$
 $\frac{kQQ}{d^{2}} = -\frac{kQq}{(d/2)^{2}}$
 $\Rightarrow q = -\frac{Q}{4}$

$$\Rightarrow$$
 q = -

13. A conducting circular loop made of a thill wire, has area $3.5 \times 10^{-3} \text{ m}^2$ and resistance 10Ω . It is placed perpendicular to a time dependent magnetic field $B(t) = (0.4T)\sin(50\pi t)$. The field is uniform in

space. Then the net charge flowing through the loop during t = 0 s and t = 10 ms is close to: (1) 14mC (2) 21 mC (3) 6 mC (4) 7 mC Ans. (1)

Sol.
$$Q = \frac{\Delta \phi}{R} = \frac{1}{10} A (B_f - B_i) = \frac{1}{10} \times 3.5 \times 10^{-3} \left(0.4 \sin \frac{\pi}{2} - 0 \right)$$

 $= \frac{1}{10} \left(3.5 \times 10^{-3} \right) \left(0.4 - 0 \right)$
 $= 1.4 \times 10^{-4} = 0.14 \text{ mC}$

Two masses m and $\frac{m}{2}$ are connected at the two 14. ends of a massless rigid rod of length l. The rod is suspended by a thin wire of torsional constant k at the centre of mass of the rod-mass system(see figure). Because of torsional constant k, the restoring torque is $\tau = k\theta$ for angular displacement 0. If the rod is rota ted by θ_0 and released, the tension in it when it passes through its mean position will be:



(1)
$$\frac{3K0_0}{l}$$
 (2) $\frac{K0_0}{2l}$

(3)
$$\frac{2k\theta_0^2}{l}$$
 (4) $\frac{k\theta_0^2}{l}$

Ans. (4)

Sol.
$$\omega = \sqrt{\frac{k}{I}}$$

 $\omega = \sqrt{\frac{3k}{m\ell^2}}$
 $m/2$ T m
 ℓ

collegebatch.com



15. A copper wire is stretched to make it 0.5% longer. The percentage change in its electrical resistance if its volume remains unchanged is: (1) 2.5% (2) 0.5% (3) 1.0% (4) 2.0%
Ans. (3)

Sol.
$$R = \frac{\rho \ell}{A}$$
 and volume (V) = $A\ell$.
 $R = \frac{\rho \ell^2}{V}$
 $\Rightarrow \frac{\Delta R}{R} = \frac{2\Delta \ell}{\ell} = 1\%$

16. A parallel plate capacitor is made of two square plates of side 'a', separated by a distance d (d<<a). The lower triangular portion is filled with a dielectric of dielectric constant K, as shown in the figure.</p>

Capacitance of this capacitor is :

collegebatch.com

Ε

(1)
$$\frac{1}{2} \frac{k \in_0 a^2}{d}$$
 (2) $\frac{k \in_0 a^2}{d} \ln K$
(3) $\frac{k \in_0 a^2}{d(K-1)} \ln K$ (4) $\frac{k \in_0 a^2}{2d(K+1)}$



5

JEE (Main) Examination-2019/Morning Session/09-01-2019

17. Mobility of electrons in a semiconductor is defined as the ratio of their drift velocity to the applied electric field. If, for an n-type semiconductor, the density of electrons is 10^{19} m⁻³ and their mobility is 1.6 m²/(V.s) then the resistivity of the semiconductor (since it is an n-type semiconductor

contribution of holes is ignored) is close to:

- (1) $2\Omega m$ (2) $0.4\Omega m$
- (3) $4\Omega m$ (4) $0.2\Omega m$
- Ans. (2)

Sol. $j = \sigma E = nev_d$

$$\sigma = ne \frac{v_0}{E}$$

$$\frac{1}{\sigma} = \rho = \frac{1}{n_e e \mu_e}$$

$$=\frac{1}{10^{19}\times1.6\times10^{-19}\times1.6}$$

- $= 0.4 \Omega m$
- 18. If the angular momentum of a planet of mass m, moving around the Sun in a circular orbit is L, about the center of the Sun, its areal velocity is :

(1)
$$\frac{4L}{m}$$
 (2) $\frac{L}{m}$ (3) $\frac{L}{2m}$ (4) $\frac{2L}{m}$

Ans. (3)

- **Sol.** $\frac{dA}{dt} = \frac{L}{2m}$
- **19.** A block of mass 10 kg is kept on a rough inclined plane as shown in the figure. A force of 3 N is applied on the block. The coefficient of static friction between the plane and the block is 0.6. What should be the minimum value of force P, such that the block doesnot move downward ?







20. Temperature difference of 120°C is maintained between two ends of a uniform rod AB of length 2L. Another bent rod PQ, of same cross-

section as AB and length $\frac{3L}{2}$, is connected across AB (See figure). In steady state, temperature difference between P and Q will be close to :



(1) 60°C (2) 75°C (3) 35°C (4) 45°C Ans. (4)



6

21. A heavy ball of mass M is suspended from the ceiling of a car by a light string of mass m (m<<M). When the car is at rest, the speed of transverse waves in the string is 60 ms⁻¹. When the car has acceleration a, the wave-speed increases to 60.5 ms⁻¹. The value of a, in terms of gravitational acceleration g, is closest to :

(1)
$$\frac{g}{5}$$
 (2) $\frac{g}{20}$

(3)
$$\frac{g}{10}$$
 (4) $\frac{g}{30}$

Ans. (1)

Sol.
$$60 = \sqrt{\frac{Mg}{\mu}}$$

$$60.5 = \sqrt{\frac{M(g^2 + a^2)^{1/2}}{\mu}} \Rightarrow \frac{60.5}{60} = \sqrt{\sqrt{\frac{g^2 + a^2}{g^2}}}$$
$$\left(1 + \frac{0.5}{60}\right)^4 = \frac{g^2 + a^2}{g^2} = 1 + \frac{2}{60}$$
$$\Rightarrow g^2 + a^2 = g^2 + g^2 \times \frac{2}{60}$$
$$a = g\sqrt{\frac{2}{60}} = \frac{g}{\sqrt{30}} = \frac{g}{5.47}$$
$$\approx \frac{g}{5}$$

22. A sample of radioactive material A, that has an activity of 10 mCi(1 Ci = 3.7×10^{10} decays/s), has twice the number of nuclei as another sample of a different radioactive maternal B which has an activity of 20 mCi. The correct choices for hall-lives of A and B would then be respectively :

(1) 20 days and 5 days (2) 20 days and 10 days (3) 5 days and 10 days (4) 10 days and 40 days

Ans. (1)

E

Sol. Activity $A = \lambda N$ For A 10 = $(2N_0)\lambda_A$ For B 20 = $N_0\lambda_B$ $\therefore \lambda_B = 4\lambda_A \Rightarrow (T_{1/2})_A = 4(T_{1/2})_B$ 23. Consider a tank made of glass(reiractive index 1.5) with a thick bottom. It is filled with a liquid of refractive index μ . A student finds that, irrespective of what the incident angle *i* (see figure) is for a beam of light entering the liquid, the light reflected from the liquid glass interface is never completely polarized. For this to happen, the minimum value of μ is :

(1)
$$\frac{3}{\sqrt{5}}$$
 (2) $\frac{5}{\sqrt{3}}$ (3) $\sqrt{\frac{5}{3}}$ (4) $\frac{4}{3}$

Ans. (1) **Sol.** $C < i_{b}$

here i_b is "brewester angle" and c is critical angle

$$\sin_{c} < \sin i_{b} \qquad \text{since } \tan i_{b} = \mu_{0_{rel}} = \frac{1.5}{\mu}$$

$$\frac{1}{\mu} < \frac{1.5}{\sqrt{\mu^{2} + (1.5)^{2}}} \qquad \therefore \sin i_{b} = \frac{1.5}{\sqrt{\mu^{2} + (1.5)^{2}}}$$

$$\sqrt{\mu^{2} \times (1.5)^{2}} < 1.5 \times \mu$$

$$\mu^{2} + (1.5)^{2} < (\mu \times 1.5)^{2}$$

$$\mu < \frac{3}{\sqrt{5}}$$

slab $\mu = 1.5$

24. An infinitely long current carrying wire and a small current carrying loop are in the plane of the paper as shown. The radius of the loop is a and distance of its centre from the wire is d (d»a). If the loop applies a force F on the wire then :





Ans. (3)

Sol.
Sol.

$$\infty \text{ long wire}$$

Eqvilent dipole of given loop
 $F = m \cdot \frac{dB}{dr}$
Now $\frac{dB}{dx} = \frac{d}{dx} \left(\frac{\mu_0 I}{2\pi x} \right)$
 $\propto \frac{1}{x^2}$
 $\Rightarrow \text{ So } F \propto \frac{M}{x^2} [\because M = \text{NIA}]$

$$\therefore F \propto \frac{a^2}{d^2}$$

25. Surface of certain metal is first illuminated with light of wavelength λ_1 =350 nm and then, by light of wavelength λ_2 =54D nm. It is found that the maximum speed of the photo electrons in the two cases differ by a factor of 2. The work function of the metal (in eV) is close to :

(Energir of photon = $\frac{1240}{\lambda(\text{in nm})}$ eV) (1) 1.8 (2) 1.4 (3) 2.5 (4) 5.6

Ans. (1)

Sol.
$$\frac{hc}{\lambda_{1}} = \phi + \frac{1}{2}m(2v)^{2}$$
$$\frac{hc}{\lambda_{2}} = \phi + \frac{1}{2}mv^{2}$$
$$\Rightarrow \frac{hc}{\lambda_{1}} - \phi$$
$$\Rightarrow \frac{hc}{\lambda_{2}} - \phi = 4 \Rightarrow \frac{hc}{\lambda_{1}} - \phi = \frac{4hc}{\lambda_{2}} - 4\phi$$
$$\Rightarrow \frac{4hc}{\lambda_{2}} - \frac{hc}{\lambda_{1}} = 3\phi$$
$$\Rightarrow \phi = \frac{1}{3}hc\left(\frac{4}{\lambda_{2}} - \frac{1}{\lambda_{1}}\right)$$
$$= \frac{1}{3} \times 1240\left(\frac{4 \times 350 - 540}{350 \times 540}\right)$$
$$= 1.8 \text{ eV}$$

26. A particle is moving with a velocity $\overline{v} = K(y\hat{i} + x\hat{j})$, where K is a constant. The general equation for its path is:

(1) xy = constant (2) $y^2 = x^2 + constant$ (3) $y = x^2 + constant$ (4) $y^2 = x + constant$ **Ans. (2)**

Sol.
$$\frac{dx}{dt} = ky, \frac{dy}{dt} = kx$$

Now, $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{x}{y}$
 $\Rightarrow ydy = xdx$
Integrating both side
 $y^2 = x^2 + c$

27. A convex lens is put 10 cm from a light source and it makes a sharp image on a screen, kept 10 cm from the lens. Now a glass block (refractive index 1.5) of 1.5 cm thickness is placed in contact with the light source. To get the sharp image again, the screen is shifted by a distance d. Then d is :

(1) 0.55 cm away from the lens

(2) 1.1 cm away from the lens

(3) 0.55 cm towards the lens

(4) 0 Ans. (1)





28. For a uniformly charged ring of radius R, the electric field on its axis has the largest magnitude at a distance h from its centre. Then value of h is :

(1)
$$\frac{R}{\sqrt{5}}$$
 (2) R (3) $\frac{R}{\sqrt{2}}$ (4) $R\sqrt{2}$

Ans. (3)

Sol. Electric field on axis of ring

$$\mathbf{E} = \frac{\mathbf{k}\mathbf{Q}\mathbf{h}}{\left(\mathbf{h}^2 + \mathbf{R}^2\right)^{3/2}}$$

for maximum electric field

$$\frac{dE}{dh} = 0$$

$$\Rightarrow$$
 h = $\frac{R}{\sqrt{2}}$

29. Three blocks A, B and C are lying on a smooth horizontal surface, as shown in the figure. A and B have equal masses, m while C has mass M. Block A is given an brutal speed v towards B due to which it collides with B perfectly inelastically. The combined mass collides with C also perfectly inelastically.

C, also perfectly inelastically $\frac{5}{6}$ th of the initial kinetic energy is lost in whole process. What is value of M/m ?

Ans. (1)

Sol.
$$k_i = \frac{1}{2}mv_0^2$$

From linear momentum conservation $mv_0 = (2m + M) v_f$

$$\Rightarrow v_{f} = \frac{mv_{0}}{2m + M}$$

$$\frac{k_{i}}{k_{f}} = 6$$

$$\Rightarrow \frac{\frac{1}{2}mv_{0}^{2}}{\frac{1}{2}(2m + M)\left(\frac{mv_{0}}{2m + M}\right)^{2}} = 6$$

$$\Rightarrow \frac{2m + M}{m} = 6$$

$$\Rightarrow \frac{M}{m} = 4$$

30. Drift speed of electrons, when 1.5 A of current flows in a copper wire of cross section 5 mm2, is v. If the electron density in copper is 9×10^{28} /m³ the value of v in mm/s is close to (Take charge of electron to be = 1.6×10^{-19} C)

Ans. (4)

Sol.
$$I = neAv_d$$

$$\Rightarrow v_{d} = \frac{I}{neA} = \frac{1.5}{9 \times 10^{28} \times 1.6 \times 10^{-19} \times 5 \times 10^{-6}}$$
$$= 0.02 \text{ m/s}$$



E

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Wednesday 09th JANUARY, 2019) TIME : 9 : 30 AM To 12 : 30 PM CHEMISTRY

- 1. Which one of the following statements regarding Henry's law not correct ?
 - (1) The value of $K_{\rm H}$ increases with function of the nature of the gas
 - (2) Higher the value of K_H at a given pressure, higher is the solubility of the gas in the liquids.
 - (3) The partial of the gas in vapour phase is proportional to the mole fraction of the gas in the solution.
 - (4) Different gases have different K_H (Henry's law constant) values at the same temperature.

Ans. (2)

Sol. Liquid solution

 $P_{gas} = K_H \times X_{gas}$

More is K_H less is solubility, lesser solubility is at higher temperature. So more is temperature more is K_H .

- 2. The correct decreasing order for acid strength is :-
 - (1) $NO_2CH_2COOH > NCCH_2COOH >$ $FCH_2COOH > CICH_2COOH$
 - (2) $FCH_2COOH > NCCH_2COOH >$

 $NO_2CHCOOH > CICH_2COOH$

- (3) $NO_2CH_2COOH > FCH_2COOH >$ $CNCH_2COOH > CICH_2COOH$
- (4) $CNCH_2COOH > O_2NCH_2COOH >$ $FCH_2COOH > CICH_2COOH$

Ans. (1)

Ε

Sol. EWG increasea acidic strength

$$\label{eq:compared} \begin{split} \text{NO}_2\text{CH}_2\text{COOH} > \text{NCCH}_2\text{COOH} > \\ \text{FCH}_2\text{COOH} > \text{CICH}_2\text{COOH} \end{split}$$

- 3. Two complexes $[Cr(H_2O_6)Cl_3]$ (A) and $[Cr(NH_3)_6]Cl_3$ (B) are violet and yellow coloured, respectively. The incorrect statement regarding them is :
 - (1) Δ_0 value of (A) is less than that of (B).
 - (2) Δ_0 value of (A) and (B) are calculated from the energies of violet and yellow light, respectively
 - (3) Bothe absorb energies corresponding to their complementary colors.
 - (4) Bothe are paramagnetic with three unpaired electrons.
- Ans. (2)
- Sol. Δ_0 order will be compared by spectro chemical series not by energies of violet & yellow light so Δ_0 order is

 $[Cr(H_2O)_6]Cl_3 < [Cr(NH_3)_6]Cl_3$

4. Adsorption of a gas follows Freundlich adsorption isotherm. In the given plot, x is the mass of the gas adsorbed on mass m of the

adsorbent at pressure p. $\frac{x}{m}$ is proportional to



(1)
$$P\frac{1}{4}$$
 (2) P^2 (3) P (4) $P\frac{1}{2}$

2

Ans. (4)

Sol.
$$\frac{x}{m} = K \times P^{1/n}$$

 $\log \frac{x}{m} = \log K + \frac{1}{n} \log P$
 $m = \frac{1}{n} = \frac{2}{4} = \frac{1}{2} \implies n = 2$
So, $\frac{x}{m} = K \times P^{1/2}$



JEE (Main) Examination-2019/Morning Session/09-01-2019

- 5. Correct statements among a to d regarding silicones are :
 - (a) They are polymers with hydrophobic character
 - (b) They are biocompatible.
 - (c) In general, they have high thermal stability and low dielectric strenth.
 - (d) Usually, they are resistant to oxidation and used as greases.
 - (1) (a), (b) and (c) only
 - (2) (a), and (b) only
 - (3) (a), (b), (c) and (d)
 - (4) (a), (b) and (d) only

Ans. (3)

- Sol. These are properties and uses of silicones.
- 6. For emission line of atomic hydrogen from $n_i = 8$ to $n_f =$ the plot of wave number $\overline{(v)}$ against $\left(\frac{1}{n^2}\right)$ will be (The Ry dberg constant, R_H is in wave number unit).
 - (1) Linear with slope R_H
 - (2) Linear with intercept R_H
 - (3) Non linear
 - (4) Linear with sslope R_H

Ans. (4)

Sol.
$$\frac{1}{\lambda} = \overline{v} = R_{H} z^{2} \left(\frac{1}{\eta_{I}^{2}} - \frac{1}{\eta_{2}^{2}} \right)$$
$$\overline{v} = R_{H} \times \left(\frac{1}{\eta_{I}^{2}} - \frac{1}{8^{2}} \right)$$
$$\overline{v} = R_{H} \times \frac{1}{\eta^{2}} - \frac{R_{H}}{8^{2}}$$
$$\overline{v} = R_{H} \times \frac{1}{\eta^{2}} - \frac{R_{H}}{64}$$
$$m = R_{H}$$
Linear with slope R_{H}

7. The major product the following reaction is :





Sol.

 $NH_2(a)$ will wact as nucleophile as (b) is having delocalised lonepair.



10. The highest value of the calculated spin only magnetic moment (in BM) among all the transition metal complexs is :

Ans. (1)

Sol. $\mu = \sqrt{n(n+2)}$ B.M.

n = Number of unpaired electrons

n = Maximum number of unpaired electron = 5

 $Ex : Mn^{2+}$ complex.

11. 20 mL of 0.1 MH_2SO_4 solution is added to 30 mL of 0.2 M NH_4OH solution. The pH of the resulatant mixture is : $[pk_b \text{ of } NH_4OH = 4.7]$.

(1) 9.4 (2) 5.0 (3) 9.0 (4) 5.2

Ans. (3)

Sol. 20 ml 0.1 M
$$H_2SO_4 \implies \eta_{H^+} = 4$$

30 ml 0.2 M NH₄OH \Rightarrow $\eta_{\text{NH}_4\text{OH}} = 6$

$$NH_4OH + H^+ \xrightarrow{\oplus} NH_4^{\oplus} + H_2O$$

$$\Rightarrow 6 \quad 4 \quad 0 \quad 0$$

$$\Rightarrow 2 \quad 0 \quad 4 \quad 4$$

Solution is basic buffer

$$pOH = pK_{b} + \log \frac{NH_{4}^{+}}{NH_{4}OH}$$
$$= 4.7 + \log 2$$
$$= 4.7 + 0.3 = 5$$
$$pH = 14 - 5 = 9$$

12. 0.5 moles of gas A and x moles of gas B exert a pressure of 200 Pa in a a container of volume 10 m³ at 1000 K. given R is the gas constant in JK⁻¹ mol⁻¹m, x is :

(1)
$$\frac{2R}{4+12}$$
 (2) $\frac{2R}{4-R}$ (3) $\frac{4-R}{2R}$ (4) $\frac{4+R}{2R}$

Ans. (3)

Sol.
$$n_T = (0.5 + x)$$

 $PV = n \times R \times T$
 $200 \times 10 = (0.5 + x) \times R \times 1000$
 $2 = (0.5 + x) R$
 $\frac{2}{R} = \frac{1}{2} + x$
 $\frac{4}{R} - 1 = 2x$
 $\frac{4-R}{2R} = x$

13. Consider the reversible isothermal expansion of an ideal gas in a closed system at two different temperatures T_1 and T_2 ($T_1 < T_2$). The correct graphical depiction of the dependence of work done (w) on the final volume (V) is:



collegebatch.com

Sol. $w = -nRT ln \frac{V_2}{V}$ $w = -nRT \ln \frac{V_b}{V_b}$ $|w| = nRT \ln \frac{V_b}{V_c}$ $|\mathbf{w}| = \mathbf{nRT} (\ln \mathbf{V}_{\rm h} - \ln \mathbf{V}_{\rm i})$ $|w| = nRT \ln V_{b} - nRT \ln V_{i}$ Y = m x - CSo, slope of curve 2 is more than curve 1 and intercept of curve 2 is more negative then curve 1. 14. The major product of following reaction is : $R - C \equiv N \xrightarrow{(1)AlH(i-Bu_2)} ?$ (2)H₂O (1) RCHO (2) RCOOH (3) RCH_2NH_2 (4) RCONH₂ Ans. (1)

Sol.
$$R-C=N \xrightarrow{AlH(i-Bu_2)} R-CH=N-\xrightarrow{H_2O} R-CH=C$$

- **15.** In general, the properties that decrease and increase down a group in the periodic table, respectively, are :
 - (1) electronegativity and electron gain enthalpy.
 - (2) electronegativity and atomic radius.
 - (3) atomic radius and electronegativity.
 - (4) electron gain enthalpy and electronegativity.

Ans. (2)

- **Sol.** Electronegativity decreases as we go down the group and atomic radius increases as we go down the group.
- 16. A solution of sodium sulfate contains 92 g of Na⁺ ions per kilogram of water. The molality of Na⁺ ions in that solution in mol kg⁻¹ is:
 (1) 16
 (2) 8
 (3) 4
 (4) 12

Ans. (4)

Sol. $n_{Na^+} = \frac{92}{23} = 4$

So molality = 4

17. A water sample has ppm level concentration of the following metals: Fe= 0.2; Mn = 5.0; Cu = 3.0; Zn = 5.0. The metal that makes the water sample unsuitable drinking is : (1) Zn (2) Fe (3) Mn (4) Cu Ans. (3) **Sol.** (i) Zn = 0.2(ii) Fe = 0.2(iii) Mn = 5.0(iv) Cu = 3.018. The increasing order of pKa of the following amino acids in aqueous solution is : Gly Asp Lys Arg (1) Asp < Gly < Arg < Lys(2) Arg < Lys < Gly < Asp(3) Gly < Asp < Arg < Lys(4) Asp < Gly < Lys < ArgAns. (4) Sol. Order of acidic strength :

Aspartic acid

Arginine

So, pK_a

19. According to molecular orbital theory, which of the following is true with respect to Li_2^+ and

 $Li_{2}^{-}?$

- (1) Both are unstable
- (2) Li_2^+ is unstable and Li_2^- is stable
- (3) Li_2^+ is stable and Li_2^- is unstable
- (4) Both are stabel

Ans. (4)

Sol. Both Li_2^+ and Li_2^- has 0.5 bond order and hence both are stable.

4



20. The following results were obtained during kinetic studies of the reaction : $2A + B \rightarrow Products$ Sol.

Experment	[A] (in mol L ⁻¹)	[B] (in mol L ⁻¹)	Initial Rate of reaction (in mol $L^{-1} min^{-1}$)
(I)	0.10	0.20	6.93×10^{-3}
(II)	0.10	0.25	6.93×10^{-3}
(III)	0.20	0.30	1.386×10^{-2}

The time (in minutes) required to consume half of A is :

(1) 10 (2) 5 (3) 100 (4) 1 Ans. (2) Sol. $6.93 \times 10^{-3} = K \times (0.1)^x (0.2)^y$

 $6.93 \times 10^{-3} = K \times (0.1)^{x} (0.25)^{y}$ So y = 0 and 1.386 × 10⁻² = K × (0.2)^{x} (0.30)^{y}

 $\frac{1}{2} = \left(\frac{1}{2}\right)^x \quad \boxed{\mathbf{x} = 1}$

So $r = K \times (0.1) \times (0.2)^0$ 6.93 × 10⁻³ = K × 0.1 × (0.2)⁰

 $K = 6.93 \times 10^{-2}$

$$t_{1/2} = \frac{0.693}{2K} = \frac{0.693}{0.693 \times 10^{-1} \times 2} = \frac{10}{2} = 5$$

21. The major product of the following reaction is:



Ans. (2)

Ε



During AES Br is o/p directing and major product will be formed on less hindrance p position :

22. Arrange the following amines in the decreasing order of basicity:



Ans. (4)

Sol. Order of basic strength :



23. Which amongst the following is the strongest acid ?

(1)
$$CHI_3$$
 (2) $CHCI_3$
(3) $CHBr_3$ (4) $CH(CN)_3$

Ans. (4)

- Sol. CN makes anino most stable so answer is $CH(CN)_3$
- 24. The anodic half-cell of lead-acid battery is recharged unsing electricity of 0.05 Faraday. The amount of $PbSO_4$ electrolyzed in g during the process in : (Molar mass of $PbSO_4 = 303$ g mol⁻¹)

(1) 22.8 (2) 15.2 (3) 7.6 (4) 11.4 **Ans. (2)**



JEE (Main) Examination-2019/Morning Session/09-01-2019

- Sol. (A) $PbSO_4(s) + 2OH \longrightarrow PbO_2 + H_2SO_4 + 2e^{-1}$ 0.05/2 mole 0.05F (C) $PbSO_4 + 2e^- + 2H^+ \longrightarrow Pb(s) + H_2SO_4$ 0.05/2 mole 0.05 F $n_{T}(PbSO_{4}) = 0.05$ mole $m_{PbSO_4} = 0.05 \times 303 = 15.2 \text{ gm}$ 25. The one that is extensively used as a piezoelectric material is : (1) Quartz (2) Amorphous silica (3) Mica (4) Tridymite Ans. (1) Sol. Quartz (Information) 26. Aluminium is usually found in +3 oxidation stagte. In contarast, thallium exists in +1 and +3 oxidation states. This is due to : (1) lanthanoid contraction (2) lattice effect (3) diagonal relationship (4) inert pair effect Ans. (4) Sol. Inert pair effect is promenent character of p-
- 27. The correct match between Item -I and Item-II is :

	Item – I (drug)		Item – II (test)
(A)	Chloroxylenol	(P)	Carbylamine Test
(B)	Norethindrone	(Q)	Sodium Hydrogen carbonateTest
(C)	Sulphapyridine	(R)	Ferric chloride test
(D)	Penicillin	(S)	Bayer's test

- (1) $A \rightarrow Q$; $B \rightarrow P$; $C \rightarrow S$; $D \rightarrow R$
- (2) $A \rightarrow R$; $B \rightarrow P$; $C \rightarrow S$; $D \rightarrow Q$
- (3) $A \rightarrow R$; $B \rightarrow S$; $C \rightarrow P$; $D \rightarrow Q$
- (4) $A \rightarrow Q$; $B \rightarrow S$; $C \rightarrow P$; $D \rightarrow R$

Ans. (3)



28. The ore that contains both iron and copper is:

- (1) malachite
- (2) dolomite
- (3) azurite
- (4) copper pyrites

Ans. (4)

6



29. The compounds A and B in the following reaction are, respectively:

$$\underbrace{HCHO+HCI}_{HCHO+HCI} \land \underbrace{AgCN}_{AgCN} \land AgCN}_{AgCN}$$

(1) A = Benzyl alcohol, B = Benzyl isocyanide

(2) A = Benzyl alcohol, B = Benzyl cyanide

(3) A = Benzyl chloride, B = Benzyl cyanide

(4) A = Benzyl chloride, B = Benzyl isocyanideAns. (4)



- **30.** The isotopes of hydrogen are :
 - (1) Tritium and protium only
 - (2) Deuterium and tritium only
 - (3) Protium and deuterum only
 - (4) Protium, deuterium and tritium

Ans. (4)

Sol. Isotopes of hydrogen is : Proteium Deuterium Tritium

Ε

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Wednesday 09th JANUARY, 2019) TIME : 02 : 30 PM To 05 : 30 PM M ATHEM ATICS

1. Let f be a differentiable function from R to R such that $|f(x) - f(y)| \le 2|x - y|^{\frac{3}{2}}$, for all x, y ε R. If

$$f(0) = 1$$
 then $\int_{0}^{1} f^{2}(x) dx$ is equal to

(1) 0 (2)
$$\frac{1}{2}$$
 (3) 2 (4) 1

Ans. (4)

Sol. $|f(\mathbf{x}) - f(\mathbf{y})| \le 2|\mathbf{x} - \mathbf{y}|^{3/2}$ divide both sides by $|\mathbf{x} - \mathbf{y}|$

$$\left|\frac{f(\mathbf{x}) - f(\mathbf{y})}{\mathbf{x} - \mathbf{y}}\right| \le 2 \cdot |\mathbf{x} - \mathbf{y}|^{1/2}$$

apply limit $x \to y$ $|f'(y)| \le 0 \Rightarrow f'(y) = 0 \Rightarrow f(y) = c \Rightarrow f(x) = 1$ $\int_{0}^{1} 1.dx = 1$

2. If
$$\int_{0}^{\frac{\pi}{3}} \frac{\tan \theta}{\sqrt{2k \sec \theta}} d\theta = 1 - \frac{1}{\sqrt{2}}, (k > 0), \text{ then the}$$

value of k is :

(1) 2 (2)
$$\frac{1}{2}$$
 (3) 4 (4) 1

Ans. (1)

Sol.
$$\frac{1}{\sqrt{2k}} \int_{0}^{\pi/3} \frac{\tan \theta}{\sqrt{\sec \theta}} d\theta = \frac{1}{\sqrt{2k}} \int_{0}^{\pi/3} \frac{\sin \theta}{\sqrt{\cos \theta}} d\theta$$
$$= -\frac{1}{\sqrt{2k}} 2\sqrt{\cos \theta} \Big|_{0}^{\pi/3} = -\frac{\sqrt{2}}{\sqrt{k}} \left(\frac{1}{\sqrt{2}} - 1\right)$$
given it is $1 - \frac{1}{\sqrt{2}} \Longrightarrow k = 2$

3. The coefficient of t⁴ in the expansion of

$$\left(\frac{1-t^{6}}{1-t}\right)^{3}$$
 is
(1) 12 (2) 15 (3) 10 (4) 14
Ans. (2)
Sol. $(1-t^{6})^{3}(1-t)^{-3}$
 $(1-t^{18}-3t^{6}+3t^{12})(1-t)^{-3}$
 \Rightarrow cofficient of t⁴ in $(1-t)^{-3}$ is
 $^{3+4+1}C_{4} = ^{6}C_{2} = 15$
4. For each xcR, let [x] be the greatest integer less
than or equal to x. Then

$$\lim_{x\to 0^{-}} \frac{x([x]+|x|)\sin[x]}{|x|}$$
 is equal to
(1) - sin1 (2) 0 (3) 1 (4) sin1
Ans. (1)
Sol.
$$\lim_{x\to 0^{-}} \frac{x([x]+|x|)\sin[x]}{|x|}$$

 $x \to 0^{-}$
 $[x]=-1 \Rightarrow \lim_{x\to 0^{-}} \frac{x(-x-1)\sin(-1)}{-x} = -\sin 1$
 $|x|=-x$
5. If both the roots of the quadratic equation
 $x^{2} - mx + 4 = 0$ are real and distinct and they lie
in the interval [1,5], then m lies in the interval:
(1) (4,5) (2) (3,4) (3) (5,6) (4) (-5,-4)
Ans. (Bonus/I)
Sol. $x^{2} - mx + 4 = 0$
 $\alpha,\beta \in [1,5]$
(1) $D > 0 \Rightarrow m^{2} - 16 > 0$
 $\Rightarrow m \in (-\infty,-4) \cup (4,\infty)$
(2) $f(1) \ge 0 \Rightarrow 5 - m \ge 0 \Rightarrow m \in (-\infty,5]$
(3) $f(5) \ge 0 \Rightarrow 29 - 5m \ge 0 \Rightarrow m \in (-\infty,5]$
(4) $1 < \frac{-b}{2a} < 5 \Rightarrow 1 < \frac{m}{2} < 5 \Rightarrow m \in (2,10)$
 $\Rightarrow m \in (4,5)$
No option correct : Bonus
* If we consider $\alpha,\beta \in (1,5)$ then option (1) is
correct.

Е

collegebatch.com

JEE (Main) Examination-2019/ Evening Session/ 09-01-2019

6. If $A = \begin{bmatrix} e^{t} & e^{-t} \cos t & e^{-t} \sin t \\ e^{t} & -e^{-t} \cos t - e^{-t} \sin t & -e^{-t} \sin t + e^{-t} \cos t \\ e^{t} & 2e^{-t} \sin t & -2e^{-t} \cos t \end{bmatrix}$ Then A is-(1) Invertible only if $t = \frac{\pi}{2}$ (2) not invertible for any teR

(2) invertible for all tɛR

(4) invertible only if $t=\pi$

Ans. (3)

Sol.
$$|\mathbf{A}| = e^{-t} \begin{vmatrix} 1 & \cos t & \sin t \\ 1 & -\cos t - \sin t & -\sin t + \cos t \\ 1 & 2\sin t & -2\cos t \end{vmatrix}$$

- $= e^{-t}[5\cos^2 t + 5\sin^2 t] \ \forall \ t \in R$ $= 5e^{-t} \neq 0 \ \forall \ t \in R$
- 7. The area of the region

$$A = \left[(x, y) : 0 \le y \le x |x| + 1 \text{ and } -1 \le x \le 1 \right]$$

in sq. units, is :

(1)
$$\frac{2}{3}$$
 (2) $\frac{1}{3}$ (3) 2 (4) $\frac{4}{3}$

Ans. (3)

Sol. The graph is a follows



$$\int_{-1}^{0} (-x^{2} + 1) dx + \int_{0}^{1} (x^{2} + 1) dx = 2$$

8. Let z_0 be a root of the quadratic equation, $x^2 + x + 1 = 0$. If $z = 3 + 6iz_0^{81} - 3iz_0^{93}$, then arg z is equal to:

(1)
$$\frac{\pi}{4}$$
 (2) $\frac{\pi}{3}$ (3) 0 (4) $\frac{\pi}{6}$

Ans. (1)

Sol. $z_0 = \omega$ or ω^2 (where ω is a non-real cube root of unity) $z = 3 + 6i(\omega)^{81} - 3i(\omega)^{93}$

$$z = 3 + 31$$

 $\Rightarrow \arg z = \frac{\pi}{4}$

 $-2 \pm 2i$

9. Let $\vec{a} = \hat{i} + \hat{j} + \sqrt{2}\hat{k}$, $\vec{b} = b_1\hat{i} + b_2\hat{j} + \sqrt{2}\hat{k}$ and $\vec{c} = 5\hat{i} + \hat{j} + \sqrt{2}\hat{k}$ be three vectors such that the projection vector of \vec{b} on \vec{a} is \vec{a} . If $\vec{a} + \vec{b}$ is perpendicular to \vec{c} , then $|\vec{b}|$ is equal to:

(1)
$$\sqrt{22}$$
 (2) 4 (3) $\sqrt{32}$ (4) 6
Ans. (4)

Sol. Projection of
$$\vec{b}$$
 on $\vec{a} = \frac{\vec{a}.\vec{b}}{|\vec{a}|} = |\vec{a}|$
 $\Rightarrow b_1 + b_2 = 2 \qquad \dots(1)$
and $(\vec{a} + \vec{b}) \perp \vec{c} \Rightarrow (\vec{a} + \vec{b}).\vec{c} = 0$
 $\Rightarrow 5b_1 + b_2 = -10 \qquad \dots(2)$
from (1) and (2) $\Rightarrow b_1 = -3$ and $b_2 =$
then $|\vec{b}| = \sqrt{b_1^2 + b_2^2 + 2} = 6$

10. Let A(4,-4) and B(9,6) be points on the parabola, $y^2 + 4x$. Let C be chosen on the arc AOB of the parabola, where O is the origin, such that the area of \triangle ACB is maximum. Then, the area (in sq. units) of \triangle ACB, is:

5

(1)
$$31\frac{3}{4}$$
 (2) 32 (3) $30\frac{1}{2}$ (4) $31\frac{1}{4}$

Ans. (4)







12. An urn contains 5 red and 2 green balls. A ball is drawn at random from the urn. If the drawn ball is green, then a red ball is added to the urn and if the drawn ball is red, then a green ball is added to the urn; the original ball is not returned to the urn. Now, a second ball is drawn at random from it. The probability that the second ball is red, is :

(1)
$$\frac{26}{49}$$
 (2) $\frac{32}{49}$ (3) $\frac{27}{49}$ (4) $\frac{21}{49}$

Ans. (2)

Ε

- JEE (Main) Examination-2019/ Evening Session/ 09-01-2019
 - Sol. E_1 : Event of drawing a Red ball and placing a green ball in the bag E_2 : Event of drawing a green ball and placing a red ball in the bag E: Event of drawing a red ball in second draw $P(E) = P(E_1) \times P\left(\frac{E}{E_1}\right) + P(E_2) \times P\left(\frac{E}{E_2}\right)$ $= \frac{5}{7} \times \frac{4}{7} + \frac{2}{7} \times \frac{6}{7} = \frac{32}{49}$
 - 13. If $0 \le x < \frac{\pi}{2}$, then the number of values of x for which $\sin x \sin 2x + \sin 3x = 0$, is
 - (1) 2 (2) 1
 - (3) 3 (4) 4

Ans. (1)

Sol.
$$\sin x - \sin 2x + \sin 3x = 0$$

 $\Rightarrow (\sin x + \sin 3x) - \sin 2x = 0$
 $\Rightarrow 2\sin x. \cos x - \sin 2x = 0$
 $\Rightarrow \sin 2x(2\cos x - 1) = 0$
 $\Rightarrow \sin 2x = 0 \text{ or } \cos x = \frac{1}{2}$
 $\Rightarrow x = 0, \frac{\pi}{3}$

14. The equation of the plane containing the straight line $\frac{x}{2} = \frac{y}{3} = \frac{z}{4}$ and perpendicular to the plane containing the straight lines

$$\frac{x}{3} = \frac{y}{4} = \frac{z}{2} \text{ and } \frac{x}{4} = \frac{y}{2} = \frac{z}{3} \text{ is:}$$
(1) $x + 2y - 2z = 0$ (2) $x - 2y + z = 0$
(3) $5x + 2y - 4z = 0$ (4) $3x + 2y - 3z = 0$

Ans. (2)



JEE (Main) Examination-2019/ Evening Session/ 09-01-2019

Sol. Vector along the normal to the plane containing the lines

$$\frac{x}{3} = \frac{y}{4} = \frac{z}{2}$$
 and $\frac{x}{4} = \frac{y}{2} = \frac{z}{3}$
is $(8\hat{i} - \hat{j} - 10\hat{k})$

vector perpendicular to the vectors $2\hat{i} + 3\hat{j} + 4\hat{k}$ and $8\hat{i} - \hat{j} - 10\hat{k}$ is $26\hat{i} - 52\hat{j} + 26\hat{k}$ so, required plane is 26x - 52y + 26z = 0x - 2y + z = 0

- 15. Let the equations of two sides of a triangle be 3x 2y + 6 = 0 and 4x + 5y 20 = 0. If the orthocentre of this triangle is at (1,1), then the equation of its third side is :
 - $(1) \ 122y 26x 1675 = 0$
 - $(2) \ 26x + 61y + 1675 = 0$
 - (3) 122y + 26x + 1675 = 0
 - $(4) \ 26x 122y 1675 = 0$

Ans. (4)

Sol. Equation of AB is 3x - 2y + 6 = 0equation of AC is 4x + 5y - 20 = 0Equation of BE is 2x + 3y - 5 = 0



Equation of CF is 5x - 4y - 1 = 0 \Rightarrow Equation of BC is 26x - 122y = 1675

16. If
$$x = 3$$
 tan t and $y = 3$ sec t, then the value of

$$\frac{d^2 y}{dx^2} \text{ at } t = \frac{\pi}{4}, \text{ is:}$$
(1) $\frac{3}{2\sqrt{2}}$ (2) $\frac{1}{3\sqrt{2}}$ (3) $\frac{1}{6}$ (4) $\frac{1}{6\sqrt{2}}$

Ans. (4)

Sol.
$$\frac{dx}{dt} = 3 \sec^2 t$$
$$\frac{dy}{dt} = 3 \sec t \tan t$$
$$\frac{dy}{dt} = 3 \sec t \tan t$$
$$\frac{dy}{dx} = \frac{\tan t}{\sec t} = \sin t$$
$$\frac{d^2 y}{dx^2} = \cos t \frac{dt}{dx}$$
$$= \frac{\cos t}{3 \sec^2 t} = \frac{\cos^3 t}{3} = \frac{1}{3 \cdot 2\sqrt{2}} = \frac{1}{6\sqrt{2}}$$

17. If $x = \sin^{-1}(\sin 10)$ and $y=\cos^{-1}(\cos 10)$, then y-x is equal to:

(1)
$$\pi$$
 (2) 7π (3) 0 (4) 10 (1)

Ans. (1)



Ans. (2)

Sol. Line x = ay + b, $z = cy + d \Rightarrow \frac{x - b}{a} = \frac{y}{1} = \frac{z - d}{c}$

Line x = a'z + b', y = c'z + d'

$$\Rightarrow \frac{x-b'}{a'} = \frac{y-d'}{c'} = \frac{z}{1}$$

(4) 4

Given both the lines are perpendicular \Rightarrow aa' + c' + c = 0

19. The number of all possible positive integral values of α for which the roots of the quadratic equation, $6x^2-11x+\alpha = 0$ are rational numbers is :

(3) 3

Sol. $6x^2 - 11x + \alpha = 0$ given roots are rational \Rightarrow D must be perfect square $\Rightarrow 121 - 24\alpha = \lambda^2$ \Rightarrow maximum value of α is 5 $\alpha = 1 \Rightarrow \lambda \notin I$ $\alpha = 2 \Rightarrow \lambda \notin I$ $\alpha = 3 \Rightarrow \lambda \in I$ $\Rightarrow 3$ integral values $\alpha = 4 \Rightarrow \lambda \in I$ $\alpha = 5 \Rightarrow \lambda \in I$





20. A hyperbola has its centre at the origin, passes through the point (4,2) and has transverse axis of length 4 along the x-axis. Then the eccentricity of the hyperbola is :

(1)
$$\frac{2}{\sqrt{3}}$$
 (2) $\frac{3}{2}$ (3) $\sqrt{3}$ (4) 2

Ans. (1)



$$\frac{x^{2}}{a^{2}} - \frac{y^{2}}{b^{2}} = 1$$

$$2a = 4 \qquad a = \frac{x^{2}}{4} - \frac{y^{2}}{b^{2}} = 1$$

Passes through (4,2)

$$4 - \frac{4}{b^2} = 1 \Longrightarrow b^2 = \frac{4}{3} \Longrightarrow e = \frac{2}{\sqrt{3}}$$

21. Let $A = \{x \in \mathbb{R} : x \text{ is not a positive integer}\}$

2

Define a function $f: A \rightarrow R$ as $f(x) = \frac{2x}{x-1}$ then f is

- (1) injective but not surjective
- (2) not injective
- (3) surjective but not injective
- (4) neither injective nor surjective

Ans. (1)

E

Sol. $f(x) = 2\left(1 + \frac{1}{x - 1}\right)$ $f'(x) = -\frac{2}{(x - 1)^2}$ $\Rightarrow f$ is one-one but not onto 22. If $f(x) = \int \frac{5x^8 + 7x^6}{(x^2 + 1 + 2x^7)^2} dx, (x \ge 0)$ and f(0) = 0, then the value of f(1) is : $(1) -\frac{1}{2}$ (2) $\frac{1}{2}$ (3) $-\frac{1}{4}$ (4) $\frac{1}{4}$ Ans. (4)

Sol.
$$\int \frac{5x^{8} + 7x^{6}}{(x^{2} + 1 + 2x^{7})^{2}} dx$$

$$= \int \frac{5x^{-6} + 7x^{-8}}{(\frac{1}{x^{7}} + \frac{1}{x^{5}} + 2)^{2}} dx = \frac{1}{2 + \frac{1}{x^{5}} + \frac{1}{x^{7}}} + C$$

As $f(0) = 0$, $f(x) = \frac{x^{7}}{2x^{7} + x^{2} + 1}$
 $f(1) = \frac{1}{4}$
23. If the circles $x^{2} + y^{2} - 16x - 20y + 164 = r^{2}$ and
 $(x - 4)^{2} + (y - 7)^{2} = 36$ intersect at two distinct
points, then:
(1) $0 < r < 1$ (2) $1 < r < 11$
(3) $r > 11$ (4) $r = 11$
Ans. (2)
Sol. $x^{2} + y^{2} - 16x - 20y + 164 = r^{2}$
 $A(8,10), R_{1} = r$
 $(x - 4)^{2} + (y - 7)^{2} = 36$
 $B(4,7), R_{2} = 6$
 $|R_{1} - R_{2}| < AB < R_{1} + R_{2}$
 $\Rightarrow 1 < r < 11$
24. Let S be the set of all triangles in the xy-plane,
each having one vertex at the origin and the other
two vertices lie on coordinate axes with integral
coordinates. If each triangle in S has area 50sq.
units, then the number of elements in the set S is:
(1) 9 (2) 18 (3) 32 (4) 36
Ans. (4)
Sol. Let $A(\alpha,0)$ and $B(0,\beta)$
be the vectors of the given triangle AOB
 $\Rightarrow |\alpha\beta| = 100$
 \Rightarrow Number of triangles
 $= 4 \times (number of divisors of 100)$
 $= 4 \times 9 = 36$
25. The sum of the follwing series
 $1 + 6 + \frac{9(1^{2} + 2^{2} + 3^{2})}{7} + \frac{12(1^{2} + 2^{2} + 3^{2} + 4^{2})}{9}$
 $+ \frac{15(1^{2} + 2^{2} + ... + 5^{2})}{11} + up to 15 terms, is:$

(1) 7820 (2) 7830 (3) 7520 (4) 7510 Ans. (1)

5

collegebatch.com

JEE (Main) Examination-2019/ Evening Session/ 09-01-2019

27.

Sol.
$$T_n = \frac{(3+(n-1)\times 3)(1^2+2^2+...+n^2)}{(2n+1)}$$

$$T_{n} = \frac{3 \cdot \frac{n(n+1)(2n+1)}{6}}{2n+1} = \frac{n^{2}(n+1)}{2}$$

$$S_{15} = \frac{1}{2} \sum_{n=1}^{15} \left(n^3 + n^2 \right) = \frac{1}{2} \left[\left(\frac{15(15+1)}{2} \right)^2 + \frac{15 \times 16 \times 31}{6} \right]$$
$$= 7820$$

26. Let a, b and c be the 7th, 11th and 13th terms respectively of a non-constant A.P. If these are also the three consecutive terms of a G.P., then $\frac{a}{c}$ is equal to:

(1)
$$\frac{1}{2}$$
 (2) 4

(3) 2 (4)
$$\frac{7}{13}$$

Ans. (2)

Sol. a = A + 6d b = A + 10d c = A + 12d a,b,c are in G.P. $\Rightarrow (A + 10d)^2 = (A + 6d) (a + 12d)$ $\Rightarrow \frac{A}{d} = -14$

$$\frac{a}{c} = \frac{A+6d}{A+12d} = \frac{6+\frac{A}{d}}{12+\frac{A}{d}} = \frac{6-14}{12-14} = 4$$

x-4y+7z = g3y - 5z = h-2x + 5y - 9z = kis consistent, then : (1) g + h + k = 0(2) 2g + h + k = 0(3) g + h + 2k = 0(4) g + 2h + k = 0Ans. (2) **Sol.** $P_1 \equiv x - 4y + 7z - g = 0$ $P_2 \equiv 3x - 5y - h = 0$ $P_3 \equiv -2x + 5y - 9z - k = 0$ Here $\Delta = 0$ $2P_1 + P_2 + P_3 = 0$ when 2g + h + k = 0Let $f:[0,1] \rightarrow \mathbb{R}$ be such that $f(xy) = f(x) \cdot f(y)$ for all 28. x,y, $\varepsilon[0,1]$, and $f(0) \neq 0$. If y = y(x) satisfies the differential equation, $\frac{dy}{dx} = f(x)$ with y(0) = 1, then $y\left(\frac{1}{4}\right) + y\left(\frac{3}{4}\right)$ is equal to (2) 3 (4) 2 (1) 4(3) 5 Ans. (2) **Sol.** f(xy) = f(x). f(y)f(0) = 1 as $f(0) \neq 0$ $\Rightarrow f(\mathbf{x}) = 1$ $\frac{\mathrm{dy}}{\mathrm{dx}} = f(\mathbf{x}) = 1$ \Rightarrow y = x + c At, x = 0, $y = 1 \implies c = 1$ y = x + 1 $\Rightarrow y\left(\frac{1}{4}\right) + y\left(\frac{3}{4}\right) = \frac{1}{4} + 1 + \frac{3}{4} + 1 = 3$ 29. A data consists of n observations: x_1, x_2, \dots, x_n . If $\sum_{i=1}^n (x_i + 1)^2 = 9n$ and $\sum_{i=1}^{n} (x_i - 1)^2 = 5n$, then the standard deviation of

If the system of linear equations

this data is :

(1) 5 (2)
$$\sqrt{5}$$
 (3) $\sqrt{7}$ (4) 2
Ans. (2)



collegebatch.com

Sol. $\sum (x_i + 1)^2 = 9n$...(1) $\sum (x_i - 1)^2 = 5n$...(2) $(1) + (2) \Rightarrow \sum (x_1^2 + 1) = 7n$ $\Rightarrow \frac{\sum x_i^2}{n} = 6$ (1) - (2) $\Rightarrow 4\Sigma x_i = 4n$ $\Rightarrow \Sigma x_i = n$ $\Rightarrow \frac{\Sigma x_i}{n} = 1$ \Rightarrow variance = 6 - 1 = 5 \Rightarrow Standard diviation $=\sqrt{5}$ The number of natural numbers less than 7,000 30. which can be formed by using the digits 0,1,3,7,9(repitition of digits allowed) is equal to : (1) 250(2) 374 (3) 372 (4) 375 Ans. (2) Sol. $\begin{vmatrix} a_1 & a_2 & a_3 \end{vmatrix}$ Number of numbers $= 5^3 - 1$ $\mathbf{a}_4 \mid \mathbf{a}_1 \mid \mathbf{a}_2 \mid \mathbf{a}_3$ 2 ways for a_4 Number of numbers = 2×5^3 Required number = $5^3 + 2 \times 5^3 - 1$ = 374



Ε



TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Wednesday 09th JANUARY, 2019) TIME : 2 : 30 PM To 05 : 30 PM PHYSICS

1. Two plane mirrors arc inclined to each other such that a ray of light incident on the first mirror (M₁) and parallel to the second mirror (M₂) is finally reflected from the second mirror (M₂) parallel to the first mirror (M₁). The angle between the two mirrors will be : (1) 90° (2) 45° (3) 75° (4) 60°



Assuming angles between two mirrors be θ as per geometry, sum of anlges of Δ $3\theta = 180^{\circ}$ $\theta = 60^{\circ}$

2. In a Young's double slit experiment, the slits are placed 0.320 mm apart. Light of wavelength λ = 500 nm is incident on the slits. The total number of bright fringes that are observed in the angular range $-30^{\circ} \le \theta \le 30^{\circ}$ is:

(3) 321

(4) 640

(2) 641

Ans. (2)

E

(1) 320



Pam difference $d\sin\theta = n\lambda$ where d = seperation of slits λ = wave length n = no. of maximas $0.32 \times 10^{-3} \sin 30 = n \times 500 \times 10^{-9}$ n = 320 Hence total no. of maximas observed in angular range $-30^{\circ} \le \theta \le 30^{\circ}$ is maximas = 320 + 1 + 320 = 641

3. At a given instant, say t = 0, two radioactive substances A and B have equal activities. The ratio $\frac{R_B}{R_A}$ of their activities after time t itself

is
$$m_2$$
, the half-life of B is :

(1)
$$\frac{ln2}{2}$$
 (2) $2ln2$ (3) $\frac{ln2}{4}$ (4) $4ln2$

Ans. (3)

Sol. Half life of A = ln2

$$t_{1/2} = \frac{\ell n 2}{\lambda}$$

$$\lambda_A = 1$$
at $t = 0$ $R_A = R_B$
 $N_A e^{-\lambda AT} = N_B e^{-\lambda BT}$
 $N_A = N_B$ at $t = 0$
at $t = t$
 $\frac{R_B}{R_A} = \frac{N_0 e^{-\lambda_B t}}{N_0 e^{-\lambda_A t}}$
 $e^{-(\lambda_B - \lambda_A)t} = e^{-t}$
 $\lambda_B - \lambda_A = 3$
 $\lambda_B = 3 + \lambda_A = 4$
 $t_{1/2} = \frac{\ell n 2}{\lambda_B} = \frac{\ell n 2}{4}$

4. Ge and Si diodes start conducting at 0.3 V and 0.7 V respectively. In the following figure if Ge diode connection are reversed, the value of V_o changes by : (assume that the Ge diode has large breakdown voltage)





Sol. Initially Ge & Si are both forward biased so current will effectivily pass through Ge diode with a drop of 0.3 V

if "Ge" is revesed then current will flow through "Si" diode hence an effective drop of (0.7 - 0.3)= 0.4 V is observed.

5. A rod of mass 'M' and length '2L' is suspended at its middle by a wire. It exhibits torsional oscillations; If two masses each of 'm' are attached at distance 'L/2' from its centre on both sides, it reduces the oscillation frequency by 20%. The value of ratio m/M is close to :

(1) 0.17 (2) 0.37 (3) 0.57 (4) 0.77Ans. (2)

Sol. Frequency of torsonal oscillations is given by \mathbf{k}

$$f = \frac{n}{\sqrt{I}}$$

$$f_1 = \frac{k}{\sqrt{\frac{M(2L)^2}{12}}}$$

$$f_2 = \frac{k}{\sqrt{\frac{M(2L)^2}{12} + 2m\left(\frac{L}{2}\right)^2}}$$

$$f_2 = 0.8 f_1$$

$$\frac{m}{M} = 0.375$$

6. A 15 g mass of nitrogen gas is enclosed in a vessel at a temperature 27°C. Amount of heat transferred to the gas, so that rms velocity of molecules is doubled, is about :

[Take R = 8.3 J/ K mole]

(1) 10 kJ (2) 0.9 kJ (3) 6 kJ (4) 14 kJ Ans. (1)

Sol. $Q = nC_v\Delta T$ as gas in closed vessel

$$Q = \frac{15}{28} \times \frac{5 \times R}{2} \times (4T - T)$$
$$Q = 10000 \text{ J} = 10 \text{ kJ}$$

collegebatch.com

A particle is executing simple harmonic motion (SHM) of amplitude A, along the x-axis, about x = 0. When its potential Energy (PE) equals kinetic energy (KE), the position of the particle will be :

(1)
$$\frac{A}{2}$$
 (2) $\frac{A}{2\sqrt{2}}$ (3) $\frac{A}{\sqrt{2}}$ (4) A

Ans. (3)

E

Sol. Potential energy (U) = $\frac{1}{2}kx^2$

Kinetic energy (K) =
$$\frac{1}{2}kA^2 - \frac{1}{2}kx^2$$

According to the question, U = k

:
$$\frac{1}{2}kx^2 = \frac{1}{2}kA^2 - \frac{1}{2}kx^2$$

$$\mathbf{x} = \pm \frac{\mathbf{A}}{\sqrt{2}}$$

 \therefore Correct answer is (3)

A musician using an open flute of length 50 cm produces second harmonic sound waves. A person runs towards the musician from another end of a hall at a speed of 10 km/h. If the wave speed is 330 m/s, the frequency heard by the running person shall be close to :

8.

Sol. Frequency of the sound produced by flute,

$$f = 2\left(\frac{v}{2\ell}\right) = \frac{2 \times 330}{2 \times 0.5} = 660 \text{Hz}$$

Velocity of observer, $v_0 = 10 \times \frac{5}{18} = \frac{25}{9}$ m/s

 \therefore frequency detected by observer, f' =

$$\begin{bmatrix} \frac{\mathbf{v} + \mathbf{v}_0}{\mathbf{v}} \end{bmatrix} \mathbf{f}$$

$$\therefore \mathbf{f} = \begin{bmatrix} \frac{25}{9} + 330}{330} \end{bmatrix} \mathbf{660}$$

= 335.56 × 2 = 671.12 ∴ closest answer is (4)

9. In a communication system operating at wavelength 800 nm, only one percent of source frequency is available as signal bandwidth. The number of channels accomodated for transmitting TV signals of band width 6 MHz are (Take velocity of light c = 3 × 10⁸m/s,h = 6.6 × 10⁻³⁴ J-s) (1) 3.75 × 10⁶ (2) 4.87 × 10⁵ (3) 3.86 × 10⁶ (4) 6.25 × 10⁵
Ans. (4)

2

Sol.
$$f = \frac{3 \times 10^8}{8 \times 10^{-7}} = \frac{30}{8} \times 10^{14} \text{ Hz}$$

= 3.75 × 10¹⁴ Hz
1% of f = 0.0375 × 10¹⁴ Hz
= 3.75 × 10¹² Hz = 3.75 × 10⁶ MHz
number of channels = $\frac{3.75 \times 10^6}{6}$ = 6.25 × 10⁵
 \therefore correct answer is (4)
10. Two point charges $q_1(\sqrt{10} \,\mu\text{C})$ and $q_2(-25 \,\mu\text{C})$
are placed on the x-axis at x = 1 m and x = 4 m
respectively. The electric field (in V/m) at a
point y = 3 m on y-axis is,
 $\left[\text{take} \frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \,\text{Nm}^2 \text{C}^{-2} \right]$
(1) $(-63\hat{i} + 27\hat{j}) \times 10^2$ (2) $(81\hat{i} - 81\hat{j}) \times 10^2$
(3) $(63\hat{i} - 27\hat{j}) \times 10^2$ (4) $(-81\hat{i} + 81\hat{j}) \times 10^2$
Ans. (3)
 $\int_{y=3}^{E_1} \frac{\theta_2}{\sqrt{10}} \frac{\xi_2}{\sqrt{x-3}} \frac{\xi_3}{x-4m} \frac{\xi_4}{x-4m} \frac{\xi_5}{x}$
Let $\vec{E}_1 \& \vec{E}_2$ are the vaues of electric field due
to $q_1 \& q_2$ respectively magnitude of
 $E_2 = \frac{1}{4\pi\epsilon_0} \frac{q_2}{r^2}$
 $E_2 = \frac{9 \times 10^9 \times (25) \times 10^{-6}}{(4^2 + 3^2)} \,\text{V/m}$
 $E_2 = 9 \times 10^3 \,\text{V/m}$

$$\therefore \vec{E}_{2} = 9 \times 10^{3} \left(\cos \theta_{2} \hat{i} - \sin \theta_{2} \hat{j} \right)$$

$$\therefore \tan \theta_{2} = \frac{3}{4}$$

$$\therefore \vec{E}_{2} = 9 \times 10^{3} \left(\frac{4}{5} \hat{i} - \frac{3}{5} \hat{j} \right) = \left(72 \hat{i} - 54 \hat{j} \right) \times 10^{2}$$

Magnitude of $E_{1} = \frac{1}{4\pi \in_{0}} \frac{\sqrt{10} \times 10^{-6}}{(1^{2} + 3^{2})}$

$$= \left(9 \times 10^{9} \right) \times \sqrt{10} \times 10^{-7}$$

$$= 9\sqrt{10} \times 10^{2}$$

$$\therefore \vec{E}_{1} = 9\sqrt{10} \times 10^{2} \left[\cos \theta_{1} \left(-\hat{i} \right) + \sin \theta_{1} \hat{j} \right]$$

$$\therefore \tan \theta_{1} = 3$$

$$3 \underbrace{\int \sqrt{10}}_{\theta_{1}} \underbrace{\int \frac{1}{\sqrt{10}} \left(-\hat{i} \right) + \frac{3}{\sqrt{10}} \hat{j} \right]}_{\theta_{1}}$$

$$E_{1} = 9 \times \sqrt{10} \times 10^{2} \left[\frac{1}{\sqrt{10}} \left(-\hat{i} \right) + \frac{3}{\sqrt{10}} \hat{j} \right]$$

$$E_{1} = 9 \times 10^{2} \left[-\hat{i} + 3\hat{j} \right] = \left[-9\hat{i} + 27\hat{j} \right] 10^{2}$$

$$\therefore \vec{E} = \vec{E}_{1} + \vec{E}_{2} = \left(63\hat{i} - 27\hat{j} \right) \times 10^{2} \text{ V/m}$$

$$\therefore \text{ correct answer is (3)}$$

A parallel plate capacitor with square plates is filled with four dielectrics of dielectric constants K₁, K₂, K₃, K₄ arranged as shown in the figure. The effective dielectric constant K will be :

$$K_{1} = \frac{K_{1}}{K_{2}} = \frac{K_{1}}{K_{3}} = \frac{K_{4}}{K_{4}} = \frac{K_{1}}{K_{2}}$$

$$(1) \quad K = \frac{(K_{1} + K_{2})(K_{3} + K_{4})}{2(K_{1} + K_{2} + K_{3} + K_{4})}$$

$$(2) \quad K = \frac{(K_{1} + K_{2})(K_{3} + K_{4})}{(K_{1} + K_{2} + K_{3} + K_{4})}$$

$$(3) \quad K = \frac{(K_{1} + K_{4})(K_{2} + K_{3})}{2(K_{1} + K_{2} + K_{3} + K_{4})}$$

$$(4) \quad K = \frac{(K_{1} + K_{3})(K_{2} + K_{4})}{K_{1} + K_{2} + K_{3} + K_{4}}$$

Ans. (Bonus)



Е



Now if
$$k_{eq} = k$$
, $C_{eq} = \frac{k \in_0 L^2}{d}$ (ii)

on comparing equation (i) to equation (ii), we get

$$k_{eq} = \frac{k_1 k_2 (k_3 + k_4) + k_3 k_4 (k_1 + k_2)}{(k_1 + k_2) (k_3 + k_4)}$$

This does not match with any of the options so probably they have assumed the wrong combination

$$C_{13} = \frac{k_1 \in_0 L \frac{L}{2}}{d/2} + k_3 \in_0 \frac{L \frac{L}{2}}{d/2}$$
$$= (k_1 + k_3) \frac{\in_0 L^2}{d}$$
$$C_{24} = (k_2 + k_4) \frac{\in_0 L^2}{d}$$

$$C_{eq} = \frac{C_{13}C_{24}}{C_{13}C_{24}} = \frac{(k_1 + k_3)(k_2 + k_4)}{(k_1 + k_2 + k_3 + k_4)} \frac{\epsilon_0 L^2}{d}$$
$$= \frac{k \epsilon_0 L^2}{d}$$
$$k = \frac{(k_1 + k_3)(k_2 + k_4)}{(k_1 + k_2 + k_3 + k_4)}$$

However this is one of the four options. It must be a "Bonus" logically but of the given options probably they might go with (4)

12. A rod of length 50cm is pivoted at one end. It is raised such that if makes an angle of 30° from the horizontal as shown and released from rest. Its angular speed when it passes through the horizontal (in rad s⁻¹) will be (g = 10ms^{-2})



Work done by gravity from initial to final position is,

$$W = mg\frac{\ell}{2}\sin 30^{\circ}$$

 $\frac{\mathrm{mg}\ell}{4}$

According to work energy theorem

 $W = \frac{1}{2}I\omega^2$



Е
$$\Rightarrow \frac{1}{2} \frac{m\ell^2}{3} \omega^2 = \frac{mg\ell}{4}$$
$$\omega = \sqrt{\frac{3g}{2\ell}} = \sqrt{\frac{3 \times 10}{2 \times 0.5}}$$
$$\omega = \sqrt{30} \text{ rad/sec}$$

 \therefore correct answer is (1)

13. One of the two identical conducting wires of length L is bent in the form of a circular loop and the other one into a circular coil of N identical turns. If the same current is passed in both, the ratio of the magnetic field at the central of the loop (B_L) to that at the centre of B.

the coil (B_C), i.e. R
$$\frac{B_L}{B_C}$$
 will be :
(1) $\frac{1}{N}$ (2) N² (3) $\frac{1}{N^2}$ (4) N

Ans. (3)

Sol.
Loop
R

$$L = 2\pi R$$
 $L = N \times 2\pi r$
 $R = Nr$
 $B_L = \frac{\mu_0 i}{2R}$ $B_C = \frac{\mu_0 N i}{2r}$
 $B_C = \frac{\mu_0 N^2 i}{2R}$
 $\frac{B_L}{B_C} = \frac{1}{N^2}$

14. The energy required to take a satellite to a height 'h' above Earth surface (radius of Earth = 6.4×10^3 km) is E₁ and kinetic energy required for the satellite to be in a circular orbit at this height is E₂. The value of h for which E₁ and E₂ are equal, is:

(1) 1.28×10^4 km	(2) 6.4×10^3 km
(3) 3.2×10^3 km	(4) 1.6×10^3 km

Ans. (3)

Ε

Sol. $U_{surface} + E_1 = U_h$

KE of satelite is zero at earth surface & at height h

$$-\frac{GM_{e}m}{R_{e}} + E_{1} = -\frac{GM_{e}m}{(Re+h)}$$
$$E_{1} = GM_{e}m\left(\frac{1}{R_{e}} - \frac{1}{R_{e}+h}\right)$$
$$E_{1} = \frac{GM_{e}m}{(R_{e}+h)} \times \frac{h}{R_{e}}$$

Gravitational attraction $F_G = ma_C = \frac{mv^2}{(R_e + h)}$

$$E_{2} \Rightarrow \frac{mv^{2}}{(R_{e} + h)} = \frac{GM_{e}m}{(R_{e} + h)^{2}}$$
$$mv^{2} = \frac{GM_{e}m}{(R_{e} + h)}$$
$$E_{2} = \frac{mv^{2}}{2} = \frac{GM_{e}m}{2(R_{e} + h)}$$
$$E_{1} = E_{2}$$
$$\frac{h}{R_{e}} = \frac{1}{2} \Rightarrow h = \frac{R_{e}}{2} = 3200 \text{km}$$

15. The energy associated with electric field is (U_E) and with magnetic field is (U_B) for an electromagnetic wave in free space. Then :

(1)
$$U_E = \frac{U_B}{2}$$
 (2) $U_E < U_B$
(3) $U_E = U_B$ (4) $U_E > U_B$

Ans. (3)

Sol. Average energy density of magnetic field,

$$u_{\rm B} = \frac{B_0^2}{2\mu_0}$$
, B_0 is maximum value of magnetic

field.

Average energy density of electric field,

$$u_{\rm E} = \frac{\varepsilon_0 \in_0^2}{2}$$

now,
$$\epsilon_0 = CB_0$$
 , $C^2 = \frac{1}{\mu_0 \epsilon_0}$

$$\mathbf{u}_{\mathrm{E}} = \frac{\epsilon_0}{2} \times \mathrm{C}^2 \mathrm{B}_{\mathrm{C}}^2$$

5

collegebatch.com

$$= \frac{\epsilon_0}{2} \times \frac{1}{\mu_0 \epsilon_0} \times B_0^2 = \frac{B_0^2}{2\mu_0} = u_{\rm E}$$

 $u_E = u_B$

since energy density of electric & magnetic field is same, energy associated with equal volume will be equal.

- $u_{\rm E} = u_{\rm B}$
- 16. A series AC circuit containing an inductor (20 mH), a capacitor (120 μ F) and a resistor (60 Ω) is driven by an AC source of 24 V/50 Hz. The energy dissipated in the circuit in 60 s is :
 - (1) 2.26 × 10³ J (2) 3.39×10^3 J
 - (3) 5.65×10^2 J (4) 5.17×10^2 J
- Ans. (4)

Sol. $R = 60\Omega$ f = 50Hz, $\omega = 2\pi f = 100 \pi$

$$x_{C} = \frac{1}{\omega C} = \frac{1}{100\pi \times 120 \times 10^{-6}}$$

$$x_{C} = 26.52 \ \Omega$$

$$x_{L} = \omega L = 100\pi \times 20 \times 10^{-3} = 2\pi\Omega$$

$$x_{C} - x_{L} = 20.24 \approx 20$$

$$x_{C} - x_{L} = 20\Omega$$

$$z = \sqrt{R^{2} + (x_{C} - x_{L})^{2}}$$

$$z = 20\sqrt{10}\Omega$$

$$\cos\phi = \frac{R}{z} = \frac{3}{\sqrt{10}}$$

$$P_{avg} = VI \cos\phi, I = \frac{V}{z}$$

$$= \frac{V^{2}}{z} \cos\phi$$

$$= 8.64 \text{ watt}$$

$$Q = P.t = 8.64 \times 60 = 5.18 \times 10^{2}$$

17. Expression for time in terms of G (universal gravitational constant), h (Planck constant) and c (speed of light) is proportional to :

(1)
$$\sqrt{\frac{Gh}{c^3}}$$
 (2) $\sqrt{\frac{hc^5}{G}}$
(3) $\sqrt{\frac{c^3}{Gh}}$ (4) $\sqrt{\frac{Gh}{c^5}}$

Ans. (4)

Sol.
$$F = \frac{GM^2}{R^2} \Rightarrow G = [M^{-1}L^3T^{-2}]$$

$$E = hv \Rightarrow h = [ML^2T^{-1}]$$

$$C = [LT^{-1}]$$

$$t \propto G^x h^y C^z$$

$$[T] = [M^{-1}L^3T^{-2}]^x [ML^2T^{-1}]^y [LT^{-1}]^z$$

$$[M^0L^0T^1] = [M^{-x} + yL^{3x} + 2y + zT^{-2x} - y - z]$$
on comparing the powers of M, L, T
$$-x + y = 0 \Rightarrow x = y$$

$$3x + 2y + z = 0 \Rightarrow 5x + z = 0 \qquad \dots(i)$$

$$-2x - y - z = 1 \Rightarrow 3x + z = -1 \qquad \dots(i)$$
on solving (i) & (ii) $x = y = \frac{1}{2}, z = -\frac{5}{2}$

$$t \propto \sqrt{\frac{Gh}{C^5}}$$
18. The magnetic field associated with a light wave is given, at the origin, by
$$B = B_y [\sin(3 + 4x + 10^7)ct + \sin(6 + 28x + 10^7)ct]$$

is given, at the origin, by $B = B_0 [sin(3.14 \times 10^7)ct + sin(6.28 \times 10^7)ct]$. If this light falls on a silver plate having a work function of 4.7 eV, what will be the maximum kinetic energy of the photo electrons ? (c = 3 × 10⁸ms⁻¹, h = 6.6 × 10⁻³⁴ J-s)

$$\begin{array}{c} (2 - 3 \times 10^{-1} \text{ ms}^{-1}, 11 - 0.0 \times 10^{-1} \text{ s}^{-3}) \\ (1) \ 7.72 \ \text{eV} \\ (3) \ 12.5 \ \text{eV} \\ \end{array} \qquad \begin{array}{c} (2) \ 8.52 \ \text{eV} \\ (4) \ 6.82 \ \text{eV} \end{array}$$

Ans. (1)

Sol. $B = B_0 \sin (\pi \times 10^7 C)t + B_0 \sin (2\pi \times 10^7 C)t$ since there are two EM waves with different frequency, to get maximum kinetic energy we take the photon with higher frequency

$$B_1 = B_0 \sin(\pi \times 10^7 C)t$$
 $v_1 = \frac{10^7}{2} \times C$

$$\begin{split} B_2 &= B_0 \sin(2\pi \times 10^7 \text{C}) \text{t } \nu_2 = 10^7 \text{C} \\ \text{where C is speed of light } C &= 3 \times 10^8 \text{ m/s} \\ \nu_2 &> \nu_1 \\ \text{so KE of photoelectron will be maximum for } \\ \text{photon of higher energy.} \\ \nu_2 &= 10^7 \text{C Hz} \\ \text{h}\nu &= \phi + \text{KE}_{\text{max}} \\ \text{energy of photon} \\ E_{\text{ph}} &= \text{h}\nu = 6.6 \times 10^{-34} \times 10^7 \times 3 \times 10^9 \\ E_{\text{ph}} &= 6.6 \times 3 \times 10^{-19} \text{J} \\ &= \frac{6.6 \times 3 \times 10^{-19}}{1.6 \times 10^{-19}} \text{eV} = 12.375 \text{eV} \\ \text{KE}_{\text{max}} &= E_{\text{ph}} - \phi \\ &= 12.375 - 4.7 = 7.675 \text{ eV} \approx 7.7 \text{ eV} \end{split}$$



19. Charge is distributed within a sphere of radius R with a volume charge density $\rho(r) = \frac{A}{r^2}e^{-2r/a}$, where A and a are constants. If Q is the total charge of this charge distribution, the radius R is :

(1)
$$\frac{a}{2}\log\left(1-\frac{Q}{2\pi aA}\right)$$
 (2) $a\log\left(1-\frac{Q}{2\pi aA}\right)$
(3) $a\log\left(\frac{1}{1-\frac{Q}{2\pi aA}}\right)$ (4) $\frac{a}{2}\log\left(\frac{1}{1-\frac{Q}{2\pi aA}}\right)$

Ans. (4)

Sol.

$$Q = \int \rho dv$$

$$= \int_{0}^{R} \frac{A}{r^{2}} e^{-2r/a} (4\pi r^{2} dr)$$

$$= \int_{0}^{R} \frac{A}{r^{2}} e^{-2r/a} (4\pi r^{2} dr)$$

$$= 4\pi A \int_{0}^{R} e^{-2r/a} dr$$

$$= 4\pi A \left(\frac{e^{-2r/a}}{-\frac{2}{a}}\right)_{0}^{R}$$

$$= 4\pi A \left(-\frac{a}{2}\right) (e^{-2R/a} - 1)$$

$$Q = 2\pi a A (1 - e^{-2R/a})$$

$$R = \frac{a}{2} \log \left(\frac{1}{1 - \frac{Q}{2\pi a A}}\right)$$

20. Two Carrnot engines A and B are operated in series. The first one, A, receives heat at T_1 (= 600 K) and rejects to a reservoir at temperature T_2 . The second engine B receives heat rejected by the first engine and, in turn, rejects to a heat reservoir at T_3 (= 400 K). Calculate the temperature T_2 if the work outputs of the two engines are equal : (1) 400 K (2) 600 K (3) 500 K (4) 300 K

Ans. (3)

Ε





- 23. The position co-ordinates of a particle moving in a 3-D coordinate system is given by $x = a \cos \omega t$ $y = a \sin \omega t$ and $z = a\omega t$ The speed of the particle is : (2) $\sqrt{3}$ a ω (1) $a\omega$ (3) $\sqrt{2}$ a ω $(4) 2a\omega$ Ans. (3) **Sol.** $v_x = -a\omega sin\omega t \implies v_y = a\omega cos\omega t$ \Rightarrow v = $\sqrt{v_x^2 + v_y^2 + v_z^2}$ $v_z = a\omega$ $v = \sqrt{2}a\omega$ 24. In the given circuit the internal resistance of the 18 V cell is negligible. If $R_1 = 400 \Omega$, $R_3 = 100 \Omega$ and $R_4 = 500 \Omega$ and the reading of an ideal voltmeter across R_4 is 5V, then the value R_2 will be : R R. 18V (1) 300 Ω (2) 230 Ω (3) 450 Ω (4) 550 Ω Ans. (1) $R_{3}=100\Omega$ R₄=500Ω ₩₩ ₩₩ $R_1 = 400\Omega$ Ŵ i, R_2 Sol. 18V $V_4 = 5V$ $i_1 = \frac{V_4}{R_4} = 0.01 \text{ A}$ $V_3 = i_1 R_3 = 1V$ $V_3 + V_4 = 6V = V_2$ $V_1 + V_3 + V_4 = 18V$ $V_1 = 12 V$ $i = \frac{V_1}{R_1} = 0.03$ Amp. K_1 $i_2 = 0.02$ Amp $V_2 = 6V$ $R_2 = \frac{V_2}{i_2} = \frac{6}{0.02} = 300\Omega$
 - **25.** A mass of 10 kg is suspended vertically by a rope from the roof. When a horizontal force is applied on the rope at some point, the rope deviated at an angle of 45° at the roof point. If the suspended mass is at equilibrium, the magnitude of the force applied is (g = 10 ms⁻²)

(1) 200 N (2) 100 N (3) 140 N (4) 70 N Ans. (2)



at equation

$$\tan 45^\circ = \frac{100}{\mathrm{F}}$$

$$F = 100 N$$

26. In a car race on straight road, car A takes a time t less than car B at the finish and passes finishing point with a speed 'v' more than that of car B. Both the cars start from rest and travel with constant acceleration a_1 and a_2 respectively. Then 'v' is equal to

(1)
$$\frac{a_1 + a_2}{2}t$$
 (2) $\sqrt{2a_1a_2}t$
(3) $\frac{2a_1a_2}{a_1 + a_2}t$ (4) $\sqrt{a_1a_2}t$

Ans. (4)

Sol. For A & B let time taken by A is t_0

$$\begin{array}{c} & X \\ & & \\ & & \\ u = 0 \\ & & v_A = a_1 t_0 \\ & v_B = a_2 (t_0 + t) \end{array}$$

from ques.

$$v_{A} - v_{B} = v = (a_{1} - a_{2})t_{0} - a_{2}t \qquad \dots(i)$$

$$x_{B} = x_{A} = \frac{1}{2}a_{1}t_{0}^{2} = \frac{1}{2}a_{2}(t_{0} + t)^{2}$$

$$\Rightarrow \sqrt{a_{1}}t_{0} = \sqrt{a_{2}}(t_{0} + t)$$

$$\Rightarrow (\sqrt{a_{2}} - \sqrt{a_{2}})t_{0} = \sqrt{a_{2}}t \qquad \dots(ii)$$

8



putting t_0 in equation

$$v = (a_1 - a_2) \frac{\sqrt{a_2}t}{\sqrt{a_1} - \sqrt{a_2}} - a_2 t$$
$$= \left(\sqrt{a_1} + \sqrt{a_2}\right) \sqrt{a_2}t - a_2 t \implies v = \sqrt{a_1 a_2}t$$
$$\Rightarrow \sqrt{a_1 a_2}t + a_2 t - a_2 t$$

- 27. A power transmission line feeds input power at 2300 V to a step down transformer with its primary windings having 4000 turns. The output power is delivered at 230 V bv the transformer. If the current in the primary of the transformer is 5A and its efficiency is 90%, the output current would be :
 - (1) 25 A (2) 50 A (3) 35 A (4) 45 A
- Ans. (4)

Sol.
$$\eta = \frac{P_{out}}{P_{in}} = \frac{V_s I_s}{V_p I_p}$$

 $\Rightarrow 0.9 = \frac{23 \times I_s}{230 \times 5}$
 $\Rightarrow I_s = 45A$

28. The top of a water tank is open to air and its water level is maintained. It is giving out 0.74 m³ water per minute through a circular opening of 2 cm radius in its wall. The depth of the centre of the opening from the level of water in the tank is close to :

(1) 9.6 m (2) 4.8 m (3) 2.9 m (4) 6.0 m Ans. (2)

Sol. In flow volume = outflow volume

$$\Rightarrow \frac{0.74}{60} = (\pi \times 4 \times 10^{-4}) \times \sqrt{2gh}$$
$$\Rightarrow \sqrt{2gh} = \frac{74 \times 100}{240\pi}$$
$$\Rightarrow \sqrt{2gh} = \frac{740}{24\pi}$$
$$\Rightarrow 2gh = \frac{740 \times 740}{24 \times 24 \times 10} (\pi^2 = 10)$$
$$\Rightarrow h = \frac{74 \times 74}{2 \times 24 \times 24}$$
$$\Rightarrow h \approx 4.8m$$

29. The pitch and the number of divisions, on the circular scale, for a given screw gauge are 0.5 mm and 100 respectively. When the screw gauge is fully tightened without any object, the zero of its circular scale lies 3 divisions below the mean line. The readings of the main scale and the circular scale, for a thin sheet, are 5.5 mm and 48 respectively, the thickness of this sheet is :

(1) 5.755 m
(2) 5.725 mm
(3) 5.740 m
(4) 5.950 mm

Ans. (2)

Sol. LC =
$$\frac{\text{Pitch}}{\text{No. of division}}$$

LC = 0.5×10^{-2} mm +ve error = $3 \times 0.5 \times 10^{-2}$ mm = 1.5×10^{-2} mm = 0.015 mm Reading = MSR + CSR - (+ve error) = 5.5 mm + ($48 \times 0.5 \times 10^{-2}$) - 0.015= 5.5 + 0.24 - 0.015 = 5.725 mm

30. A particle having the same charge as of electron moves in a circular path of radius 0.5 cm under the influence of a magnetic field of 0.5 T. If an electric field of 100 V/m makes it to move in a straight path, then the mass of the particle is (Given charge of electron =1.6 × 10⁻¹⁹C) (1) 2.0 × 10⁻²⁴ kg
(2) 1.6 × 10⁻¹⁹ kg
(3) 1.6 × 10⁻²⁷ kg

(4)
$$9.1 \times 10^{-31}$$
 kg

Ans. (1)

Sol.
$$\frac{mv^2}{R} = qvB$$

 $mv = qBR \dots(i)$ Path is straight line it qE = qvB E = vB \dots(ii) From equation (i) & (ii)

$$m = \frac{qB^2R}{E}$$
$$m = 2.0 \times 10^{-24} \text{ kg}$$

9



E

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Wednesday 09th JANUARY, 2019) TIME : 2 : 30 PM To 05 : 30 PM CHEMISTRY

1. lood reducing nature of H₃PO₂ ttributed to the 5. The major product of the following reaction is: presence of: OH (1) One P-OH bond (2) One P-H bond CH₃ (3) Two P-H bonds (4) Two P-OH bonds $AlCl_3,\Delta$ Ans. (3) **Sol.** H_3PO_2 is good reducing agent due to presence OH H₂C of two P-H bonds. (1)2. The complex that has highest cry splitting energy (Δ), is : (1) $K_3[Co(CN)_6]$ CH₃ (2) $[Co(NH_3)_5(H_2O)]Cl_3$ (3) $K_2[CoCl_4]$ (2)0: (4) $[Co(NH_3)_5Cl]Cl_2$ Ans. (1) **Sol.** As complex $K_3[Co(CN)_6]$ have CN^- ligand CH₃ which is strongfield ligand amongst the given ligands in other complexes. 3. The metal that forms nitride by reacting directly (3)0 with N_2 of air, is : (1) K (2) Cs (3) Li (4) Rb Ans. (3) OH Sol. Only Li react directly with N_2 out of alkali HC3 metals $6Li + N_2 \rightarrow 2Li_3N$ (4)4. In which of the following processes, the bond order has increased and paramagnetic character has changed to diamagnetic ? Ans. (4) (1) $N_2 \rightarrow N_2^+$ (2) NO \rightarrow NO⁺ Sol. (3) $O_2 \rightarrow O_2^{2-}$ (4) $O_2 \rightarrow O_2^+$ CH₂ Ans. (2) CH₃ Sol. Changein Bond Order OH Process magnetic Change nature $N_2 \rightarrow N_2^+$ $Dia \rightarrow para$ $3 \rightarrow 2.5$ $NO \rightarrow NO^+$ Para \rightarrow Dia $2.5 \rightarrow 3$ CH₃ $O_2 \rightarrow O_2^{-2}$ Para \rightarrow Dia $2 \rightarrow 1$ $O_2 \rightarrow O_2^+$ $Para \rightarrow Para$ $2 \rightarrow 2.5$





- 6. The transition element that has lowest enthalpy of atomisation, is :
 - (1) Zn
 - (2) Cu
 - (3) V
 - (4) Fc
- Ans. (2)
- **Sol.** Since Zn is not a transition element so transition element having lowest atomisation energy out of Cu, V, Fe is Cu.
- 7. Which of the following combination of statements is true regarding the interpretation of the atomic orbitals ?
 - (a) An electron in an orbital of high angular momentum stays away from the nucleus than an electron in the orbital of lower angular momentum.
 - (b) For a given value of the principal quantum number, the size of the orbit is inversely proportional to the azimuthal quantum number.
 - (c) According to wave mechanics, the ground

state angular momentum is h equal to $\frac{h}{2\pi}$.

- (d) The plot of ψ Vs r for various azimuthal quantum numbers, shows peak shifting towards higher r value.
- (1) (b), (c) (2) (a), (d) (3) (a), (b) (4) (a), (c)

Ans. (4)

- Sol. Refer Theory
- 8. The tests performed on compound X and their inferences are:
 - Test Inference (a) 2,4 - DNP test Coloured precipitate (b) Iodoform test Yellow precipitate (c) Azo-dye test No dye formation



Ans. (2)

 \rightarrow 2,4 – DNP test is given by aldehyde on Sol. ketone \rightarrow Iodoform test is given by compound having

$$CH_3 - C - group.$$

9. The major product formed in the following reaction is:

2













Sol. Aldehyde reacts at a faster rate than keton during aldol and stericall less hindered anion will be a better nucleophile so sefl aldol at

 $CH_3 - C - H$ will be the major product.

10. For the reaction, 2A + B → products, when the concentrations of A and B both wrere doubled, the rate of the reaction increased from 0.3 mol L⁻¹s⁻¹ to 2.4 mol L⁻¹s⁻¹. When the concentration of A alone is doubled, the rate increased from 0.3 mol L⁻¹s⁻¹ to0.6 mol L⁻¹s⁻¹

Which one of the following statements is correct ?

- (1) Order of the reaction with respect to Bis2
- (2) Order of the reaction with respect to Ais2
- (3) Total order of the reaction is 4
- (4) Order of the reaction with respect to B is 1

Ans. (1)

Ε

- Sol. $r = K[A]^{x}[B]^{y}$ $\Rightarrow 8 = 2^{3} = 2^{x+y}$ $\Rightarrow x + y = 3 ...(1)$ $\Rightarrow 2 = 2^{x}$ $\Rightarrow x = 1, y = 2$ Order w.r.t. A = 1Order w.r.t. B = 2
- **11.** The correct sequence of amino acids present in the tripeptide given below is :



(1) Leu - Ser - Thr
(2) Thr - Ser - Leu
(3) Thr - Ser - Val
(4) Val - Ser - Thr
Ans. (4)

Sol. Leusine $\rightarrow \begin{array}{c} CH-COOH \\ I \\ NH_2 \end{array}$

Serine

NO- CH_2 - CH - COOHNH₂

Threenine $H_3C-CH-CH-COOH$ I IOH NH_2

12. The correct statement regarding the given Ellingham diagram is:





- At 800°C, Cu can be used for the extraction of Zn from ZnO
- (2) At 500 C, coke can be used for the extraction of Zn from ZnO
- Coke cannot be used for the extraction of Cu from Ca₂O.
- (4) At 1400°C, Al can be used for the extraction of Zn from ZnO
- Ans. (4)
- **Sol.** According to the given diagram Al can reduce ZnO.

 $3ZnO+2Al \rightarrow 3Zn+Al_2O_3$

- 13. For the following reaction, the mass of water produced from 445 g of $C_{57}H_{110}O_6$ is :
- $2C_{57}H_{110}O_6(s) + 163O_2(g) \rightarrow 114CO_2(g) + 110 H_2OP(1)$ (1) 495 g (2) 490 g (3) 890 g (4) 445 g **Ans. (1)**

(1)

Sol. moles of $C_{57}H_{110}O_6(s) = \frac{445}{890} = 0.5$ moles

 $2\mathrm{C}_{57}\mathrm{H}_{110}\mathrm{O}_{6}(\mathrm{s}) + 163~\mathrm{O}_{2}(\mathrm{g}) \rightarrow 114~\mathrm{CO}_{2}(\mathrm{g}) + 110~\mathrm{H}_{2}\mathrm{O}(l)$

$$n_{\rm H_2O} = \frac{110}{4} = \frac{55}{2}$$

$$m_{H_{2}O} = \frac{55}{2} \times 18$$

= 495gm

- 14. The correct match between Item I and Item II is :
- Item IItem II(A) Benzaldehyde(P) Mobile phase(B) Alumina(Q) Adsorbent(C) Acetonitrile(R) Adsorbate(1) (A) \rightarrow (Q);(B) \rightarrow (R);(C) \rightarrow (P)(2) (A) \rightarrow (P); (B) \rightarrow (R); (C) \rightarrow (Q)(3) (A) \rightarrow (Q); (B) \rightarrow (P); (C) \rightarrow (R)(4) (A) \rightarrow (R); (B) \rightarrow (Q); (C) \rightarrow (P)Ans. (4)

Sol.

15. The increasing basicity order of the following compounds is : (A) $CH_3CH_2NH_2$ (B) $\begin{array}{c} CH_2CH_3 \\ I \\ CH_3CH_2NH \end{array}$ (C) $\begin{array}{c} CH_3 \\ I \\ H_3C-N-CH_3 \end{array}$ (D) $\begin{array}{c} CH_3 \\ I \\ Ph-N-H \end{array}$ (1) (D)<(C)<(A)<(B) (2) (A)<(B)<(D)<(C) (3) (A)<(B)<(C)< (D) (4) (D)<(C)<(B)<(A) Ans. (1)

Sol.

$$\begin{array}{cccc} CH_3 & CH_3 & CH_2-CH_3 \\ I & I & I \\ Ph-N-H < CH_3-N-CH_3 < CH_3-CH_2-NH < CH_3-CH_2-NH_2 \\ \uparrow & \uparrow & \uparrow \\ Ione pair & more steric \\ delocalized & hinderence \\ less solutions \\ energy \end{array}$$

16. For coagulation of arscnious sulphide sol, which one of the following salt solution will be most effective ?

(1)
$$AlCl_3$$
 (2) $NaCl$
(3) $BaCl_2$ (4) Na_3PO_4

Ans. (1)

Sol. Sulphide is -ve charged colloid so cation with maximum charge will be most effective for coagulation.

 $Al^{3+} > Ba^{2+} > Na^+$ coagulating power.

At 100°C, copper (Cu) has FCC unit cell structure with cell edge length of x Å. What is the approximate density of Cu (in g cm⁻³) at this temperature ?

[Atomic Mass of Cu = 63.55u]

(1)
$$\frac{105}{x^3}$$
 (2) $\frac{211}{x^3}$ (3) $\frac{205}{x^3}$ (4) $\frac{422}{x^3}$

Ans. (4)

4

collegebatch.com

Sol. FCC unit cell Z = 4

$$d = \frac{63.5 \times 4}{6 \times 10^{23} \times x^3 \times 10^{-24}} \text{ g/cm}^3$$
$$d = \frac{63.5 \times 4 \times 10}{6} \text{ g/cm}^3$$
$$d = \frac{423.33}{x^3} \approx \left(\frac{422}{x^3}\right)$$

18. The major product obtained in the following reaction is :











Ans. (3)



19. Which of the following conditions in drinking water causes methemoglobinemia ? (1) > 50ppm of load (2) > 100 ppm of sulphate (3) > 50 ppm of chloride (4) > 50 ppm of nitrate Ans. (4) Sol. Concentration of nitrate >50 ppm in drinking water causes methemoglobinemia 20. Homoleptic octahedral complexes of a metal ion 'M^{3+'} with three monodentate ligands and L_1, L_2, L_3 absorb wavelengths in the region of green, blue and red respectively. The increasing order of the ligand strength is :

(1)
$$L_2 < L_1 < L_3$$

(2) $L_3 < L_2 < L_1$
(3) $L_3 < L_1 < L_2$
(4) $L_1 < L_2 < L_3$

Ans. (3)

Sol. Order of λ_{abs} - $L_3 > L_1 > L_2$

So Δ_0 order will be $L_2 > L_1 > L_3$ (as $\Delta_0 \propto \frac{1}{\lambda_{abs}}$)

So order of ligand strength will be $L_2 > L_1 > L_3$

21. The product formed in the reaction of cumene with O_2 followed by treatment with dil. HCl are :



Ans. (3)



Sol. Cummene hydroperoxide reaction



22. The temporary hardness of water is due to :-(1) Ca(HCO₃)₂(2) NaCl

(3) Na_2SO_4 (4) $CaCl_2$

Ans. (1)

- Sol. $Ca(HCO_3)_2$ is reponsible for temporary hardness of water
- 23. The entropy change associated with the conversion of 1 kg of ice at 273 K to water vapours at 383 K is :

(Specific heat of water liquid and water vapour are 4.2 kJ K⁻¹ kg⁻¹ and 2.0 kJ K⁻¹ kg⁻¹; heat of liquid fusion and vapourisation of water are 344 kJ kg⁻¹ and 2491 kJ kg⁻¹, respectively). (log 273 = 2.436, log 373 = 2.572, log 383 = 2.583) (1) 7.90 kJ kg⁻¹ K⁻¹ (2) 2.64 kJ kg⁻¹ K⁻¹ (3) 8.49 kJ kg⁻¹ K⁻¹ (4) 4.26 kJ kg⁻¹ K⁻¹

Ans. (4)

$$\Delta S_{1} = \frac{\Delta H_{\text{fusion}}}{273} = \frac{334}{273} = 1.22$$

$$\Delta S_{2} = 4.2 \ell N \left(\frac{363}{273}\right) = 1.31$$

$$\Delta S_{3} = \frac{\Delta H_{\text{vap}}}{373} = \frac{2491}{373} = 6.67$$

$$\Delta S_{4} = 2.0 \ell n \left(\frac{383}{373}\right) = 0.05$$

$$\Delta S_{\text{total}} = 9.26 \text{ kJ kg}^{-1} \text{ K}^{-1}$$

24. The pH of rain water, is approximately : (1) 6.5 (2) 7.5 (3) 5.6 (4) 7.0

Ans. (3)

- Sol. pH of rain water is approximate 5.6
- 25. If the standard electrode potential for a cell is2 V at 300 K, the equilibrium constant (K) for the reaction

 $Zn(s) + Cu^{2+}(aq) \implies Zn^{2+}(aq) + Cu(s)$

at 300 K is approximately.

$$(R = 8 JK^{-1} mol^{-1}, F = 96000 C mol^{-1})$$

(1)
$$e^{-160}$$
 (2) e^{-200}
(3) e^{-160} (4) e^{-80}

$$(\mathbf{J}) \in \mathbf{I}^{\mathbf{I}}$$

Ans. (1)

Sol.
$$\Delta G^{\circ} = -RT \ln k = -nFE_{cell}^{\circ}$$

 $\ln k = \frac{n \times F \times E^{\circ}}{R \times T} = \frac{2 \times 96000 \times 2}{8 \times 300}$
 $\ln k = 160$
 $k = e^{160}$

26. A solution containing 62 g ethylene glycol in 250 g water is cooled to -10° C. If K_f for water is 1.86 K kg mol⁻¹, the amount of water (in g) separated as ice is :

(1) 32 (2) 48 (3) 16 (4) 64

Sol.
$$\Delta T_f = K_f \cdot m$$

$$10 = 1.86 \times \frac{62/62}{W_{kg}}$$

W = 0.186 kg

- $\Delta W = (250 186) = 64 \text{ gm}$
- 27. When the first electron gain enthalpy $(\Delta_{eg}H)$ of oxygen is -141 kJ/mol, its second electron gain enthalpy is :
 - (1) almost the same as that of the first
 - (2) negative , but less negative than the first
 - (3) a positive value
 - (4) a more negative value than the first

Ans. (3)

Sol. Second electron gain enthalpy is always positive for every element.

 $O^{-}_{(g)}$ + e⁻ $\rightarrow O^{-2}_{(g)}$; ΔH = positive



Е

28. The major product of the following reaction is :



Ans. (3)



29. Which of the following compounds is not aromatic ?

Вr



Sol.

Do not have $(4n + 2) \pi$ electron It has $4n \pi$ electrons

So it is Anti aromatic.

30. Consider the following reversible chemical reactions :

$$A_2(g) + Br_2(g) \xrightarrow{K_1} 2AB(g) \dots (1)$$

 $\begin{array}{l} 6AB(g) \xrightarrow{K_2} 3A_2(g) + 3B_2(g) \ \dots \ (2) \end{array}$ The relation between K_1 and K_2 is : (1) $K_2 = K_1^3$ (2) $K_2 = K_1^{-3}$

(3)
$$K_1K_2 = 3$$
 (4) $K_1K_2 = \frac{1}{3}$

Ans. (2)

Sol.
$$A_2(g) + B_2(g) \xrightarrow{k_1} 2AB$$
 ...(1)
 \Rightarrow eq. (1) $\times 3$
 $6 AB(g) \xrightarrow{3} 3A_2(g) + 3B_2(g)$
 $\Rightarrow \left(\frac{1}{k_1}\right)^3 = k_2 \Rightarrow k_2 = (k_1)^{-3}$

collegebatch.com

Ε

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Thursday 10th JANUARY, 2019) TIME : 9 : 30 AM To 12 : 30 PM MATHEMATICS

 Consider a triangular plot ABC with sides AB=7m, BC=5m and CA=6m. A vertical lamp-post at the mid point D of AC subtends an angle 30° at B. The height (in m) of the lamp-post is:

(1)
$$7\sqrt{3}$$
 (2) $\frac{2}{3}\sqrt{21}$ (3) $\frac{3}{2}\sqrt{21}$ (4) $2\sqrt{21}$

Ans. (2)

Sol.

 $A \xrightarrow{7}{3} D 3$

BD = hcot30° = h
$$\sqrt{3}$$

So, 7² + 5² = 2(h $\sqrt{3}$)² + 3²)
 \Rightarrow 37 = 3h² + 9.
 \Rightarrow 3h² = 28
 \Rightarrow h = $\sqrt{\frac{28}{3}} = \frac{2}{3}\sqrt{21}$

2. Let $f : R \rightarrow R$ be a function such that $f(x) = x^3 + x^2 f'(1) + x f''(2) + f'''(3), x \in R.$ Then f(2) equal :

Then f(2) equal : (1) 8(2) - 2(3) - 4(4) 30Ans. (2) **Sol.** $f(x) = x^3 + x^2 f'(1) + x f''(2) + f'''(3)$ $\Rightarrow f'(\mathbf{x}) = 3\mathbf{x}^2 + 2\mathbf{x}f'(1) + f''(\mathbf{x})$(1) \Rightarrow f''(x) = 6x + 2f'(1).....(2) $\Rightarrow f'''(\mathbf{x}) = 6$(3) put x = 1 in equation (1) : f'(1) = 3 + 2f'(1) + f''(2).....(4) put x = 2 in equation (2) : f''(2) = 12 + 2f'(1).....(5) from equation (4) & (5) : -3 - f'(1) = 12 + 2f'(1) $\Rightarrow 3f'(1) = -15$ \Rightarrow $f'(1) = -5 \Rightarrow f''(2) = 2 \dots (2)$

put x = 3 in equation (3) : f'''(3) = 6∴ $f(x) = x^3 - 5x^2 + 2x + 6$ f(2) = 8 - 20 + 4 + 6 = -2

3. If a circle C passing through the point (4,0) touches the circle $x^2 + y^2 + 4x - 6y = 12$ externally at the point (1, -1), then the radius of C is :

(1)
$$\sqrt{57}$$
 (2) 4 (3) $2\sqrt{5}$ (4) 5

Ans. (4)

- Sol. $x^2 + y^2 + 4x 6y 12 = 0$ Equation of tangent at (1, -1) x - y + 2(x + 1) - 3(y - 1) - 12 = 0 3x - 4y - 7 = 0∴ Equation of circle is $(x^2 + y^2 + 4x - 6y - 12) + \lambda(3x - 4y - 7) = 0$ It passes through (4, 0) : $(16 + 16 - 12) + \lambda(12 - 7) = 0$ $\Rightarrow 20 + \lambda(5) = 0$ $\Rightarrow \lambda = -4$ ∴ $(x^2 + y^2 + 4x - 6y - 12) - 4(3x - 4y - 7) = 0$ or $x^2 + y^2 - 8x + 10y + 16 = 0$ Radius = $\sqrt{16 + 25 - 16} = 5$
- 4. In a class of 140 students numbered 1 to 140, all even numbered students opted mathematics course, those whose number is divisible by 3 opted Physics course and theose whose number is divisible by 5 opted Chemistry course. Then the number of students who did not opt for any of the three courses is :

(1) 102 (2) 42 (3) 1 (4) 38

Ans. (4)

Sol. Let n(A) = number of students opted Mathematics = 70, n(B) = number of students opted Physics = 46, n(C) = number of students opted Chemistry

$$= 28,$$

$$n(A \cap B) = 23,$$



E

 $n(B \cap C) = 9.$ $n(A \cap C) = 14,$ $n(A \cap B \cap C) = 4,$ Now $n(A \cup B \cup C)$ $= n(A) + n(B) + n(C) - n(A \cap B) - n(B \cap C)$ $-n(A \cap C) + n(A \cap B \cap C)$ = 70 + 46 + 28 - 23 - 9 - 14 + 4 = 102So number of students not opted for any course = Total – $n(A \cup B \cup C)$ = 140 - 102 = 38

5. The sum of all two digit positive numbers which when divided by 7 yield 2 or 5 as remainder is :

> (4) 1356 (1) 1365 (2) 1256 (3) 1465

Ans. (4)

Sol. $\sum_{r=2}^{13} (7r+2) = 7 \cdot \frac{2+13}{2} \times 6 + 2 \times 12$ $= 7 \times 90 + 24 = 654$ $\sum_{i=1}^{13} (7r+5) = 7\left(\frac{1+13}{2}\right) \times 13 + 5 \times 13 = 702$ Total = 654 + 702 = 1356Let $\vec{a} = 2\hat{i} + \lambda_1\hat{j} + 3\hat{k}$, $\vec{b} = 4\hat{i} + (3 - \lambda_2)\hat{j} + 6\hat{k}$ and 6.

 $\vec{c} = 3\hat{i} + 6\hat{j} + (\lambda_3 - 1)\hat{k}$ be three vectors such that $\vec{b} = 2\vec{a}$ and \vec{a} is perpendicular to \vec{c} . Then a possible value of $(\lambda_1, \lambda_2, \lambda_3)$ is :-

 $(1)\left(\frac{1}{2}, 4, -2\right)$ (2) $\left(-\frac{1}{2}, 4, 0\right)$ (3) (1,3,1)(4) (1.5.1)

Ans. (2)

Sol.
$$4\hat{i} + (3 - \lambda_2)\hat{j} + 6\hat{k} = 4\hat{i} + 2\lambda_1\hat{j} + 6\hat{k}$$

 $\Rightarrow 3 - \lambda_2 = 2\lambda_1 \Rightarrow 2\lambda_1 + \lambda_2 = 3 \qquad \dots (1)$
Given $\vec{a}.\vec{c} = 0$
 $\Rightarrow 6 + 6\lambda_1 + 3(\lambda_3 - 1) = 0$
 $\Rightarrow 2\lambda_1 + \lambda_3 = -1 \qquad \dots (2)$
Now $(\lambda_1, \lambda_2, \lambda_3) = (\lambda_1, 3 - 2\lambda_1, -1 - 2\lambda_1)$
Now check the options, option (2) is correct
7. The equation of a tangent to the hyperbola

 $4x^2-5y^2 = 20$ parallel to the line x-y = 2 is : (1) x-y+9 = 0(2) x-y+7 = 0(3) x-y+1 = 0(4) x-y-3 = 0

Ans. (3) **Sol.** Hyperbola $\frac{x^2}{5} - \frac{y^2}{4} = 1$ slope of tangent = 1equation of tangent $y = x \pm \sqrt{5-4}$ \Rightarrow y = x ± 1 \Rightarrow y = x + 1 or y = x - 1 If the area enclosed between the curves $y=kx^2$ 8. and $x=ky^2$, (k>0), is 1 square unit. Then k is: (1) $\frac{1}{\sqrt{3}}$ (2) $\frac{2}{\sqrt{3}}$ (3) $\frac{\sqrt{3}}{2}$ (4) $\sqrt{3}$ Ans. (1) **Sol.** Area bounded by $y^2 = 4ax \& x^2 = 4by$, $a, b \neq 0$ is $\frac{16ab}{3}$ by using formula : $4a = \frac{1}{4b} = 4b, k > 0$ Area = $\frac{16.\frac{1}{4k}.\frac{1}{4k}}{3} = 1$ $\Rightarrow k^2 = \frac{1}{2}$ $\Rightarrow k = \frac{1}{\sqrt{3}}$ Let $f(x) = \begin{cases} \max\{|x|, x^2\}, & |x| \le 2\\ 8-2|x|, & 2 < |x| \le 4 \end{cases}$ 9. Let S be the set of points in the interval (-4,4)at which f is not differentiable. Then S: (1) is an empty set (2) equals $\{-2, -1, 1, 2\}$ (3) equals $\{-2, -1, 0, 1, 2\}$ (4) equals $\{-2, 2\}$ Ans. (3) S

Fol.
$$f(x) = \begin{cases} 8+2x, -4 \le x < -2 \\ x^2, -2 \le x \le -1 \\ |x|, -1 < x < 1 \\ x^2, 1 \le x \le 2 \\ 8-2x, 2 < x \le 4 \end{cases}$$

2



10. If the parabolas y²=4b(x-c) and y²=8ax have a common normal, then which one of the following is a valid choice for the ordered triad (a,b,c)

(1) (1, 1, 0)
(2)
$$\left(\frac{1}{2}, 2, 3\right)$$

(3) $\left(\frac{1}{2}, 2, 0\right)$
(4) (1, 1, 3)

Ans. (1,2,3,4)

Sol. Normal to these two curves are $y = m(x - c) - 2bm - bm^3$, $y = mx - 4am - 2am^3$ If they have a common normal $(c + 2b) m + bm^3 = 4am + 2am^3$ Now $(4a - c - 2b) m = (b - 2a)m^3$ We get all options are correct for m = 0(common normal x-axis) Ans. (1), (2), (3), (4) Remark :

If we consider question as

If the parabolas $y^2 = 4b(x - c)$ and $y^2 = 8ax$ have a common normal other than x-axis, then which one of the following is a valid choice for the ordered triad (a, b, c) ?

When $m \neq 0$: $(4a - c - 2b) = (b - 2a)m^2$

 $m^2 = \frac{c}{2a-b} - 2 > 0 \Longrightarrow \frac{c}{2a-b} > 2$

Now according to options, option 4 is correct

11. The sum of all values of
$$\theta \in \left(0, \frac{\pi}{2}\right)$$
 satisfying
 $\sin^2 2\theta + \cos^4 2\theta = \frac{3}{2}$ is :

(1)
$$\frac{\pi}{2}$$
 (2) π (3) $\frac{3\pi}{8}$ (4) $\frac{5\pi}{4}$

Ans. (1)

Ε

Sol.
$$\sin^2 2\theta + \cos^4 2\theta = \frac{3}{4}, \ \theta \in \left(0, \frac{\pi}{2}\right)$$

$$\Rightarrow 1 - \cos^2 2\theta + \cos^4 2\theta = \frac{3}{4}$$

$$\Rightarrow 4\cos^4 2\theta - 4\cos^2 2\theta + 1 = 0$$

$$\Rightarrow (2\cos^2 2\theta - 1)^2 = 0$$

$$\Rightarrow \cos^2 2\theta = \frac{1}{2} = \cos^2 \frac{\pi}{4}$$

$$\Rightarrow 2\theta = n\pi \pm \frac{\pi}{4}, n \in I$$
$$\Rightarrow \theta = \frac{n\pi}{2} \pm \frac{\pi}{8}$$
$$\Rightarrow \theta = \frac{\pi}{8}, \frac{\pi}{2} - \frac{\pi}{8}$$
Sum of solutions $\frac{\pi}{2}$

12. Let z_1 and z_2 be any two non-zero complex numbers such that $3|z_1| = 4 |z_2|$.

If
$$z = \frac{3z_1}{2z_2} + \frac{2z_2}{3z_1}$$
 then :
(1) $|z| = \frac{1}{2}\sqrt{\frac{17}{2}}$ (2) $\operatorname{Re}(z) = 0$
(3) $|z| = \sqrt{\frac{5}{2}}$ (4) $\operatorname{Im}(z) = 0$

Ans. (Bonus)

Sol.
$$3|z_1| = 4|z_2|$$

 $\Rightarrow \frac{|z_1|}{|z_2|} = \frac{4}{3}$
 $\Rightarrow \frac{|3z_1|}{|2z_2|} = 2$
Let $\frac{3z_1}{2z_2} = a = 2\cos\theta + 2i\sin\theta$
 $z = \frac{3z_1}{2z_2} + \frac{2z_2}{3z_1} = a + \frac{1}{a}$
 $= \frac{5}{2}\cos\theta + \frac{3}{2}i\sin\theta$
Now all options are incorrect

Remark :

There is a misprint in the problem actual problem should be :

"Let z_1 and z_2 be any non-zero complex number such that $3|z_1| = 2|z_2|$.

If
$$z = \frac{3z_1}{2z_2} + \frac{2z_2}{3z_1}$$
, then"

Given

$$3|z_1| = 2|z_2|$$

Now $\left|\frac{3z_1}{2z_2}\right| = 1$

collegebatch.com

Let
$$\frac{3z_1}{2z_2} = a = \cos \theta + i \sin \theta$$

 $z = \frac{3z_1}{2z_2} + \frac{2z_2}{3z_1}$
 $= a + \frac{1}{a} = 2\cos \theta$
 $\therefore \text{ Im}(z) = 0$
Now option (4) is correct.
If the system of equations

x+y+z = 5x+2y+3z = 9 $x+3y+\alpha z = \beta$

13.

has infinitely many solutions, then $\beta\text{-}\alpha$ equals:

(1) 5 (2) 18 (3) 21 (4) 8 Ans. (4)

Sol. $D = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 3 & \alpha \end{vmatrix} = \begin{vmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 0 & 2 & \alpha - 1 \end{vmatrix} = (\alpha - 1) - 4 = (\alpha - 5)$

for infinite solutions $D = 0 \Rightarrow \alpha = 5$

$$D_x = 0 \Longrightarrow \begin{vmatrix} 5 & 1 & 1 \\ 9 & 2 & 3 \\ \beta & 3 & 5 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} 0 & 0 & 1 \\ -1 & -1 & 3 \\ \beta - 15 & -2 & 5 \end{vmatrix} = 0$$

$$\Rightarrow 2 + \beta - 15 = 0 \Rightarrow \beta - 13 = 0$$

on $\beta = 13$ we get $D_y = D_z = 0$
 $\alpha = 5, \beta = 13$

14. The shortest distance between the point $\left(\frac{3}{2}, 0\right)$ and the curve $y = \sqrt{x}, (x > 0)$ is :

(1)
$$\frac{\sqrt{5}}{2}$$
 (2) $\frac{5}{4}$ (3) $\frac{3}{2}$ (4) $\frac{\sqrt{3}}{2}$

Ans. (1)

Sol. Let points
$$\left(\frac{3}{2}, 0\right)$$
, $(t^2, t), t > 0$

Distance $=\sqrt{t^2 + (t^2 - \frac{3}{2})^2}$ $=\sqrt{t^4 - 2t^2 + \frac{9}{4}} = \sqrt{(t^2 - 1)^2 + \frac{5}{4}}$ So minimum distance is $\sqrt{\frac{5}{4}} = \frac{\sqrt{5}}{2}$

15. Consider the quadratic equation $(c-5)x^2-2cx + (c-4) = 0$, $c \neq 5$. Let S be the set of all integral values of c for which one root of the equation lies in the interval (0,2) and its other root lies in the interval (2,3). Then the number of elements in S is :

(1) 11 (2) 18 (3) 10 (4) 12 Ans. (1)

Sol.
$$f(x) = (c - 5)x^2 - 2cx + c - 4$$

$$f(0)f(2) < 0 \qquad \dots (1)$$

$$f(2)f(3) < 0 \qquad \dots (2)$$

from (1) & (2)

$$f(x) = (c - 5)x^2 - 2cx + c - 4$$

$$(c - 4)(c - 24) < 0$$

& $(c - 24)(4c - 49) < 0$

$$\Rightarrow \frac{49}{4} < c < 24$$

:. $s = \{13, 14, 15, ..., 23\}$ Number of elements in set S = 11

16.
$$\sum_{i=1}^{20} \left(\frac{{}^{20}C_{i-1}}{{}^{20}C_i + {}^{20}C_{i-1}} \right) = \frac{k}{21}, \text{ then } k \text{ equals } :$$

(1) 200 (2) 50 (3) 100 (4) 400
Ans. (3)

Sol.
$$\sum_{i=1}^{20} \left(\frac{{}^{20}C_{i-1}}{{}^{20}C_{i} + {}^{20}C_{i-1}} \right)^{3} = \frac{k}{21}$$
$$\Rightarrow \sum_{i=1}^{20} \left(\frac{{}^{20}C_{i-1}}{{}^{21}C_{i}} \right)^{3} = \frac{k}{21}$$
$$\Rightarrow \sum_{i=1}^{20} \left(\frac{i}{21} \right)^{3} = \frac{k}{21}$$
$$\Rightarrow \frac{1}{(21)^{3}} \left[\frac{20(21)}{2} \right]^{2} = \frac{k}{21}$$
$$\Rightarrow 100 = k$$

4

collegebatch.com

17. Let $d \in \mathbb{R}$, and

$$A = \begin{bmatrix} -2 & 4+d & (\sin \theta) - 2 \\ 1 & (\sin \theta) + 2 & d \\ 5 & (2\sin \theta) - d & (-\sin \theta) + 2 + 2d \end{bmatrix},$$

 $\theta \in [0,2\pi]$. If the minimum value of det(A) is 8, then a value of d is :

(1) -7 (2)
$$2(\sqrt{2}+2)$$

(3) -5 (4) $2(\sqrt{2}+1)$

Ans. (3)

Sol. detA =
$$\begin{vmatrix} -2 & 4+d & \sin\theta - 2 \\ 1 & \sin\theta + 2 & d \\ 5 & 2\sin\theta - d & -\sin\theta + 2 + 2d \end{vmatrix}$$
$$(R_1 \rightarrow R_1 + R_3 - 2R_2)$$
$$= \begin{vmatrix} 1 & 0 & 0 \\ 1 & \sin\theta + 2 & d \\ 5 & 2\sin\theta - d & 2 + 2d - \sin\theta \end{vmatrix}$$
$$= (2 + \sin\theta)(2 + 2d - \sin\theta) - d(2\sin\theta - d)$$
$$= 4 + 4d - 2\sin\theta + 2\sin\theta + 2d\sin\theta - \sin^2\theta - 2d\sin\theta + d^2$$
$$= d^2 + 4d + 4 - \sin^2\theta$$
$$= (d + 2)^2 - \sin^2\theta$$
For a given d, minimum value of det(A) = (d + 2)^2 - 1 = 8
$$\Rightarrow d = 1 \text{ or } -5$$

18. If the third term in the binomial expansion of $(1+x^{\log_2 x})^5$ equals 2560, then a possible value of x is :

(1)
$$2\sqrt{2}$$
 (2) $\frac{1}{8}$ (3) $4\sqrt{2}$ (4) $\frac{1}{4}$

Ans. (4)

Sol.
$$(1 + x^{\log_2 x})^5$$

 $T_3 = {}^5C_2 \cdot (x^{\log_2 x})^2 = 2560$
 $\Rightarrow 10 \cdot x^{2\log_2 x} = 2560$
 $\Rightarrow 2(\log_2 x)^2 = \log_2 256$
 $\Rightarrow 2(\log_2 x)^2 = 8$
 $\Rightarrow (\log_2 x)^2 = 4 \Rightarrow \log_2 x = 2 \text{ or } -2$
 $x = 4 \text{ or } \frac{1}{4}$

19. If the line 3x + 4y - 24 = 0 intersects the x-axis at the point A and the y-axis at the point B, then the incentre of the triangle OAB, where O is the origin, is

(1) (3, 4) (2) (2, 2) (3) (4, 4) (4) (4, 3)

Ans. (2)



20. The mean of five observations is 5 and their variance is 9.20. If three of the given five observations are 1, 3 and 8, then a ratio of other two observations is :

Ans. (1)

Sol. Let two observations are $x_1 \& x_2$

mean =
$$\frac{\sum x_i}{5} = 5 \implies 1 + 3 + 8 + x_1 + x_2 = 25$$

 $\implies x_1 + x_2 = 13$ (1)
variance $(\sigma^2) = \frac{\sum x_i^2}{5} - 25 = 9.20$
 $\implies \sum x_i^2 = 171$
 $\implies x_1^2 + x_2^2 = 97$ (2)
by (1) & (2)
 $(x_1 + x_2)^2 - 2x_1x_2 = 97$
or $x_1x_2 = 36$
 $\therefore x_1 : x_2 = 4 : 9$

Ε



(1) parallel to x-axis (2) with slope $\frac{2}{3}$ (3) with slope $\frac{3}{2}$ (4) parallel to y-axis Ans. (2) Sol. Let the centroid of $\triangle PQR$ is (h, k) & P is (α , β), then $\frac{\alpha+1+3}{3} = h$ and $\frac{\beta+4-2}{3} = k$ $\alpha = (3h - 4)$ $\beta = (3k - 4)$ Point P(α , β) lies on line $2x - 3y + 4 = 0$ $\therefore 2(3h - 4) - 3(3k - 2) + 4 = 0$ \Rightarrow locus is $6x - 9y + 2 = 0$ 22. If $\frac{dy}{dx} + \frac{3}{\cos^2 x}y = \frac{1}{\cos^2 x}, x \in \left(-\frac{\pi}{3}, \frac{\pi}{3}\right)$, and $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$, then $y\left(-\frac{\pi}{4}\right)$ equals : (1) $\frac{1}{3} + e^{6}$ (2) $\frac{1}{3}$ (3) $-\frac{4}{3}$ (4) $\frac{1}{3} + e^{3}$ Ans. (1) Sol. $\frac{dy}{dx} + 3\sec^2 x.y = \sec^2 x$ I.F. $= e^{3\int \sec^2 x.dx} = e^{3\tan x}$ or $y.e^{3\tan x} = \int \sec^2 x.e^{3\tan x}dx$ or $y.e^{3\tan x} = \int \sec^2 x.e^{3\tan x}dx$ or $y.e^{3\tan x} = \frac{1}{3}e^{3} + C$ (1) Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3}.e^{3} = \frac{1}{3}e^{3} + C$ $\therefore C = e^{3}$	21.	A point P moves on the $IfQ(1,4)$ and $R(3,-2)$ are locus of the centroid of	e line $2x - 3y + 4 = 0$. e fixed points, then the f $\triangle PQR$ is a line :
(3) with slope $\frac{3}{2}$ (4) parallel to y-axis Ans. (2) Sol. Let the centroid of $\triangle PQR$ is (h, k) & P is (α, β) , then $\frac{\alpha+1+3}{3} = h$ and $\frac{\beta+4-2}{3} = k$ $\alpha = (3h-4)$ $\beta = (3k-4)$ Point P(α, β) lies on line $2x - 3y + 4 = 0$ $\therefore 2(3h-4) - 3(3k-2) + 4 = 0$ $\Rightarrow locus is 6x - 9y + 2 = 022. If \frac{dy}{dx} + \frac{3}{\cos^2 x}y = \frac{1}{\cos^2 x}, x \in (\frac{-\pi}{3}, \frac{\pi}{3}), andy(\frac{\pi}{4}) = \frac{4}{3}, then y(-\frac{\pi}{4}) equals :(1) \frac{1}{3} + e^6 (2) \frac{1}{3}(3) -\frac{4}{3} (4) \frac{1}{3} + e^3Ans. (1)Sol. \frac{dy}{dx} + 3\sec^2 x.y = \sec^2 xI.F. = e^{3\int \sec^2 x dx} = e^{3\tan x}or y.e^{3\tan x} = \int \sec^2 x.e^{3\tan x} dxor y.e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C(1)Giveny(\frac{\pi}{4}) = \frac{4}{3}\therefore \frac{4}{3}e^3 = \frac{1}{3}e^3 + C\therefore C = e^3$		(1) parallel to x-axis	(2) with slope $\frac{2}{3}$
Ans. (2) Sol. Let the centroid of $\triangle PQR$ is (h, k) & P is (α, β) , then $\frac{\alpha+1+3}{3} = h$ and $\frac{\beta+4-2}{3} = k$ $\alpha = (3h - 4)$ $\beta = (3k - 4)$ Point P(α, β) lies on line $2x - 3y + 4 = 0$ $\Rightarrow 10cus is 6x - 9y + 2 = 0$ 22. If $\frac{dy}{dx} + \frac{3}{cos^2 x}y = \frac{1}{cos^2 x}, x \in (\frac{-\pi}{3}, \frac{\pi}{3})$, and $y(\frac{\pi}{4}) = \frac{4}{3}$, then $y(-\frac{\pi}{4})$ equals : (1) $\frac{1}{3} + e^6$ (2) $\frac{1}{3}$ (3) $-\frac{4}{3}$ (4) $\frac{1}{3} + e^3$ Ans. (1) Sol. $\frac{dy}{dx} + 3sec^2 x.y = sec^2 x$ I.F. $= e^{3\int sec^2 xdx} = e^{3inx}$ or $y.e^{3inx} = \int sec^2 x.e^{3inx} dx$ or $y.e^{3inx} = \frac{1}{3}e^{3inx} + C$ (1) Given $y(\frac{\pi}{4}) = \frac{4}{3}$ $\therefore \frac{4}{3}e^3 = \frac{1}{3}e^3 + C$ $\therefore C = e^3$		(3) with slope $\frac{3}{2}$	(4) parallel to y-axis
$\frac{\alpha + 1 + 3}{3} = h \text{and} \frac{\beta + 4 - 2}{3} = k$ $\alpha = (3h - 4) \qquad \beta = (3k - 4)$ Point P(α , β) lies on line $2x - 3y + 4 = 0$ $\therefore 2(3h - 4) - 3(3k - 2) + 4 = 0$ $\Rightarrow \text{ locus is } 6x - 9y + 2 = 0$ 22. If $\frac{dy}{dx} + \frac{3}{\cos^2 x} y = \frac{1}{\cos^2 x}, x \in \left(\frac{-\pi}{3}, \frac{\pi}{3}\right)$, and $y\left(\frac{\pi}{4}\right) = \frac{4}{3}, \text{ then } y\left(-\frac{\pi}{4}\right) \text{ equals }:$ (1) $\frac{1}{3} + e^6$ (2) $\frac{1}{3}$ (3) $-\frac{4}{3}$ (4) $\frac{1}{3} + e^3$ Ans. (1) Sol. $\frac{dy}{dx} + 3\sec^2 x.y = \sec^2 x$ I.F. $= e^{3\int \sec^2 x.dx} = e^{3\tan x}$ or $y.e^{3\tan x} = \int \sec^2 x.e^{3\tan x} dx$ or $y.e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C$ (1) Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3}.e^3 = \frac{1}{3}e^3 + C$ $\therefore C = e^3$	Ans. Sol.	(2) Let the centroid of Δ (α , β), then	PQR is (h, k) & P is
$\alpha = (3h - 4) \qquad \beta = (3k - 4)$ Point P(α , β) lies on line 2x - 3y + 4 = 0 $\therefore 2(3h - 4) - 3(3k - 2) + 4 = 0$ \Rightarrow locus is $6x - 9y + 2 = 0$ 22. If $\frac{dy}{dx} + \frac{3}{\cos^2 x} y = \frac{1}{\cos^2 x}, x \in \left(\frac{-\pi}{3}, \frac{\pi}{3}\right)$, and $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$, then $y\left(-\frac{\pi}{4}\right)$ equals : (1) $\frac{1}{3} + e^6$ (2) $\frac{1}{3}$ (3) $-\frac{4}{3}$ (4) $\frac{1}{3} + e^3$ Ans. (1) Sol. $\frac{dy}{dx} + 3\sec^2 x.y = \sec^2 x$ I.F. $= e^{3\int \sec^2 x dx} = e^{3\tan x}$ or $y.e^{3\tan x} = \int \sec^2 x.e^{3\tan x} dx$ or $y.e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C$ (1) Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3}.e^3 = \frac{1}{3}e^3 + C$ $\therefore C = e^3$		$\frac{\alpha+1+3}{3} = h \text{and} $	$\frac{\beta+4-2}{3} = k$
22. If $\frac{dy}{dx} + \frac{3}{\cos^2 x} y = \frac{1}{\cos^2 x}, x \in \left(\frac{-\pi}{3}, \frac{\pi}{3}\right)$, and $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$, then $y\left(-\frac{\pi}{4}\right)$ equals : (1) $\frac{1}{3} + e^6$ (2) $\frac{1}{3}$ (3) $-\frac{4}{3}$ (4) $\frac{1}{3} + e^3$ Ans. (1) Sol. $\frac{dy}{dx} + 3\sec^2 x.y = \sec^2 x$ I.F. $= e^{3\int \sec^2 x.dx} = e^{3\tan x}$ or $y.e^{3\tan x} = \int \sec^2 x.e^{3\tan x} dx$ or $y.e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C$ (1) Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3}.e^3 = \frac{1}{3}e^3 + C$ $\therefore C = e^3$		$\alpha = (3h - 4)$ Point P(α , β) lies on li $\therefore 2(3h - 4) - 3(3k - 3)$ \Rightarrow locus is $6x - 9y + 3$	$\beta = (3k - 4)$ ne 2x - 3y + 4 = 0 2) + 4 = 0 2 = 0
$y\left(\frac{\pi}{4}\right) = \frac{4}{3}, \text{ then } y\left(-\frac{\pi}{4}\right) \text{ equals :}$ (1) $\frac{1}{3} + e^{6}$ (2) $\frac{1}{3}$ (3) $-\frac{4}{3}$ (4) $\frac{1}{3} + e^{3}$ Ans. (1) Sol. $\frac{dy}{dx} + 3\sec^{2} x \cdot y = \sec^{2} x$ I.F. $= e^{3\int \sec^{2} x \cdot y} = \sec^{2} x$ or $y \cdot e^{3\tan x} = \int \sec^{2} x \cdot e^{3\tan x} dx$ or $y \cdot e^{3\tan x} = \int \sec^{2} x \cdot e^{3\tan x} dx$ (1) Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3} \cdot e^{3} = \frac{1}{3} e^{3} + C$ $\therefore C = e^{3}$	22.	If $\frac{dy}{dx} + \frac{3}{\cos^2 x}y = \frac{1}{\cos^2 x}$	$\overline{x}, x \in \left(\frac{-\pi}{3}, \frac{\pi}{3}\right)$, and
(1) $\frac{1}{3} + e^{6}$ (2) $\frac{1}{3}$ (3) $-\frac{4}{3}$ (4) $\frac{1}{3} + e^{3}$ Ans. (1) Sol. $\frac{dy}{dx} + 3\sec^{2} x.y = \sec^{2} x$ I.F. $= e^{3\int \sec^{2} x dx} = e^{3\tan x}$ or $y.e^{3\tan x} = \int \sec^{2} x.e^{3\tan x} dx$ or $y.e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C$ (1) Given $y(\frac{\pi}{4}) = \frac{4}{3}$ $\therefore \frac{4}{3}.e^{3} = \frac{1}{3}e^{3} + C$ $\therefore C = e^{3}$		$y\left(\frac{\pi}{4}\right) = \frac{4}{3}$, then $y\left(-\frac{\pi}{4}\right)$	equals :
(3) $-\frac{4}{3}$ (4) $\frac{1}{3} + e^{3}$ Ans. (1) Sol. $\frac{dy}{dx} + 3\sec^{2} x \cdot y = \sec^{2} x$ I.F. $= e^{3\int \sec^{2} x \cdot x} = e^{3\tan x}$ or $y \cdot e^{3\tan x} = \int \sec^{2} x \cdot e^{3\tan x} dx$ or $y \cdot e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C$ (1) Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3}e^{3} = \frac{1}{3}e^{3} + C$ $\therefore C = e^{3}$		(1) $\frac{1}{3} + e^6$	(2) $\frac{1}{3}$
Ans. (1) Sol. $\frac{dy}{dx} + 3\sec^2 x \cdot y = \sec^2 x$ I.F. $= e^{3\int \sec^2 x dx} = e^{3\tan x}$ or $y \cdot e^{3\tan x} = \int \sec^2 x \cdot e^{3\tan x} dx$ or $y \cdot e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C$ (1) Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3} \cdot e^3 = \frac{1}{3}e^3 + C$ $\therefore C = e^3$		$(3) -\frac{4}{3}$	(4) $\frac{1}{3} + e^3$
Sol. $\frac{dy}{dx} + 3\sec^2 x \cdot y = \sec^2 x$ I.F. $= e^{3\int \sec^2 x dx} = e^{3\tan x}$ or $y \cdot e^{3\tan x} = \int \sec^2 x \cdot e^{3\tan x} dx$ or $y \cdot e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C$ (1) Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3} \cdot e^3 = \frac{1}{3}e^3 + C$ $\therefore C = e^3$	Ans.	(1)	
I.F. = $e^{3\int \sec^2 x dx} = e^{3\tan x}$ or $y \cdot e^{3\tan x} = \int \sec^2 x \cdot e^{3\tan x} dx$ or $y \cdot e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C$ (1) Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3} \cdot e^3 = \frac{1}{3}e^3 + C$ $\therefore C = e^3$	Sol.	$\frac{\mathrm{d}y}{\mathrm{d}x} + 3\sec^2 x \cdot y = \sec^2 x$	
or $y \cdot e^{3\tan x} = \int \sec^2 x \cdot e^{3\tan x} dx$ or $y \cdot e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C$ (1) Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3} \cdot e^3 = \frac{1}{3}e^3 + C$ $\therefore C = e^3$		I.F. = $e^{3\int \sec^2 x dx} = e^{3\tan x}$	
or $y \cdot e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C$ (1) Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3} \cdot e^3 = \frac{1}{3}e^3 + C$ $\therefore C = e^3$		or $y.e^{3\tan x} = \int \sec^2 x.e^{3\tan x}$	^x dx
Given $y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3} \cdot e^{3} = \frac{1}{3}e^{3} + C$ $\therefore C = e^{3}$ 6		or $y \cdot e^{3\tan x} = \frac{1}{3}e^{3\tan x} + C$	(1)
$y\left(\frac{\pi}{4}\right) = \frac{4}{3}$ $\therefore \frac{4}{3} \cdot e^{3} = \frac{1}{3}e^{3} + C$ $\therefore C = e^{3}$		Given	
$\therefore \frac{4}{3} \cdot e^3 = \frac{1}{3} e^3 + C$ $\therefore C = e^3$		$y\left(\frac{\pi}{4}\right) = \frac{4}{3}$	
6		$\therefore \frac{4}{3} \cdot e^3 = \frac{1}{3} e^3 + C$ $\therefore C = e^3$	
	6		

Now put
$$x = -\frac{\pi}{4}$$
 in equation (1)
 $\therefore y \cdot e^{-3} = \frac{1}{3}e^{-3} + e^{3}$
 $\therefore y = \frac{1}{3} + e^{6}$
 $\therefore y\left(-\frac{\pi}{4}\right) = \frac{1}{3} + e^{6}$
The plane passing through the point (4, -1, 2)
and parallel to the lines $\frac{x+2}{3} = \frac{y-2}{-1} = \frac{z+1}{2}$
and $\frac{x-2}{1} = \frac{y-3}{2} = \frac{z-4}{3}$ also passes through
the point :

z+12

(1) (-1, -1, -1)(2) (-1, -1, 1)(3)(1, 1, -1)(4)(1, 1, 1)

23.

Sol. Let \vec{n} be the normal vector to the plane passing through (4, -1, 2) and parallel to the lines $L_1 \& L_2$

then
$$\vec{n} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -1 & 2 \\ 1 & 2 & 3 \end{vmatrix}$$

 $\therefore \vec{n} = -7\hat{i} - 7\hat{j} + 7\hat{k}$
 \therefore Equation of plane

:. Equation of plane is -1(x - 4) - 1(y + 1) + 1(z - 2) = 0 $\therefore x + y - z - 1 = 0$ Now check options

Let $I = \int_{a}^{b} (x^4 - 2x^2) dx$. If I is minimum then 24. the ordered pair (a, b) is :

(1)
$$(-\sqrt{2},0)$$
 (2) $(-\sqrt{2},\sqrt{2})$
(3) $(0,\sqrt{2})$ (4) $(\sqrt{2},-\sqrt{2})$

Ans. (2)

Sol. Let
$$f(x) = x^2(x^2 - 2)$$

$$-\sqrt{2}$$

As long as f(x) lie below the x-axis, definite integral will remain negative,

so correct value of (a, b) is $(-\sqrt{2},\sqrt{2})$ for minimum of I



25. If 5, 5r, 5r² are the lengths of the sides of a triangle, then r cannot be equal to :

(1)
$$\frac{3}{2}$$
 (2) $\frac{3}{4}$ (3) $\frac{5}{4}$ (4) $\frac{7}{4}$

Ans. (4)

Sol.
$$r = 1$$
 is obviously true.
Let $0 < r < 1$
 $\Rightarrow r + r^2 > 1$
 $\Rightarrow r^2 + r - 1 > 0$
 $\left(r - \frac{-1 - \sqrt{5}}{2}\right) \left(r - \left(\frac{-1 + \sqrt{5}}{2}\right)\right)$
 $\Rightarrow r - \frac{-1 - \sqrt{5}}{2}$ or $r > \frac{-1 + \sqrt{5}}{2}$
 $r \in \left(\frac{\sqrt{5} - 1}{2}, 1\right)$
 $\frac{\sqrt{5} - 1}{2} < r < 1$

When r > 1

$$\Rightarrow \frac{\sqrt{5}+1}{2} > \frac{1}{r} > 1$$
$$\Rightarrow r \in \left(\frac{\sqrt{5}-1}{2}, \frac{\sqrt{5}+1}{2}\right)$$

Now check options

- 26. Consider the statement : "P(n): $n^2 n + 41$ is prime." Then which one of the following is true?
 - (1) P(5) is false but P(3) is true
 - (2) Both P(3) and P(5) are false
 - (3) P(3) is false but P(5) is true
 - (4) Both P(3) and P(5) are true

Ans. (4)

Ε

Sol. $P(n) : n^2 - n + 41$ is prime P(5) = 61 which is prime P(3) = 47 which is also prime 27. Let A be a point on the line $\vec{r} = (1-3\mu)\hat{i} + (\mu-1)\hat{j} + (2+5\mu)\hat{k}$ and B(3, 2, 6) be a point in the space. Then the value of μ for which the vector \overrightarrow{AB} is parallel to the plane x - 4y + 3z = 1 is :

(1)
$$\frac{1}{2}$$
 (2) $-\frac{1}{4}$ (3) $\frac{1}{4}$ (4) $\frac{1}{8}$

Ans. (3)

Sol. Let point A is

$$(1-3\mu)\hat{i} + (\mu-1)\hat{j} + (2+5\mu)\hat{k}$$

and point B is (3, 2, 6)
then $\overline{AB} = (2+3\mu)\hat{i} + (3-\mu)\hat{j} + (4-5\mu)\hat{k}$
which is parallel to the plane $x - 4y + 3z = 1$
 $\therefore 2 + 3\mu - 12 + 4\mu + 12 - 15\mu = 0$
 $8\mu = 2$
 $\mu = \frac{1}{4}$

28. For each $t \in \mathbb{R}$, let [t] be the greatest integer less than or equal to t. Then,

$$\lim_{x \to 1+} \frac{(1-|x|+\sin|1-x|)\sin\left(\frac{\pi}{2}[1-x]\right)}{|1-x|[1-x]}$$
(1) equals -1
(2) equals 1
(3) does not exist
(4) equals 0

Ans. (4)

Sol.
$$\lim_{x \to 1^{+}} \frac{(1 - |x| + \sin|1 - x|) \sin\left(\frac{\pi}{2}[1 - x]\right)}{|1 - x|[1 - x]}$$
$$= \lim_{x \to 1^{+}} \frac{(1 - x) + \sin(x - 1)}{(x - 1)(-1)} \sin\left(\frac{\pi}{2}(-1)\right)$$
$$= \lim_{x \to 1^{+}} \left(1 - \frac{\sin(x - 1)}{(x - 1)}\right)(-1) = (1 - 1)(-1) = 0$$



29. An unbiased coin is tossed. If the outcome is a head then a pair of unbiased dice is rolled and the sum of the numbers obtained on them is noted. If the toss of the coin results in tail then a card from a well-shuffled pack of nine cards numbered 1,2,3,...,9 is randomly picked and the number on the card is noted. The probability that the noted number is either 7 or 8 is :

(1) $\frac{13}{36}$ (2) $\frac{19}{36}$ (3) $\frac{19}{72}$ (4) $\frac{15}{72}$

Ans. (3)

Sol. Start
$$H \rightarrow \text{Sum 7 or } 8 \Rightarrow \frac{11}{36}$$

 $1/2$ $T \rightarrow \text{Number is 7 or } 8 = \frac{2}{9}$

 $P(A) = \frac{1}{2} \times \frac{11}{36} + \frac{1}{2} \times \frac{2}{9} = \frac{19}{72}$

30. Let $n \ge 2$ be a natural number and $0 < \theta < \pi/2$.

Then $\int \frac{(\sin^n \theta - \sin \theta)^{\frac{1}{n}} \cos \theta}{\sin^{n+1} \theta} d\theta$ is equal to :

(Where C is a constant of integration)

$$(1) \frac{n}{n^{2}-1} \left(1 - \frac{1}{\sin^{n+1}\theta}\right)^{\frac{n+1}{n}} + C$$

$$(2) \frac{n}{n^{2}+1} \left(1 - \frac{1}{\sin^{n-1}\theta}\right)^{\frac{n+1}{n}} + C$$

$$(3) \frac{n}{n^{2}-1} \left(1 - \frac{1}{\sin^{n-1}\theta}\right)^{\frac{n+1}{n}} + C$$

$$(4) \frac{n}{n^{2}-1} \left(1 + \frac{1}{\sin^{n-1}\theta}\right)^{\frac{n+1}{n}} + C$$

Ans. (3)

Sol.
$$\int \frac{(\sin^n \theta - \sin \theta)^{1/n} \cos \theta}{\sin^{n+1} \theta} d\theta$$
$$= \int \frac{\sin \theta \left(1 - \frac{1}{\sin^{n-1} \theta}\right)^{1/n}}{\sin^{n+1} \theta} d\theta$$
Put $1 - \frac{1}{\sin^{n-1} \theta} = t$ So $\frac{(n-1)}{\sin^n \theta} \cos \theta d\theta = dt$ Now $\frac{1}{n-1} \int (t)^{1/n} dt$
$$= \frac{1}{(n-1)} \frac{(t)^{\frac{1}{n+1}}}{\frac{1}{n+1}} + C$$
$$= \frac{1}{(n-1)} \left(1 - \frac{1}{\sin^{n-1} \theta}\right)^{\frac{1}{n+1}} + C$$

8

collegebatch.com

TEST PAPER OF JEE(MAIN) EXAMINATION - 2019 (Held On Thursday 10th JANUARY, 2019) TIME : 09 : 30 AM To 12 : 30 PM PHYSICS

1. A uniform metallic wire has a resistance of 18 Ω and is bent into an equilateral triangle. Then, the resistance between any two vertices of the triangle is :

(1) 8 Ω (3) 4 Ω (2) 12 Ω (4) 2Ω Ans. (3)



R_{eq} between any two vertex will be

$$\frac{1}{R_{eq}} = \frac{1}{12} + \frac{1}{6} \Longrightarrow R_{eq.} = 4\Omega$$

2. A satellite is moving with a constant speed v in circular orbit around the earth. An object of mass 'm' is ejected from the satellite such that it just escapes from the gravitational pull of the earth. At the time of ejection, the kinetic energy of the object is :

(1)
$$\frac{3}{2}$$
 mv²
(2) mv²
(3) 2mv²
(4) $\frac{1}{2}$ mv²

Ans. (2)

Sol. At height r from center of earth. orbital velocity

$$=\sqrt{\frac{\mathrm{GM}}{\mathrm{r}}}$$

... By energy conservation

KE of 'm' +
$$\left(-\frac{GMm}{r}\right) = 0 + 0$$

(At infinity, PE = KE = 0)

$$\Rightarrow$$
 KE of 'm' = $\frac{GMm}{r} = \left(\sqrt{\frac{GM}{r}}\right)^2 m = mv^2$

3. A solid metal cube of edge length 2 cm is moving in a positive y direction at a constant speed of 6 m/s. There is a uniform magnetic field of 0.1 T in the positive z-direction. The potential difference between the two faces of the cube perpendicular to the x-axis, is : (1) 6 mV (2) 1 mV (3) 12 mV (4) 2 mV

Ans. (3)

Sol. Potential difference between two faces perpendicular to x-axis will be

$$\ell . (\vec{V} \times \vec{B}) = 12mV$$

4. A parallel plate capacitor is of area 6 cm² and a separation 3 mm. The gap is filled with three dielectric materials of equal thickness (see figure) with dielectric constants K_1 , = 10, $K_2 = 12$ and $K_3 = 14$. The dielectric constant of a material which when fully inserted in above capacitor, gives same capacitance would be :

$$\mathbf{K}_{1}$$
 \mathbf{K}_{2} \mathbf{K}_{3} mm

(3) 36

(4) 14

Ans. (1)

(1) 12

Sol. Let dielectric constant of material used be K.

(2) 4

$$\therefore \frac{10 \in A/3}{d} + \frac{12 \in A/3}{d} + \frac{14 \in A/3}{d} = \frac{K \in A/3}{d}$$
$$\Rightarrow K = 12$$

5. A 2 W carbon resistor is color coded with green, black, red and brown respectively. The maximum current which can be passed through this resistor is :

Sol. $P = i^2 R$.

 \therefore for i_{max} , R must be minimum from color coding R = $50 \times 10^2 \Omega$

$$\therefore$$
 $i_{max} = 20mA$



collegebatch.com

6. In a Young's double slit experiment with slit separation 0.1 mm, one observes a bright fringe at angle 1/40 rad by using light of wavelength λ₁. When the light of wavelength λ₂ is used a bright fringe is seen at the same angle in the same set up. Given that λ₁ and λ₂ are in visible range (380 nm to 740 nm), their values are :

(1) 380 nm, 500 nm
(2) 625 nm, 500 nm
(3) 380 nm, 525 nm

Ans. (2)

Sol. Path difference = $d \sin \theta \approx d\theta$

 $= 0.1 \times \frac{1}{40}$ mm = 2500 nm

or bright fringe, path difference must be integral multiple of λ .

- $\therefore 2500 = n\lambda_1 = m\lambda_2$
- :. $\lambda_1 = 625, \lambda_2 = 500$ (from m=5) (for n = 4)
- 7. A magnet of total magnetic moment $10^{-2}\hat{i} A \cdot m^2$ is placed in a time varying magnetic field, $B\hat{i}$ (cost ω t) where B = 1 Tesla and $\omega = 0.125$ rad/s. The work done for reversing the direction of the magnetic moment at t = 1second, is :

(1) 0.007 J (2) 0.014 J (3) 0.01 J (4) 0.028 J Ans. (2)

- Sol. Work done, $W = (\Delta \vec{\mu}) \cdot \vec{B}$ = 2 × 10⁻² × 1 cos(0.125) = 0.02 J
 - \therefore correct answer is (2)
- 8. To mop-clean a floor, a cleaning machine presses a circular mop of radius R vertically down with a total force F and rotates it with a constant angular speed about its axis. If the force F is distributed uniformly over the mop and if coefficient of friction between the mop and the floor is μ , the torque, applied by the machine on the mop is :
 - (1) $\frac{2}{3}\mu FR$ (2) $\mu FR/3$ (3) $\mu FR/2$ (4) $\mu FR/6$ (1)

Ans. (1)

2



Consider a strip of radius x & thickness dx, Torque due to friction on this strip.

$$\int d\tau = \int_{0}^{R} \frac{x\mu F \cdot 2\pi x dx}{\pi R^{2}}$$
$$\tau = \frac{2\mu F}{R^{2}} \cdot \frac{R^{3}}{3}$$
$$\tau = \frac{2\mu FR}{3}$$

 \therefore correct answer is (1)

9. Using a nuclear counter the count rate of emitted particles from a radioactive source is measured. At t = 0 it was 1600 counts per second and t = 8 seconds it was 100 counts per second. The count rate observed, as counts per second, at t = 6 seconds is close to :

Ans. (3)

Sol. at
$$t = 0$$
, $A_0 = \frac{dN}{dt} = 1600C/s$
at $t = 8s$, $A = 100 C/s$
 $\frac{A}{A_0} = \frac{1}{16}$ in 8 sec
Therefor half life is $t_{1/2} = 2$ sec
 \therefore Activity at $t = 6$ will be $1600 \left(\frac{1}{2}\right)^3$
 $= 200 C/s$
 \therefore correct answer is (3)

collegebatch.com

10. If the magnetic field of a plane electromagnetic wave is given by (The speed of light = 3×10^8 /m/s)

B=100 × 10⁻⁶ sin
$$\left[2\pi \times 2 \times 10^{15} \left(t - \frac{x}{c}\right)\right]$$
 then the

maximum electric field associated with it is :

- (1) 4×10^4 N/C (2) 4.5×10^4 N/C
- (3) 6×10^4 N/C (4) 3×10^4 N/C
- Ans. (4)
- Sol. $E_0 = B_0 \times C$ = 100 × 10⁻⁶ × 3 × 10⁸ = 3 × 10⁴ N/C \therefore correct answer is 3 × 10⁴ N/C
- 11. A charge Q is distributed over three concentric spherical shells of radii a, b, c (a < b < c) such that their surface charge densities are equal to one another. The total potential at a point at distance r from their common centre, where r < a, would be :

(1)
$$\frac{Q}{4\pi\epsilon_0(a+b+c)}$$

(2) $\frac{Q(a+b+c)}{4\pi\epsilon_0(a^2+b^2+c^2)}$
(3) $\frac{Q}{12\pi\epsilon_0}\frac{ab+bc+ca}{abc}$
(4) $\frac{Q}{4\pi\epsilon_0}\frac{(a^2+b^2+c^2)}{(a^3+b^3+c^3)}$

Ans. (2)

Ε



Potential at point P,
$$V = \frac{kQ_a}{a} + \frac{kQ_b}{b} + \frac{kQ_c}{c}$$

 $\therefore Q_a : Q_b : Q_c :: a^2 : b^2 : c^2$
 $\{\text{since } \sigma_a = \sigma_b = \sigma_c\}$
 $\therefore Q_a = \left[\frac{a^2}{a^2 + b^2 + c^2}\right]Q$
 $Q_b = \left[\frac{b^2}{a^2 + b^2 + c^2}\right]Q$
 $Q_c = \left[\frac{c^2}{a^2 + b^2 + c^2}\right]Q$
 $V = \frac{Q}{4\pi \epsilon_0} \left[\frac{(a + b + c)}{a^2 + b^2 + c^2}\right]$

- \therefore correct answer is (2) Water flows into a large tank with flat bottom
- 12. Water flows into a large tank with flat bottom at the rate of 10^{-4} m³s⁻¹. Water is also leaking out of a hole of area 1 cm² at its bottom. If the height of the water in the tank remains steady, then this height is:

(1) 4 cm (2) 2.9 cm (3) 1.7 cm (4) 5.1 cm **Ans. (4)**



Since height of water column is constant therefore, water inflow rate (Q_{in})

= water outflow rate

$$Q_{in} = 10^{-4} \text{ m}^3 \text{s}^{-1}$$

 $Q_{out} = Au = 10^{-4} \times \sqrt{2 \text{ gh}}$
 $10^{-4} = 10^{-4} \sqrt{20 \times \text{h}}$
 $h = \frac{1}{20} \text{ m}$
 $h = 5 \text{ cm}$
∴ correct answer is (4)



13. A piece of wood of mass 0.03 kg is dropped from the top of a 100 m height building. At the same time, a bullet of mass 0.02 kg is fired vertically upward, with a velocity 100 ms⁻¹, from the ground. The bullet gets embedded in the wood. Then the maximum height to which the combined system reaches above the top of the building before falling below is : (g =10ms⁻²) (1) 30 m (2) 10 m (3) 40 m (4) 20 m

Ans. (3)

Sol. 100 m € 0.03kg 100 m 100m/s 0.02kg

Time taken for the particles to collide,

$$t = \frac{d}{V_{rel}} = \frac{100}{100} = 1 \sec t$$

Speed of wood just before collision = gt = 10 m/s & speed of bullet just before collision v-gt = 100 - 10 = 90 m/s Now, conservation of linear momentum just before and after the collision --(0.02) (1v) + (0.02) (9v) = (0.05)v $\Rightarrow 150 = 5v$ $\Rightarrow v = 30$ m/s

Max. height reached by body $h = \frac{v^2}{2g}$

$$h = \frac{30 \times 30}{2 \times 10} = 45m$$

$$\therefore$$
 Height above tower = 40 m

14. The density of a material in SI units is 128 kg m⁻³. In certain units in which the unit of length is 25 cm and the unit of mass is 50 g, the numerical value of density of the material is :

(1) 410 (2) 640 (3) 16 (4) 40 Ans. (4)

Sol.
$$\frac{128 \text{kg}}{\text{m}^3} = \frac{125(50 \text{g})(20)}{(25 \text{cm})^3 (4)^3}$$

= $\frac{128}{64} (20) \text{ units}$
= 40 units

15. To get output '1' at R, for the given logic gate circuit the input values must be :



Sol. $p = \overline{x} + v$

$$Q = \overline{\overline{y}.x} = y + \overline{x}$$

$$Q = \overline{\overline{y}.x} = y + \overline{x}$$

$$O/P = \overline{P+Q}$$
To make O/P
$$P + Q$$
 must be 'O
SO, $y = 0$

$$x = 1$$

16. A block of mass m is kept on a platform which starts from rest with constant acceleration g/2 upward, as shown in fig. Work done by normal reaction on block in time t is :

$$m \qquad a = \frac{g}{2}$$
(1) 0
(2) $\frac{3mg^2t^2}{8}$
(3) $-\frac{mg^2t^2}{8}$
(4) $\frac{mg^2t^2}{8}$
Ans. (2)
Sol. $N - mg = \frac{mg}{2} \Rightarrow N = \frac{3mg}{2}$
Now, work done $W = \vec{N}\vec{S} = \left(\frac{3mg}{2}\right)\left(\frac{1}{2}gt^2\right)$
 $\Rightarrow W = \frac{3mg^2t^2}{4}$

4



- 17. A heat source at $T = 10^3$ K is connected to another heat reservoir at $T=10^2$ K by a copper slab which is 1 m thick. Given that the thermal conductivity of copper is 0.1 WK⁻¹ m⁻¹, the energy flux through it in the steady state is :
 - (1) 90 Wm⁻² (2) 200 Wm⁻²
 - (3) 65 Wm⁻² (4) 120 Wm⁻²

Ans. (1)



$$\Rightarrow \frac{1}{A} \left(\frac{dQ}{dt} \right) = \frac{(0.1)(900)}{1} = 90 W / m^2$$

18. A TV transmission tower has a height of 140 m and the height of the receiving antenna is 40 m. What is the maximum distance upto which signals can be broadcasted from this tower in LOS(Line of Sight) mode ? (Given : radius of earth $= 6.4 \times 10^6$ m).

(1)	80 km	(2)	48	km
(3)	40 km	(4)	65	km

Ans. (4)

E

Sol. Maximum distance upto which signal can be broadcasted is

$$d_{max} = \sqrt{2Rh_{T}} + \sqrt{2Rh_{R}}$$

where h_T and h_R are heights of transmiter tower and height of receiver respectively. Putting all values -

$$d_{max} = \sqrt{2 \times 6.4 \times 106} \left[\sqrt{104} + \sqrt{40} \right]$$

on solving, $d_{max} = 65 \text{ km}$

19. A potentiometer wire AB having length L and resistance 12 r is joined to a cell D of emf ɛ and internal resistance r. A cell C having emf ɛ/2 and internal resistance 3r is connected. The length AJ at which the galvanometer as shown in fig. shows no deflection is :



Ans. (4)

Sol.
$$i = \frac{\varepsilon}{13r}$$

 $i\left(\frac{x}{L}12r\right) = \frac{\varepsilon}{2}$
 $\frac{\varepsilon}{13r}\left[\frac{x}{L}.12r\right] = \frac{\varepsilon}{2} \implies \boxed{x = \frac{13L}{24}}$

20. An insulating thin rod of length ℓ has a x linear charge density $p(x) = \rho_0 \frac{x}{\ell}$ on it. The rod is rotated about an axis passing through the origin (x = 0) and perpendicular to the rod. If the rod makes n rotations per second, then the time averaged magnetic moment of the rod is :

(1)
$$\frac{\pi}{4}$$
 np ℓ^3 (2) np ℓ^3 (3) π np ℓ^3 (4) $\frac{\pi}{3}$ np ℓ^3

Ans. (1)

Sol. :
$$M = NIA$$

da = $\lambda dx \& A = \pi x^2$

$$\int dm = \int (x) \frac{\rho_0 x}{\ell} dx . \pi x^2$$
$$M = \frac{n\rho_0 \pi}{\ell} . \int_0^\ell x^3 . dx = \frac{n\rho_0 \pi}{\ell} . \left[\frac{L^4}{4} \right]$$
$$M = \frac{n\rho_0 \pi \ell^3}{4} \text{ or } \frac{\pi}{4} n\rho \ell^3$$



21. Two guns A and B can fire bullets at speeds 1 km/s and 2 km/s respectively. From a point on a horizontal ground, they are fired in all possible directions. The ratio of maximum areas covered by the bullets fired by the two guns, on the ground is :

 $(1) 1 : 2 \quad (2) 1 : 4 \quad (3) 1 : 8 \quad (4) 1 : 16$

Ans. (4)

Sol. $R = \frac{u^{2} \sin 2\theta}{g}$ $A = \pi R^{2}$ $A \propto R^{2}$ $A \propto u^{4}$ $\frac{A_{1}}{A_{2}} = \frac{u_{1}^{4}}{u_{2}^{4}} = \left[\frac{1}{2}\right]^{4} = \frac{1}{16}$

22. A string of length 1 m and mass 5 g is fixed at both ends. The tension in the string is 8.0 N. The siring is set into vibration using an external vibrator of frequency 100 Hz. The separation between successive nodes on the string is close to :

(1) 16.6 cm	(2) 20.0 cm
(3) 10.0 cm	(4) 33.3 cm

Ans. (2)

Sol. Velocity of wave on string

$$V = \sqrt{\frac{T}{\mu}} = \sqrt{\frac{8}{5} \times 1000} = 40 \text{ m/s}$$

Now, wavelength of wave $\lambda = \frac{v}{n} = \frac{40}{100} m$

Separation b/w successive nodes, $\frac{\lambda}{2} = \frac{20}{100}$ m = 20 cm

23. A train moves towards a stationary observer with speed 34 m/s. The train sounds a whistle and its frequency registered by the observer is f_1 . If the speed of the train is reduced to 17 m/s, the frequency registered is f_2 . If speed of sound is 340 m/s, then the ratio f_1/f_2 is :

(1) 18/17 (2) 19/18 (3) 20/19 (4) 21/20 Ans. (2)

Sol.
$$f_{app} = f_0 \left[\frac{v_2 \pm v_0}{v_2 \mp v_s} \right]$$

 $f_1 = f_0 \left[\frac{340}{340 - 34} \right]$
 $f_2 = f_0 \left[\frac{340}{340 - 17} \right]$
 $\frac{f_1}{f_2} = \frac{340 - 17}{340 - 34} = \frac{323}{306} \implies \frac{f_1}{f_2} = \frac{19}{18}$

24. In an electron microscope, the resolution that can be achieved is of the order of the wavelength of electrons used. To resolve a width of 7.5×10^{-12} m, the minimum electron energy required is close to :

Ans. (3)

Sol.
$$\lambda = \frac{h}{p}$$
 { $\lambda = 7.5 \times 10^{-12}$ }
 $P = \frac{h}{\lambda}$
 $KE = \frac{P^2}{2m} = \frac{(h/\lambda)^2}{2m} = \frac{\left\{\frac{6.6 \times 10^{-34}}{7.5 \times 10^{-12}}\right\}}{2 \times 9.1 \times 10^{-31}}J$

$$KE = 25 Kev$$

25. A homogeneous solid cylindrical roller of radius R and mass M is pulled on a cricket pitch by a horizontal force. Assuming rolling without slipping, angular acceleration of the cylinder is:

R

(1)
$$\frac{3F}{2m R}$$
 (2) $\frac{F}{3m}$

(3)
$$\frac{2F}{3m R}$$
 (4) $\frac{F}{2m R}$

Ans. (3)

Sol.

$$FR = \frac{3}{2}MR^{2}\alpha$$

$$\alpha = \frac{2F}{3MR}$$

Ε



26. A plano convex lens of refractive index μ_1 and focal length f_1 is kept in contact with another plano concave lens of refractive index μ_2 and focal length f_2 . If the radius of curvature of their spherical faces is R each and $f_1 = 2f_2$, then μ_1 and μ_2 are related as :

(1)
$$\mu_1 + \mu_2 = 3$$

(2) $2\mu_1 - \mu_2 = 1$
(3) $2\mu_1 - \mu_2 = 1$

(3)
$$2\mu_2 - \mu_1 = 1$$

$$(4) \ 3\mu_2 - 2\mu_1 = 1$$

Ans. (2)

Sol.
$$\frac{1}{2f_2} = \frac{1}{f_1} = (\mu_1 - 1) \left(\frac{1}{\infty} - \frac{1}{-R} \right)$$

 $\frac{1}{f_2} = (\mu_2 - 1) \left(\frac{1}{-R} - \frac{1}{\infty} \right)$
 $\frac{(\mu_1 - 1)}{R} = \frac{(\mu_2 - 1)}{2R}$
 $2\mu_2 - \mu_2 = 1$

27. Two electric dipoles, A, B with respective dipole moments $\vec{d}_A = -4qa\hat{i}$ and $\vec{d}_B = -2qa\hat{i}$ placed on the x-axis with a separation R, as shown in the figure

$$\xrightarrow{A} \xrightarrow{R} \xrightarrow{R} \xrightarrow{K} X$$

The distance from A at which both of them produce the same potential is :

(1)
$$\frac{\sqrt{2R}}{\sqrt{2}+1}$$

(2)
$$\frac{R}{\sqrt{2}+1}$$

(3)
$$\frac{\sqrt{2}R}{\sqrt{2}-1}$$

(4)
$$\frac{R}{\sqrt{2}-1}$$

Ans. (3)

Е

Sol.
$$V = \frac{4qa}{(R+x)} = \frac{2qa}{(x^2)}$$
$$\sqrt{2}x = R + x$$
$$x = \frac{R}{\sqrt{2} - 1}$$
$$\overrightarrow{4qa} = 2qa$$
$$dist = \frac{R}{\sqrt{2} - 1} + R = \frac{\sqrt{2}R}{\sqrt{2} - 1}$$

28. In the given circuit the cells have zero internal resistance. The currents (in Amperes) passing through resistance R₁, and R₂ respectively, are:

$$R_{1} \ge 20\Omega R_{2} \ge 20\Omega$$

$$R_{1} \ge 20\Omega R_{2} \ge 20\Omega$$
(1) 2, 2
(2) 0,1
(3) 1,2
(4) 0.5,0

Ans. (4)

Sol.
$$i_1 = \frac{10}{20} = 0.5A$$

 $i_2 = 0$

29. In the cube of side 'a' shown in the figure, the vector from the central point of the face ABOD to the central point of the face BEFO will be:





Sol.
$$\vec{r}_g = \frac{a}{2}\hat{i} + \frac{a}{2}\hat{k}$$

 $\vec{r}_H = \frac{a}{2}\hat{j} + \frac{a}{2}\hat{k}$
 $\vec{r}_H - \vec{r}_g = \frac{a}{2}(\hat{j} - \hat{i})$

30. Three Carnot engines operate in series between a heat source at a temperature T_1 and a heat sink at temperature T_4 (see figure). There are two other reservoirs at temperature T_2 , and T_3 , as shown, with $T_2 > T_2 > T_3 > T_4$. The three engines are equally efficient if:

$$\begin{array}{c} & & \\ & &$$

(1)
$$T_2 = (T_1^2 T_4)^{1/3}; T_3 = (T_1 T_4^2)^{1/3}$$

(2) $T_2 = (T_1 T_4^2)^{1/3}; T_3 = (T_1^2 T_4)^{1/3}$
(3) $T_2 = (T_1^3 T_4)^{1/4}; T_3 = (T_1 T_4^3)^{1/4}$
(4) $T_2 = (T_1 T_4)^{1/2}; T_2 = (T_1^2 T_4)^{1/3}$

Ans. (1)

Sol.
$$t_1 = 1 - \frac{T_2}{T_1} = 1 - \frac{T_2}{T_2} = 1 - \frac{T_4}{T_3}$$

$$\Rightarrow \frac{T_2}{T_1} = \frac{T_3}{T_4} = \frac{T_4}{T_3}$$

$$\Rightarrow T_2 = \sqrt{T_1 T_3} = \sqrt{T_1 \sqrt{T_2 T_4}}$$

$$T_3 = \sqrt{T_2 T_4}$$

$$T_2^{3/4} = \sqrt{T_1^{1/2}} T_4^{1/4}$$

$$T_2 = T_1^{2/3} T_4^{1/3}$$

8



Е

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Thursday 10th JANUARY, 2019) TIME : 9 : 30 AM To 12 : 30 PM CHEMISTRY

1. Two pi and half sigma bonds are present in: (1) N_2^+ (2) N_2 (3) O_2^+ (4) O_2

Ans. (1)

Sol.

$$N_2^{\oplus} \Rightarrow BO = 2.5 \Rightarrow \left[\pi - Bond = 2 \& \sigma - Bond = \frac{1}{2}\right]$$

$$\begin{split} N_2 &\Rightarrow B.O. = 3.0 \Rightarrow [\pi\text{-Bond} = 2 \& \sigma\text{-Bond} = 1] \\ O_2^{\oplus} &= B.O. \Rightarrow 2.5 \Rightarrow [\pi\text{-Bond} = 1.5 \& \sigma\text{-Bond} = 1] \\ O_2 &\Rightarrow B.O. \Rightarrow 2 \Rightarrow [\pi\text{-Bond} \Rightarrow 1 \& \sigma\text{-Bond} = 1] \\ \textbf{2.} \quad \text{The chemical nature of hydrogen preoxide is :-} \end{split}$$

- (1) Oxidising and reducing agent in acidic medium, but not in basic medium.
- (2) Oxidising and reducing agent in both acidic and basic medium
- (3) Reducing agent in basic medium, but not in acidic medium
- (4) Oxidising agent in acidic medium, but not in basic medium.

Ans. (2)

- Sol. H_2O_2 act as oxidising agent and reducing agent in acidic medium as well as basic medium. H_2O_2 Act as oxidant :- $H_2O_2 + 2H^{\oplus} + 2e^{\Theta} \rightarrow 2H_2O$ (In acidic medium) $H_2O + 2e^{\Theta} \rightarrow 2OH^{\Theta}$ (In basic medium) H_2O_2 Act as reductant :-
 - $$\begin{split} H_2O_2 &\rightarrow 2H^+ + O_2 + 2e^{\Theta} \text{ (In acidic medium)} \\ H_2O_2 + 2OH^{\Theta} &\rightarrow 2H_2O + O_2 + 2e^{\Theta} \text{ (In basic medium)} \end{split}$$
- **3.** Which dicarboxylic acid in presence of a dehydrating agent is least reactive to give an anhydride :



Sol.	Adipic acid CO ₂ H–(CH ₂	$)_4$ -CO ₂ H $\xrightarrow{\text{dehydrating}}_{\text{agent}}$
4.	7 membered cyclic anhydride (Very unstable) Which premitive unit cell has unequal edge lenghs ($a \neq b \neq c$) and all axial angles differetn from 90°?	
	(1) Tetragonal	(2) Hexagonal
	(3) Monoclinic	(4) Triclinic
Ans.	(4)	
Sol.	In Triclinic unit cell	
	$a \neq b \neq c \& \alpha \neq \beta \neq \gamma$	≠ 90°
5.	Wilkinson catalyst is :	
	(1) $[(Ph_3P)_3RhCl]$	$(\mathrm{Et} = \mathrm{C}_{2}\mathrm{H}_{5})$
	(2) $[Et_3P)_3IrCl]$	
	(3) $[Et_3P)_3RhCl]$	
	(4) $[Ph_3P)_3IrCl]$	
Ans.	(1)	
Sol.	Wilkinsion catalyst is [(ph ₃ P) ₃ RhCl]
6.	The total number of isotopes of hydrogen and number of radioactive isotopes among them,	
	(1) 2 and 0	(2) 3 and 2
	(1) 2 and 0 (3) 3 and 1	(4) 2 and 1
Ang	(3) 5 and 1	(+) 2 and 1
Sol.	Total number of isoton	es of hydrogen is 3
	\rightarrow ¹ U ² U or ² D ³ U or ³ T	
	\rightarrow ₁ H, ₁ H or ₁ D, ₁ H or ₁ I	
	and only ${}_{1}^{3}$ H or ${}_{1}^{3}$ T is an	n Radioactive element.
7.	The major product of the following reaction is	
	Ph Br KOH all	$\frac{c (excess)}{\Delta}$



(1)

1



- Sol. Example of E_2 elimination and conjugated diene is formed with phenyl ring in conjugation which makes it very stable.
- 8. The total number of isomers for a square planar complex [M(F)(Cl)(SCN)(NO₂)] is :
 - (1) 12 (2) 8 (3) 16 (4) 4
- Ans. (1)
- **Sol.** The total number of isomers for a square planar complex $[M(F)(Cl)(SCN)(NO_2)]$ is 12.



- 9. Hall-Heroult's process is given by "
 - (1) $\operatorname{Cr}_2\operatorname{O}_3 + 2\operatorname{Al} \rightarrow \operatorname{Al}_2\operatorname{O}_3 + 2\operatorname{Cr}$
 - (2) $\operatorname{Cu}^{2+}(\operatorname{aq.}) + \operatorname{H}_2(g) \rightarrow \operatorname{Cu}(s) + 2\operatorname{H}^+(\operatorname{aq})$
 - (3) $ZnO + C \xrightarrow{Coke, 1673K} Zn + CO$
 - (4) $2Al_2O_3 + 3C \rightarrow 4Al + 3CO_2$
- Ans. (4)
- Sol. In Hall-Heroult's process is given by $2Al_2O_3 + 3C \longrightarrow 4Al + 3CO_2$ $2Al_2O_3(\ell) \rightleftharpoons 4Al^{3+}(\ell) + 6O^{2\Theta}(\ell)$ At cathode :- $4Al_{(\ell)}^{3+} + 12e^{\Theta} \rightarrow 4Al(\ell)$ At cathode : $6O_{(\ell)}^{2\Theta} \rightarrow 3O_2(g) + 12e^{\Theta}$ $3C + 3O_2 \rightarrow 3CO_2 (\uparrow)$ 10. The value of K_p/K_C for the following reactions at 300K are, respectively : (At 300K, RT = 24.62 dm³atm mol⁻¹) $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$ $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ $N_2(g) + 3H_2(g) \oiint 2NH_3(g)$ (1) 1, 24.62 dm³atm mol⁻¹, $606.0 \text{ dm}^6 \text{atm}^2 \text{mol}^{-2}$
 - (2) 1, 4.1 \times 10⁻² dm⁻³atm⁻¹ mol⁻¹, 606.0 dm⁶ atm² mol⁻²
 - (3) 606.0 dm⁶atm²mol⁻²,
 - $1.65 \times 10^{-3} \text{ dm}^3 \text{atm}^{-2} \text{ mol}^{-1}$

(4) 1, 24.62 dm³atm mol⁻¹,
$$1.65 \times 10^{-3} \text{ dm}^{-6} \text{atm}^{-2} \text{ mol}^2$$

Ans. (4)

2

- Sol. $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$ $\frac{k_p}{k_c} = (RT)^{\Delta n_g} = (RT)^0 = 1$ $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ $\frac{k_p}{k_c} = (RT)^1 = 24.62$ $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$ $\frac{k_p}{k_c} = (RT)^{-2} = \frac{1}{(RT)^2} = 1.65 \times 10^{-3}$
- 11. If dichloromethane (DCM) and water (H_2O) are used for differential extraction, which one of the following statements is correct ?
 - (1) DCM and H_2O would stay as lower and upper layer respectively in the S.F.
 - (2) DCM and H₂O will be miscible clearly
 - (3) DCM and H₂O would stay as upper and lower layer respectively in the separating funnel (S.F.)
 - (4) DCM and H₂O will make trubid/colloidal mixture

Ans. (1)

- 12. The type of hybridisation and number of lone pair(s) of electrons of Xe in XeOF₄, respectively, are :
 - (1) sp^3d and 1
 - (2) sp^3d and 2
 - (3) $sp^{3}d^{2}$ and 1
 - (4) $sp^{3}d^{2}$ and 2

Ans. (3)

Sol.
$$\underset{F}{\overset{F}{\longrightarrow}} \underset{F}{\overset{H}{\longrightarrow}} \underset{F}{\overset{F}{\longrightarrow}} sp^{3}d^{2} \Rightarrow [5\sigma \text{-bond } +1 \text{ l.p.}]$$

13. The metal used for making X-ray tube window is :

(1) Mg (2) Na (3) Ca (4) Be

Ans. (4)

Sol. "Be" Metal is used in x-ray window is due to transparent to x-rays.

Collegebatch.com

E

14. Consider the given plots for a reaction obeying Arrhenius equation $(0^{\circ}C < T < 300^{\circ}C)$: (k and E_a are rate constant and activation energy, respectively)



Choose the correct option :

- (1) Both I and II are wrong
- (2) I is wrong but II is right
- (3) Both I and II are correct
- (4) I is right but II is wrong

Ans. (4)

- **Sol.** On increasing E_a , K decreases
- **15.** Water filled in two glasses A and B have BOD values of 10 and 20, respectively. The correct statement regarding them, is :
 - (1) A is more polluted than B
 - (2) A is suitable for drinking, whereas B is not
 - (3) B is more polluted than A
 - (4) Both A and B are suitable for drinking
- Ans. (3)
- **Sol.** Two glasses "A" and "B" have BOD values 10 and "20", respectively.

Hence glasses "B" is more polluted than glasses "A".

16. The increasing order of the pKa values of the following compounds is :



Sol. Acidic strength is inversely proportional to pka.



17. Liquids A and B form an ideal solution in the entire composition range. At 350 K, the vapor pressures of pure A and pure B are 7×10^3 Pa and 12×10^3 Pa, respectively. The composition of the vapor in equilibrium with a solution containing 40 mole percent of A at this temperature is :

(1)
$$x_A = 0.37$$
; $x_B = 0.63$
(2) $x_A = 0.28$; $x_B = 0.72$
(3) $x_A = 0.76$; $x_B = 0.24$
(4) $x_A = 0.4$; $x_B = 0.6$

Ans. (2)

Sol.
$$y_A = \frac{P_A}{P_{Total}} = \frac{P_A^o x_A}{P_A^o x_A \times p_B^o x_B}$$

$$= \frac{7 \times 10^{3} \times 0.4}{7 \times 10^{3} \times 0.4 + 12 \times 10^{3} \times 0.6}$$

$$=\frac{2.8}{10}=0.28$$

 $y_{\rm B} = 0.72$

18. Consider the following reduction processes :

$$Zn^{2+} + 2e^{-} \rightarrow Zn(s); E^{\circ} = -0.76 V$$

 $Ca^{2+} + 2e^{-} \rightarrow Ca(s); E^{\circ} = -2.87 V$
 $Mg^{2+} + 2e^{-} \rightarrow Mg(s); E^{\circ} = -2.36 V$
 $Ni^{2+} + 2e^{-} \rightarrow Ni(s); E^{\circ} = -0.25 V$

The reducing power of the metals increases in the order :

Ans. (2)

Sol. Higher the oxidation potential better will be reducing power.

3



19. The major product of the following reaction is:



- (1) Boron(2) Carbon(3) Lithium(4) Beryllium
- Ans. (4)
- **Sol.** E.N. of Al = $(1.5) \ge$ Be (1.5)

21. The decreasing order of ease of alkaline hydrolysis for the following esters is :



Ans. (2)

Sol. More is the electrophilic character of carbonyl group of ester faster is the alkaline hydrolysis.

22. A process has $\Delta H = 200 \text{ Jmol}^{-1}$ and

 $\Delta S = 40 \text{ JK}^{-1}\text{mol}^{-1}$. Out of the values given below, choose the minimum temperature above which the process will be spontaneous :

(1) 5 K (2) 4 K (3) 20 K (4) 12 K Ans. (1) Sol. $\Delta G = \Delta H - T\Delta S$ _ $\Delta H = 200$

$$T = \frac{\Delta H}{\Delta S} = \frac{200}{40} = 5K$$

4

collegebatch.com

E

23. Which of the graphs shown below does not represent the relationship between incident light and the electron ejected form metal surface ?



(4) Hydrogenation of vegetable oils

Ans. (3)

E

- **Sol.** Then is no catalyst is required for combustion of coal.
- 25. The effect of lanthanoid contraction in the lanthanoid series of elements by and large means :
 - (1) decrease in both atomic and ionic radii
 - (2) increase in atomic radii and decrease in ionic radii
 - (3) increase in both atomic and ionic radii
 - (4) decrease in atomic radii and increase in ionic radii

Ans. (1)

- **Sol.** Due to Lanthanoid contraction both atomic radii and ionic radii decreases gradually in the lanthanoid series.
- **26.** The major product formed in the reaction given below will be :



5

collegebatch.com

27. The correct structure of product 'P' in the following reaction is :



Ans. (1)

Sol. Asn-Ser is dipeptide having following structure



$$Asn - Ser + (CH_3CO)_2 O \xrightarrow{NEt_3} P$$
excess

P is



28. Which hydrogen in compound (E) is easily replaceable during bromination reaction in presence of light :

$$CH_{3} - CH_{2} - CH_{\beta} = CH_{2} \\ \alpha$$

(1) β – hydrogen

- (2) γ hydrogen
- (3) δ hydrogen
- (4) α hydrogen

Ans. (2)

29. The major product 'X' formed in the following reaction is :

$$(1) \xrightarrow{O} CH_2-C-OCH_3 \xrightarrow{\text{NaBH}_4} X$$

$$(1) \xrightarrow{O} CH_2CH_2OH$$

$$(2) \xrightarrow{O} CH_2-C-H$$

$$(3) \xrightarrow{O} CH_2CH_2OH$$

$$(4) \xrightarrow{O} CH_2-C-OCH_3$$

Ans. (4)

collegebatch.com

- 30. A mixture of 100 m mol of Ca(OH)₂ and 2g of sodium sulphate was dissolved in water and the volume was made up to 100 mL. The mass of calcium sulphate formed and the concentration of OH⁻ in resulting solution, respectively, are : (Molar mass of Ca(OH)₂, Na₂SO₄ and CaSO₄ are 74, 143 and 136 g mol⁻¹, respectively; K_{sp} of Ca(OH)₂ is 5.5 × 10⁻⁶)
 - (1) 1.9 g, 0.14 mol L⁻¹
 - (2) 13.6 g, 0.14 mol L^{-1}
 - (3) 1.9 g, 0.28 mol L⁻¹
 - (4) 13.6 g, 0.28 mol L⁻¹

Ans. (3)

- - $w_{CaSO_4} = 14 \times 10^{-3} \times 136 = 1.9 gm$

$$[OH^{-}] = \frac{28}{100} = 0.28M$$



Ε

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Thursday 10th JANUARY, 2019) TIME : 2 : 30 PM To 5 : 30 PM MATHEMATICS

- Let $z = \left(\frac{\sqrt{3}}{2} + \frac{i}{2}\right)^3 + \left(\frac{\sqrt{3}}{2} \frac{i}{2}\right)^3$. If R(z) and I[z] 1. respectively denote the real and imaginary parts of z, then : (1) R(z) > 0 and I(z) > 0(2) R(z) < 0 and I(z) > 0(3) R(z) = -3(4) I(z) = 0Ans. (4) S **Sol.** $z = \left(\frac{\sqrt{3} + i}{2}\right)^3 + \left(\frac{\sqrt{3} - i}{2}\right)^3$ $z = (e^{i\pi/6})^5 + (e^{-i\pi/6})^5$ $= e^{i5\pi/6} + e^{-i5\pi/6}$ $=\cos\frac{5\pi}{6} + i\frac{\sin 5\pi}{6} + \cos\left(\frac{-5\pi}{6}\right) + i\sin\left(\frac{-5\pi}{6}\right)$ $= 2 \cos \frac{5\pi}{6} < 0$ I(z) = 0 and Re(z) < 0Option (4) Let $a_1, a_2, a_3, ..., a_{10}$ be in G.P. with $a_i > 0$ for 2. 4. i = 1, 2, ..., 10 and S be the set of pairs (r, k), r $k \in N$ (the set of natural numbers) for which $\log_e a_2^r a_3^k$ $\log_e a_5^r a_6^k$ $\log_e a_1^r a_2^k$ $\frac{\log_{e} a_{3}^{r} a_{4}^{k}}{\log_{e} a_{6}^{r} a_{7}^{k}} = 0$ $\log_e a_4^r a_5^k$ $\log_{e} a_{7}^{r} a_{8}^{k}$ Then the number of elements in S, is : (1) Infinitely many (2) 4(3) 10(4) 2Ans. (1) Sol. Apply $C_3 \rightarrow C_3 - C_2$ $C_2 \rightarrow C_2 - C_1$ We get D = 0Option (1)
- 3. The positive value of λ for which the co-efficient of x^2 in the expression

$$x^2 \left(\sqrt{x} + \frac{\lambda}{x^2}\right)^{10}$$
 is 720, is :

(1)
$$\sqrt{5}$$
 (2) 4

(3)
$$2\sqrt{2}$$
 (4) 3

Ans. (2)

ol.
$$x^{2}\left({}^{10}C_{r}\left(\sqrt{x}\right)^{10-r}\left(\frac{\lambda}{x^{2}}\right)^{r}\right)$$

 $x^{2}\left[{}^{10}C_{r}\left(x\right)^{\frac{10-r}{2}}\left(\lambda\right)^{r}\left(x\right)^{-2r}\right]$

$$x^{2} \begin{bmatrix} {}^{10}C_{r} \lambda^{r} x^{\frac{10-5r}{2}} \end{bmatrix}$$

$$\therefore r = 2$$

Hence, ${}^{10}C_{2} \lambda^{2} = 720$

$$\lambda^{2} = 16$$

$$\lambda = \pm 4$$

Option (2)
The value of $\cos \frac{\pi}{2^2} \cdot \cos \frac{\pi}{2^3} \cdot \dots \cdot \cos \frac{\pi}{2^{10}} \cdot \sin \frac{\pi}{2^{10}}$

is :

(1)
$$\frac{1}{256}$$
 (2) $\frac{1}{2}$

(3)
$$\frac{1}{512}$$
 (4) $\frac{1}{1024}$

Ans. (3)

Sol.
$$2\sin\frac{\pi}{2^{10}}\cos\frac{\pi}{2^{10}}\dots\cos\frac{\pi}{2^{2}}$$

 $\frac{1}{2^{9}}\sin\frac{\pi}{2} = \frac{1}{512}$
Option (3)


6.

5. The value of
$$\int_{-\pi/2}^{\pi/2} \frac{dx}{[x] + [\sin x] + 4}$$
, where [t]

denotes the greatest integer less than or equal to t, is :

(1)
$$\frac{1}{12}(7\pi+5)$$
 (2) $\frac{3}{10}(4\pi-3)$
(3) $\frac{1}{12}(7\pi-5)$ (4) $\frac{3}{20}(4\pi-3)$

Ans. (4)

Sol.
$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{dx}{[x] + [\sin x] + 4}$$
$$= \int_{-\frac{\pi}{2}}^{-1} \frac{dx}{-2 - 1 + 4} + \int_{-1}^{0} \frac{dx}{-1 - 1 + 4}$$
$$+ \int_{0}^{1} \frac{dx}{0 + 0 + 4} + \int_{1}^{\frac{\pi}{2}} \frac{dx}{1 + 0 + 4}$$
$$\int_{-\frac{\pi}{2}}^{-1} \frac{dx}{1} + \int_{-1}^{0} \frac{dx}{2} + \int_{0}^{1} \frac{dx}{4} + \int_{1}^{\frac{\pi}{2}} \frac{dx}{5}$$
$$\left(-1 + \frac{\pi}{2}\right) + \frac{1}{2}(0 + 1) + \frac{1}{4} + \frac{1}{5}\left(\frac{\pi}{2} - 1\right)$$

$$\left(-1+\frac{\pi}{2}\right) + \frac{1}{2}(0+1) + \frac{1}{4} + \frac{1}{5}\left(-1+\frac{1}{2}+\frac{1}{4}-\frac{1}{5}+\frac{\pi}{2}+\frac{\pi}{10}-\frac{1}{20}+\frac{1}{5}+\frac{\pi}{2}+\frac{\pi}{10}-\frac{1}{20}+\frac{1}{20}+\frac{1}{10$$

 $\frac{-9}{20} + \frac{3\pi}{5}$ Option (4)

20

in any shot, is 1/3, then the minimum number of independent shots at the target required by him so that the probability of hitting the target at least once is greater than $\frac{5}{6}$, is : (1) 6(2) 5(3) 4(4) 3 Ans. (2) **Sol.** $1 - {}^{n}C_{0}\left(\frac{1}{3}\right)^{0}\left(\frac{2}{3}\right)^{n} > \frac{5}{6}$ $\frac{1}{6} > \left(\frac{2}{3}\right)^n \implies 0.1666 > \left(\frac{2}{3}\right)^n$ $n_{min} = 5 \implies Option (2)$ 7. If mean and standard deviation of 5 observations x_1 , x_2 , x_3 , x_4 , x_5 are 10 and 3, respectively, then the variance of 6 observations x_1, x_2, \dots, x_5 and -50 is equal to : (1) 582.5(2) 507.5 (3) 586.5 (4) 509.5 Ans. (2) **Sol.** $\overline{\mathbf{x}} = 10 \implies \sum_{i=1}^{5} \mathbf{x}_{i} = 50$

If the probability of hitting a target by a shooter,

i=1
S.D. =
$$\sqrt{\frac{\sum_{i=1}^{5} x_i^2}{5} - (\overline{x})^2} = 8$$

⇒ $\sum_{i=1}^{5} (x_i)^2 = 109$

variance =
$$\frac{\sum_{i=1}^{5} (x_i)^2 + (-50)^2}{6} - \left(\sum_{i=1}^{5} \frac{x_i - 50}{6}\right)$$

Option (2)

8. The length of the chord of the parabola $x^2 = 4y$ having equation $x - \sqrt{2}y + 4\sqrt{2} = 0$ is :

(1)
$$2\sqrt{11}$$
 (2) $3\sqrt{2}$

(3)
$$6\sqrt{3}$$
 (4) $8\sqrt{2}$

Ans. (3)

collegebatch.com

Sol. $x^2 = 4y$ $x - \sqrt{2}y + 4\sqrt{2} = 0$ Solving together we get $x^2 = 4 \left(\frac{x + 4\sqrt{2}}{\sqrt{2}} \right)$ $\sqrt{2}x^2 + 4x + 16\sqrt{2}$ $\sqrt{2}x^2 - 4x - 16\sqrt{2} = 0$ $x_1 + x_2 = 2\sqrt{2}$; $x_1x_2 = \frac{-16\sqrt{2}}{\sqrt{2}} = -16$ Similarly, $\left(\sqrt{2}y - 4\sqrt{2}\right)^2 = 4y$ 10. $2y^2 + 32 - 16y = 4y$ $2y^2 - 20y + 32 = 0$ $y_1 + y_2 = 10$ $y_1 y_2 = 16$ Sol. $\ell_{AB} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ $= \sqrt{\left(2\sqrt{2}\right)^2 + 64 + (10)^2 - 4(16)}$ $=\sqrt{8+64+100-64}$ $=\sqrt{108}=6\sqrt{3}$ Option (3) 9. Let $A = \begin{bmatrix} 2 & b & 1 \\ b & b^2 + 1 & b \\ 1 & b & 2 \end{bmatrix}$ where b > 0. Then the 11. minimum value of $\frac{\det(A)}{h}$ is : $(1)\sqrt{3}$ (2) $-\sqrt{3}$ $(3) - 2\sqrt{3}$ $(4)2\sqrt{3}$ A Ans. (4)

Sol. A = $\begin{bmatrix} 2 & b & 1 \\ b & b^2 + 1 & b \\ 1 & b & 2 \end{bmatrix}$ (b > 0) $|\mathbf{A}| = 2(2b^2 + 2 - b^2) - b(2b - b) + 1 (b^2 - b^2 - 1)$ $|\mathbf{A}| = 2(\mathbf{b}^2 + 2) - \mathbf{b}^2 - 1$ $|A| = b^2 + 3$ $\frac{|\mathbf{A}|}{\mathbf{b}} = \mathbf{b} + \frac{3}{\mathbf{b}} \implies \frac{\mathbf{b} + \frac{3}{\mathbf{b}}}{2} \ge \sqrt{3}$ $b + \frac{3}{b} \ge 2\sqrt{3}$ Option (4) The tangent to the curve, $y = xe^{x^2}$ passing

through the point (1,e) also passes through the point :

(1)
$$\left(\frac{4}{3}, 2e\right)$$
 (2) (2,3e)
(3) $\left(\frac{5}{3}, 2e\right)$ (4) (3,6e)

Ans. (1)

$$y = xe^{x^{2}}$$

$$\frac{dy}{dx}\Big|_{(1,e)} = \left(e \cdot e^{x^{2}} \cdot 2x + e^{x^{2}}\right)\Big|_{(1,e)} = 2 \cdot e + e = 3e$$

$$T : y - e = 3e (x - 1)$$

$$y = 3ex - 3e + e$$

$$y = (3e)x - 2e$$

$$\left(\frac{4}{3}, 2e\right) \text{ lies on it}$$
Option (1)
The number of values of $\theta \in (0, \pi)$ for which the

values of $\theta \in (0,\pi)$ for which the system of linear equations x + 3y + 7z = 0

> -x + 4y + 7z = 0 $(\sin 3\theta)x + (\cos 2\theta)y + 2z = 0$

has a non-trivial solution, is :

- (1) One (2) Three
- (3) Four (4) Two



7 -1 4 7 = 0Sol. 2 $\sin 3\theta \cos 2\theta$ $(8 - 7 \cos 2\theta) - 3(-2 - 7 \sin 3\theta)$ + 7 (- $\cos 2\theta$ - 4 $\sin 3\theta$) = 0 $14 - 7 \cos 2\theta + 21 \sin 3\theta - 7 \cos 2\theta$ $-28 \sin 3\theta = 0$ $14 - 7 \sin 3\theta - 14 \cos 2\theta = 0$ $14 - 7 (3 \sin \theta - 4 \sin^3 \theta) - 14 (1 - 2 \sin^2 \theta) = 0$ $-21 \sin \theta + 28 \sin^3 \theta + 28 \sin^2 \theta = 0$ $7 \sin \theta \left[-3 + 4 \sin^2 \theta + 4 \sin \theta \right] = 0$ $\sin \theta$. $4 \sin^2 \theta + 6 \sin \theta - 2 \sin \theta - 3 = 0$ $2 \sin \theta (2 \sin \theta + 3) - 1 (2 \sin \theta + 3) = 0$ $\sin \theta = \frac{-3}{2}; \sin \theta = \frac{1}{2}$ Hence, 2 solutions in $(0, \pi)$ Option (4) If $\int_{0}^{x} f(t) dt = x^{2} + \int_{0}^{1} t^{2} f(t) dt$, then f'(1/2) is : 12. (1) $\frac{6}{25}$ (2) $\frac{24}{25}$ (4) $\frac{4}{5}$ (3) $\frac{18}{25}$ Ans. (2) **Sol.** $\int_{0}^{x} f(t) dt = x^{2} + \int_{0}^{1} t^{2} f(t) dt$ $f'\left(\frac{1}{2}\right) = ?$ Differentiate w.r.t. 'x' $f(x) = 2x + 0 - x^2 f(x)$ $f(x) = \frac{2x}{1+x^2} \implies f'(x) = \frac{(1+x^2)2 - 2x(2x)}{(1+x^2)^2}$ $f'(x) = \frac{2x^2 - 4x^2 + 2}{(1 + x^2)^2}$ $f'\left(\frac{1}{2}\right) = \frac{2-2\left(\frac{1}{4}\right)}{\left(1+\frac{1}{4}\right)^2} = \frac{\left(\frac{3}{2}\right)}{\frac{25}{16}} = \frac{48}{50} = \frac{24}{25}$ Option (2)

13. Let f: (-1,1)→R be a function defined by f(x) = max {-|x|, -√1-x²}. If K be the set of all points at which f is not differentiable, then K has exactly :
(1) Three elements (2) One element

(3) Five elements (4) Two elements

Sol.
$$f: (-1, 1) \rightarrow R$$

$$f(x) = \max\left\{-|x|, -\sqrt{1-x^2}\right\}$$



Non-derivable at 3 points in (-1, 1)Option (1)

- **14.** Let $S = \left\{ (x, y) \in R^2 : \frac{y^2}{1+r} \frac{x^2}{1-r} = 1 \right\}$, where
 - $r \neq \pm 1$. Then S represents :
 - (1) A hyperbola whose eccentricity is $\frac{2}{\sqrt{r+1}}$, where 0 < r < 1.
 - (2) An ellipse whose eccentricity is $\frac{1}{\sqrt{r+1}}$, where r > 1
 - (3) A hyperbola whose eccentricity is $\frac{2}{\sqrt{1-r}}$, when 0 < r < 1.
 - (4) An ellipse whose eccentricity is $\sqrt{\frac{2}{r+1}}$, when r > 1

4



Ans. (4) **Sol.** $\frac{y^2}{1+r} - \frac{x^2}{1-r} = 1$ for r > 1, $\frac{y^2}{1+r} + \frac{x^2}{r-1} = 1$ $e = \sqrt{1 - \left(\frac{r-1}{r+1}\right)}$ $=\sqrt{\frac{(r+1)-(r-1)}{(r+1)}}$ $=\sqrt{\frac{2}{r+1}}=\sqrt{\frac{2}{r+1}}$ Option (4)**15.** If $\sum_{r=0}^{25} \left\{ {}^{50}C_r \cdot {}^{50-r}C_{25-r} \right\} = K \left({}^{50}C_{25} \right)$, then K is (4) 2²⁴ (1) $2^{25} - 1$ (2) $(25)^2$ (3) 2^{25} Ans. (3) **Sol.** $\sum_{r=0}^{25} {}^{50}C_r \cdot {}^{50-r}C_{25-r}$ $=\sum_{r=0}^{25} \frac{50!}{r! (50-r)!} \times \frac{(50-r)!}{(25)! (25-r)!}$ $= \sum_{n=1}^{25} \frac{50!}{25! \cdot 25!} \times \frac{25!}{(25-r)! \cdot (r!)}$ $= {}^{50}C_{25}\sum_{r}^{25}{}^{25}C_{r} = (2^{25}){}^{50}C_{25}$ \therefore K = 2²⁵ Option (3) 16. Let N be the set of natural numbers and two functions f and g be defined as $f,g: N \rightarrow N$ such that : $f(n) = \begin{pmatrix} \frac{n+1}{2} & \text{if } n \text{ is odd} \\ \frac{n}{2} & \text{if } n \text{ is even} \end{pmatrix}$ and $g(n) = n - (-1)^n$. The fog is : (1) Both one-one and onto (2) One-one but not onto

- (3) Neither one-one nor onto
- (4) onto but not one-one

Ans. (4)

Sol. $f(x) = \begin{cases} \frac{n+1}{2} & n \text{ is odd} \\ n/2 & n \text{ is even} \end{cases}$ $g(x) = n - (-1)^n \begin{cases} n+1 ; n \text{ is odd} \\ n-1 ; n \text{ is even} \end{cases}$ $f(g(n)) = \begin{cases} \frac{n}{2}; & n \text{ is even} \\ \frac{n+1}{2}; & n \text{ is odd} \end{cases}$: many one but onto Option (4) 17. The values of λ such that sum of the squares of the roots of the quadratic equation, $x^{2} + (3 - \lambda) x + 2 = \lambda$ has the least value is : (2) $\frac{4}{9}$ (1) 2(3) $\frac{15}{9}$ (4) 1 Ans. (1) **Sol.** $\alpha + \beta = \lambda - 3$ $\alpha\beta = 2 - \lambda$ $\alpha^{2} + \beta^{2} = (\alpha + \beta)^{2} - 2\alpha\beta = (\lambda - 3)^{2} - 2(2 - \lambda)$ $= \lambda^2 + 9 - 6\lambda - 4 + 2\lambda$ $= \lambda^2 - 4\lambda + 5$ $= (\lambda - 2)^2 + 1$ $\therefore \lambda = 2$ Option (1)

18. Two vertices of a triangle are (0,2) and (4,3).If its orthocentre is at the origin, then its third vertex lies in which quadrant ?

- (1) Fourth
- (2) Second
- (3) Third
- (4) First

Ans. (2)



collegebatch.com

- **Sol.** $m_{BD} \times m_{AD} = -1 \implies \left(\frac{3-2}{4-0}\right) \times \left(\frac{b-0}{a-0}\right) = -1$
 - \Rightarrow b + 4a = 0(i)



 $m_{AB} \times m_{CF} = -1 \implies \left(\frac{(b-2)}{a-0}\right) \times \left(\frac{3}{4}\right) = -1$

 $\Rightarrow 3b - 6 = -4a \Rightarrow 4a + 3b = 6 \dots (ii)$ From (i) and (ii)

$$a = \frac{-3}{4}, b = 3$$

∴ IInd quadrant.

Option (2)

- **19.** Two sides of a parallelogram are along the lines, x + y = 3 and x y + 3 = 0. If its diagonals intersect at (2,4), then one of its vertex is : (1) (2,6) (2) (2,1)
 - (3) (3,5) (4) (3,6)

Ans. (4)



 $\frac{x_1+0}{2} = 2; x_i = 4 \text{ similarly } y_1 = 5$ C \Rightarrow (4, 5) Now equation of BC is x - y = -1 and equation of CD is x + y = 9

Solving x + y = 9 and x - y = -3Point D is (3, 6) Option (4) 20. Let $\vec{\alpha} = (\lambda - 2)\vec{a} + \vec{b}$ and $\vec{\beta} = (4\lambda - 2)\vec{a} + 3\vec{b}$ be two given vectors where vectors \vec{a} and \vec{b} are non-collinear. The value of λ for which vectors $\vec{\alpha}$ and $\vec{\beta}$ are collinear, is :

$$\begin{array}{cccc} (1) & -3 & (2) & 4 \\ (3) & 3 & (4) & -4 \end{array}$$

Ans. (4)

Sol.
$$\vec{\alpha} = (\lambda - 2)\vec{\alpha} + \vec{b}$$

$$\vec{\beta} = (4\lambda - 2)\vec{\alpha} + 3\vec{b}$$
$$\frac{\lambda - 2}{4\lambda - 2} = \frac{1}{3}$$
$$3\lambda - 6 = 4\lambda - 2$$
$$\boxed{\lambda = -4}$$
$$\therefore \text{ Option (4)}$$

21. The value of $\cot\left(\sum_{n=1}^{19} \cot^{-1}\left(1 + \sum_{p=1}^{n} 2p\right)\right)$ is :

(1)
$$\frac{22}{23}$$
 (2) $\frac{23}{22}$ (3) $\frac{21}{19}$ (4) $\frac{19}{21}$

Ans. (3)

Sol.
$$\cot\left(\sum_{n=1}^{19}\cot^{-1}(1+n(n+1))\right)$$

 $\cot\left(\sum_{n=1}^{19}\cot^{-1}(n^2+n+1)\right) = \cot\left(\sum_{n=1}^{19}\tan^{-1}\frac{1}{1+n(n+1)}\right)$
 $\sum_{n=1}^{19}(\tan^{-1}(n+1)-\tan^{-1}n)$
 $\cot(\tan^{-1}20-\tan^{-1}1) = \frac{\cot A \cot \beta + 1}{\cot \beta - \cot A}$

(Where tanA=20, tanB=1)
$$\frac{1\left(\frac{1}{20}\right)+1}{1-\frac{1}{20}} = \frac{21}{19}$$

 \therefore Option (3)

22. With the usual notation, in $\triangle ABC$, if $\angle A + \angle B = 120^{\circ}$, $a = \sqrt{3} + 1$ and $b = \sqrt{3} - 1$, then the ratio $\angle A : \angle B$, is : (1) 7 : 1 (2) 5 : 3 (3) 9 : 7 (4) 3 : 1



F

JEE (Main) Examination-2019/Evening Session/10-01-2019

24. Consider the following three statements :P : 5 is a prime number.

Q : 7 is a factor of 192.

R : L.C.M. of 5 and 7 is 35.

Then the truth value of which one of the following statements is true ?

(1) $(P \land Q) \lor (\sim R)$

(2)
$$(\sim P) \land (\sim Q \land R)$$

(3) $(\sim P) \lor (Q \land R)$

$$(4) \mathbf{P} \vee (\mathbf{P} \mathbf{V} \mathbf{R})$$

Ans. (4)

- Sol. It is obvious
 - \therefore Option (4)
- 25. On which of the following lines lies the point
 - of intersection of the line, $\frac{x-4}{2} = \frac{y-5}{2} = \frac{z-3}{1}$ and the plane, x + y + z = 2 ?

(1)
$$\frac{x-2}{2} = \frac{y-3}{2} = \frac{z+3}{3}$$

(2) $\frac{x-4}{1} = \frac{y-5}{1} = \frac{z-5}{-1}$
(3) $\frac{x-1}{1} = \frac{y-3}{2} = \frac{z+4}{-5}$

(4)
$$\frac{x+3}{3} = \frac{4-y}{3} = \frac{z+1}{-2}$$

Ans. (3)

Sol. General point on the given line is $x = 2\lambda + 4$ $y = 2\lambda + 5$ $z = \lambda + 3$ Solving with plane, $2\lambda + 4 + 2\lambda + 5 + \lambda + 3 = 2$ $5\lambda + 12 = 2$ $5\lambda = -10$ $\boxed{\lambda = -2}$ \therefore Option (3)

Ans. (1)

Sol.
$$A + B = 120^{\circ}$$



$$\tan\frac{A-B}{2} = \frac{a-b}{a+b}\cot\left(\frac{C}{2}\right)$$

$$= \frac{\sqrt{3} + 1 - \sqrt{3} + 1}{2(\sqrt{3})} \cot(30^\circ) = \frac{1}{\sqrt{3}} \cdot \sqrt{3} = 1$$

$$\frac{A-B}{2} = 45^{\circ} \qquad \Rightarrow \quad A-B = 90^{\circ}$$
$$A+B = 120^{\circ}$$
$$2A = 210^{\circ}$$
$$A = 105^{\circ}$$
$$B = 15^{\circ}$$

 \therefore Option (1)

23. The plane which bisects the line segment joining the points (-3,-3,4) and (3,7,6) at right angles, passes through which one of the following points ?

(1) (4, -1, 7) (2) (4, 1, -2)(3) (-2, 3, 5) (4) (2, 1, 3)Ans. (2)



p : 3(x - 0) + 5(y - 2) + 1(z - 5) = 03x + 5y + z = 15 ∴ Option (2)

26. Let f be a differentiable function such that

$$f'(x) = 7 - \frac{3}{4} \frac{f(x)}{x}, (x > 0) \text{ and } f(1) \neq 4$$

Then
$$\lim_{x\to 0^+} x f\left(\frac{1}{x}\right)$$
:

- (1) Exists and equals 4
- (2) Does not exist
- (3) Exist and equals 0

(4) Exists and equals
$$\frac{4}{7}$$

Ans. (1)

Sol. $f'(x) = 7 - \frac{3}{4} \frac{f(x)}{x}$ (x > 0)

Given
$$f(1) \neq 4$$
 $\lim_{x \to 0^+} xf\left(\frac{1}{x}\right) = ?$

 $\frac{dy}{dx} + \frac{3}{4}\frac{y}{x} = 7 \text{ (This is LDE)}$ $IF = e^{\int \frac{3}{4x}dx} = e^{\frac{3}{4}\ln|x|} = x^{\frac{3}{4}}$ $y.x^{\frac{3}{4}} = \int 7.x^{\frac{3}{4}}dx$

$$y \cdot x^{\frac{3}{4}} = 7 \cdot \frac{x^{\frac{7}{4}}}{\frac{7}{4}} + C$$

$$f(x) = 4x + C.x^{-\frac{3}{4}}$$
$$f\left(\frac{1}{x}\right) = \frac{4}{x} + C.x^{-\frac{3}{4}}$$

$$\lim_{x \to 0^+} xf\left(\frac{1}{x}\right) = \lim_{x \to 0^+} \left(4 + C \cdot x^{\frac{7}{4}}\right) = 4$$

$$\therefore \quad \text{Option (1)}$$

27. A helicopter is flying along the curve given by $y - x^{3/2} = 7$, $(x \ge 0)$. A soldier positioned at the point $\left(\frac{1}{2}, 7\right)$ wants to shoot down the helicopter when it is nearest to him. Then this nearest distance is : (2) $\frac{1}{3}\sqrt{\frac{7}{3}}$ (1) $\frac{1}{2}$ (3) $\frac{1}{6}\sqrt{\frac{7}{3}}$ (4) $\frac{\sqrt{5}}{6}$ Ans. (3) **Sol.** $y - x^{3/2} = 7 \ (x \ge 0)$ $\frac{\mathrm{dy}}{\mathrm{dx}} = \frac{3}{2} \mathrm{x}^{1/2}$ $\left(\frac{3}{2}\sqrt{x}\right)\left(\frac{7-y}{\frac{1}{2}-x}\right) = -1 \qquad \xrightarrow{B} (x, y) \\ A\left(\frac{1}{2}, 7\right)$ $\left(\frac{3}{2}\sqrt{x}\right)\left(\frac{-x^{3/2}}{\frac{1}{2}-x}\right) = -1$ $\frac{3}{2} \cdot x^2 = \frac{1}{2} - x$ $3x^2 = 1 - 2x$ $3x^2 + 2x - 1 = 0$ $3x^2 + 3x - x - 1 = 0$ (x + 1) (3x - 1) = 0 \therefore x = -1 (rejected)

$$x = \frac{1}{3}$$

y = 7 + x^{3/2} = 7 +
$$\left(\frac{1}{3}\right)^{3/2}$$

 $\ell_{AB} = \sqrt{\left(\frac{1}{2} - \frac{1}{3}\right)^2 + \left(\frac{1}{3}\right)^3} = \sqrt{\frac{1}{36} + \frac{1}{27}}$

$$= \sqrt{\frac{3+4}{9\times 12}}$$
$$= \sqrt{\frac{7}{108}} = \frac{1}{6}\sqrt{\frac{7}{3}}$$

Option (3)



28. Ans.	If $\int x^5 e^{-4x^3} dx = \frac{1}{48} e^{-4x^3} f$ constant of integration, (1) $-4x^3 - 1$ (3) $-2x^3 - 1$ (1)	f(x)+C, where C is a then f(x) is equal to : (2) $4x^3 + 1$ (4) $-2x^3 + 1$	29.	The curve amongst the family of curves, represented by the differential equation, $(x^2 - y^2)dx + 2xy dy = 0$ which passes through (1,1) is : (1) A circle with centre on the y-axis (2) A circle with centre on the x-axis (3) An ellipse with major axis along the y-axis
Sol.	$\int x^5 \cdot e^{-4x^3} dx = \frac{1}{48} e^{-4x^3} f(x)$	x)+c		(4) A hyperbola with transverse axis along the x-axis
	Put $x^3 = t$		Ans.	(2)
	$3x^2 dx = dt$		Sol.	$(x^2 - y^2) dx + 2xy dy = 0$
	$\int x^3 \cdot e^{-4x^3} \cdot x^2 dx$			$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{y^2 - x^2}{2xy}$
	$\frac{1}{3}\int t \cdot e^{-4t} dt$			Put $y = vx \implies \frac{dy}{dx} = v + x \frac{dv}{dx}$
	$\frac{1}{3}\left[t\cdot\frac{e^{-4t}}{-4}-\int\frac{e^{-4t}}{-4}dt\right]$			Solving we get,
	$-\frac{e^{-4t}}{48}[4t+1]+c$			$\int \frac{2v}{v^2 + 1} dv = \int -\frac{dx}{x}$
				$\ln(v^2 + 1) = -\ln x + C$
	$\frac{-e^{-4x^3}}{48}[4x^3+1]+c$			$(y^2 + x^2) = Cx$ 1 + 1 = C \Rightarrow C = 2
	$\therefore f(x) = -1 - 4x^3$			
	Option (1)			$y^2 + x^2 = 2x$
	(From the given option	as (1) is most suitable)		\therefore Option (2)



		• · · / _ · • · · · · · · · · · · · · · · · · ·	
30.). If the area of an equilateral triangle inscribed in the circle, $x^2 + y^2 + 10x + 12y + c = 0$ is $27\sqrt{3}$ sq. units then c is equal to :		
	(1) 20	(2) 25	
	(3) 13	(4) -25	
Ans.	(2)		
Sol.	$3\left(\frac{1}{2}r^2.\sin 120^\circ\right) = 27\sqrt{3}$	603	
	$\frac{r^2}{2}\frac{\sqrt{3}}{2} = \frac{27\sqrt{3}}{3}$	r C(-5,-6) r 120°	
	$r^2 = \frac{108}{3} = 36$		
	Radius = $\sqrt{25+36-C}$ =	= \sqrt{36}	
	C = 25		
	\therefore Option (2)		

10

. E

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Thursday 10th JANUARY, 2019) TIME : 02 : 30 PM To 05 : 30 PM PHYSICS

1. Two forces P and Q of magnitude 2F and 3F, respectively, are at an angle θ with each other. If the force Q is doubled, then their resultant also gets doubled. Then, the angle is : (1) 30° (2) 60° (3) 90° (4) 120°

Ans. (4)

Sol. $4F^2 + 9F^2 + 12F^2 \cos \theta = R^2$ $4F^2 + 36 F^2 + 24 F^2 \cos \theta = 4R^2$ $4F^2 + 36 F^2 + 24 F^2 \cos \theta$ $= 4(13F^2 + 12F^2\cos\theta) = 52 F^2 + 48F^2\cos\theta$

$$\cos \theta = -\frac{12F^2}{24F^2} = -\frac{1}{2}$$

2. The actual value of resistance R, shown in the figure is 30Ω . This is measured in an experiment as shown using the standard

formula $R = \frac{V}{I}$, where V and I are the readings

of the voltmeter and ammeter, respectively. If the measured value of R is 5% less, then the internal resistance of the voltmeter is :



(1) 350Ω (2) 570Ω (3) 35Ω (4) 600Ω Ans. (2)

Sol. 0.95 R = $\frac{R R_{\upsilon}}{R + R_{\upsilon}}$ 0.95 × 30 = 0.05 R_v R_v = 19 × 30 = 570 Ω

An unknown metal of mass 192 g heated to a temperature of 100°C was immersed into a brass calorimeter of mass 128 g containing 240 g of water a temperature of 8.4°C Calculate the specific heat of the unknown metal if water temperature stabilizes at 21.5°C (Specific heat of brass is 394 J kg⁻¹ K⁻¹)

Ans. (4)
Sol.
$$192 \times S \times (100 - 21.5)$$

= $128 \times 394 \times (21.5 - 8.4)$

= $128 \times 394 \times (21.5 - 8.4)$ + $240 \times 4200 \times (21.5 - 8.4)$ \Rightarrow S = 916 4. A particle starts from the origin at time t = 0 and moves along the positive x-axis. The graph of velocity with respect to time is shown in figure. What is the position of the particle at time t = 5s?



Ans. (2)

$$\frac{1}{2} \times 2 \times 2 + 2 \times 2 + 3 \times 1 = 9 \text{ m}$$

5. The self induced emf of a coil is 25 volts. When the current in it is changed at uniform rate from 10 A to 25 A in 1s, the change in the energy of the inductance is :

Ans. (1)

$$L\frac{di}{dt} = 25$$
$$L \times \frac{15}{1} = 25$$
$$L = \frac{5}{3}H$$

$$\Delta E = \frac{1}{2} \times \frac{5}{3} \times (25^2 - 10^2) = \frac{5}{6} \times 525 = 437.5 \text{ J}$$

6. A current of 2 mA was passed through an unknown resistor which dissipated a power of 4.4 W. Dissipated power when an ideal power supply of 11V is connected across it is :

(1) $11 \times 10^{-5} \text{ W}$	(2) $11 \times 10^{-4} \text{ W}$
(3) 11×10^5 W	(4) 11×10^{-3} W



E

Ans. (1)

$$P = I^{2}R$$

$$4.4 = 4 \times 10^{-6} R$$

$$R = 1.1 \times 10^{6} \Omega$$

$$P' = \frac{11^{2}}{R} = \frac{11^{2}}{1.1} \times 10^{-6} = 11 \times 10^{-5} W$$

7. The diameter and height of a cylinder are measured by a meter scale to be 12.6 ± 0.1 cm and 34.2 ± 0.1 cm, respectively. What will be the value of its volume in appropriate significant figures ?

(1) 4260 ± 80 cm³
(2) 4300 ± 80 cm³
(3) 4264.4 ± 81.0 cm³
(4) 4264 ± 81 cm³

Ans. (1)

$$\frac{\Delta V}{V} = 2\frac{\Delta d}{d} + \frac{\Delta h}{h} = 2\left(\frac{0.1}{12.6}\right) + \frac{0.1}{34.2}$$
$$V = 12.6 \times \frac{\pi}{4} \times 314.2$$

8. At some location on earth the horizontal component of earth's magnetic field is 18×10^{-6} T. At this location, magnetic neeedle of length 0.12 m and pole strength 1.8 Am is suspended from its mid-point using a thread, it makes 45° angle with horizontal in equilibrium. To keep this needle horizontal, the vertical force that should be applied at one of its ends is :

(1) 3.6×10^{-5} N (2) 6.5×10^{-5} N (3) 1.3×10^{-5} N (4) 1.8×10^{-5} N

Ans. (2)



$$F = 2\mu B$$

9. The modulation frequency of an AM radio station is 250 kHz, which is 10% of the carrier wave. If another AM station approaches you for license what broadcast frequency will you allot ?

(1) 2750 kHz	(2) 2000 kHz
(3) 2250 kHz	(4) 2900 kHz

Ans. (2)

$$f_{carrier} = \frac{250}{0.1} = 2500 \text{ KHZ}$$

:. Range of signal = 2250 Hz to 2750 Hz Now check all options : for 2000 KHZ $f_{mod} = 200$ Hz

 \therefore Range = 1800 KHZ to 2200 KHZ

10. A hoop and a solid cylinder of same mass and radius are made of a permanent magnetic material with their magnetic moment parallel to their respective axes. But the magnetic moment of hoop is twice of solid cylinder. They are placed in a uniform magnetic field in such a manner that their magnetic moments make a small angle with the field. If the oscillation periods of hoop and cylinder are T_h and T_c respectively, then :

(1)
$$T_h = 0.5 T_c$$
 (2) $T_h = 2 T_c$

(3)
$$T_h = 1.5 T_c$$
 (4) $T_h = T_c$

Ans. (4)

$$T = 2\pi \sqrt{\frac{I}{\mu B}}$$
$$T_{h} = 2\pi \sqrt{\frac{mR^{2}}{(2\mu)B}}$$
$$T_{C} = 2\pi \sqrt{\frac{1/2mR^{2}}{\mu B}}$$

11. The electric field of a plane polarized electromagnetic wave in free space at time t= 0 is given by an expression

$$\vec{E}(x,y) = 10\hat{j} \cos [(6x + 8z)]$$

The magnetic field \vec{B} (x, z, t) is given by : (c is the velocity of light)

(1) $\frac{1}{c} (6\hat{k} + 8\hat{i}) \cos[(6x - 8z + 10ct)]$ (2) $\frac{1}{c} (6\hat{k} - 8\hat{i}) \cos[(6x + 8z - 10ct)]$ (3) $\frac{1}{c} (6\hat{k} + 8\hat{i}) \cos[(6x + 8z - 10ct)]$ (4) $\frac{1}{c} (6\hat{k} - 8\hat{i}) \cos[(6x + 8z + 10ct)]$

2

collegebatch.com

Ans. (2)

$$\vec{\mathbf{E}} = 10\hat{\mathbf{j}}\cos\left[\left(6\hat{\mathbf{i}} + 8\hat{\mathbf{k}}\right)\cdot\left(x\hat{\mathbf{i}} + z\hat{\mathbf{k}}\right)\right]$$

 $= 10\hat{j}\cos[\vec{K}\cdot\vec{r}]$

 $\vec{K} = 6\hat{i} + 8\hat{k}; \text{ direction of waves travel.}$ i.e. direction of 'c'.



 \therefore Direction of \hat{B} will be along $\hat{-} -4\hat{i} + 3\hat{k}$

$$\hat{C} \times \hat{E} = \frac{-41 + 3K}{5}$$

Mag. of \vec{B} will be along $\hat{C} \times \hat{E} = \frac{-4\hat{i} + 3\hat{k}}{5}$

Mag. of
$$\vec{B} = \frac{E}{C} = \frac{10}{C}$$

 $\therefore \vec{B} = \frac{10}{C} \left(\frac{-4\hat{i} + 3\hat{k}}{5} \right) = \frac{\left(-8\hat{i} + 6\hat{k}\right)}{C}$

12. Condiser the nuclear fission $Ne^{20} \rightarrow 2He^4 + C^{12}$

Given that the binding energy/nucleon of Ne^{20} , He^4 and C^{12} are, respectively, 8.03 MeV, 7.07 MeV and 7.86 MeV, identify the correct statement :

- (1) 8.3 MeV energy will be released
- (2) energy of 12.4 MeV will be supplied
- (3) energy of 11.9 MeV has to be supplied
- (4) energy of 3.6 MeV will be released

Ans. (3)

Ε

 $\begin{array}{rl} \text{Ne}^{20} & \rightarrow & 2\text{He}^4 + \text{C}^{12} \\ \text{8.03} \times 20 & & 2 \times 7.07 \times 4 + 7.86 \times 12 \\ \therefore & \text{E}_{\text{B}} = (\text{BE})_{\text{react}} & - (\text{BE})_{\text{product}} = 9.72 \text{ MeV} \end{array}$

13. Two vectors \vec{A} and \vec{B} have equal magnitudes. The magnitude of $(\vec{A} + \vec{B})$ is 'n' times the magnitude of $(\vec{A} - \vec{B})$. The angle between \vec{A} and \vec{B} is :

(1)
$$\sin^{-1}\left[\frac{n^2-1}{n^2+1}\right]$$
 (2) $\cos^{-1}\left[\frac{n-1}{n+1}\right]$
(3) $\cos^{-1}\left[\frac{n^2-1}{n^2+1}\right]$ (4) $\sin^{-1}\left[\frac{n-1}{n+1}\right]$

Ans. (3)

$$|\vec{A} + \vec{B}| = 2 \alpha \cos\theta / 2 \qquad (1)$$

$$|\vec{A} - \vec{B}| = 2 \alpha \cos\frac{(\pi - \theta)}{2} = 2 \alpha \sin\theta / 2 \qquad (2)$$

$$\Rightarrow n \left(2 \alpha \cos\frac{\theta}{2}\right) = 2\alpha \frac{\sin\theta}{2}$$

$$\Rightarrow \tan\frac{\theta}{2} = n$$

14. A particle executes simple harmonic motion with an amplitude of 5 cm. When the particle is at 4 cm from the mean position, the magnitude of its velocity in SI units is equal to that of its acceleration. Then, its periodic time in seconds is :

(1)
$$\frac{7}{3}\pi$$
 (2) $\frac{3}{8}\pi$
(3) $\frac{4\pi}{3}$ (4) $\frac{8\pi}{3}$

Ans. (4)



15. Consider a Young's double slit experiment as shown in figure. What should be the slit separation d in terms of wavelength λ such that the first minima occurs directly in front of the slit (S_1) ?



Ans. (4)

 $\sqrt{5}d - 2d = \frac{\lambda}{2}$

16. The eye can be regarded as a single refracting surface . The radius of curvature of this surface is equal to that of cornea (7.8 mm). This surface separates two media of refractive indices 1 and 1.34. Calculate the distance from the refracting surface at which a parallel beam of light will come to focus.

(1) 2 cm	(2) 1 cm
(3) 3.1 cm	(4) 4.0 cm

Ans. (3)

R = 7.8 mm

$$\mu = 1$$
 $\mu = 1.34$
 $\frac{1.34}{V} - \frac{1}{\infty} = \frac{1.34 - 1}{7.8}$
∴ V = 30.7 mm

17. Half mole of an ideal monoatomic gas is heated at constant pressure of 1atm from 20 °C to 90°C. Work done by gas is close to : (Gas constant R = 8.31 J /mol.K(1) 73 J (2) 291 J (3) 581 J (4) 146 J

Ans. (2)

WD = P
$$\Delta$$
V = nR Δ T = $\frac{1}{2} \times 8.31 \times 70$

18. A metal plate of area 1×10^{-4} m² is illuminated by a radiation of intensity 16 mW/m². The work function of the metal is 5eV. The energy of the incident photons is 10 eV and only 10% of it produces photo electrons. The number of emitted photo electrons per second and their maximum energy, respectively, will be : $[1 \text{ eV} = 1.6 \times 10^{-19}\text{J}]$

> (1) 10^{10} and 5 eV (2) 10^{14} and 10 eV(4) 1011 and 5 eV

(3)
$$10^{12}$$
 and 5 eV (4) 10^{11} and

Ans. (4)

$$I = \frac{nE}{At}$$

16×10⁻³ = $\left(\frac{n}{t}\right)_{Photon} \frac{10 \times 1.6 \times 10^{-19}}{10^{-4}} = 10^{12}$

19. Charges -q and +q located at A and B, respectively, constitute an electric dipole. Distance AB = 2a, O is the mid point of the dipole and OP is perpendicular to AB. A charge Q is placed at P where OP = y and y >> 2a. The charge Q experiences and electrostatic force F. If Q is now moved along the equatorial line

to P' such that OP'=
$$\left(\frac{y}{3}\right)$$
, the force on Q will be
close to : $\left(\frac{y}{3} >> 2a\right)$
Q P'
Q P'
A Q P'
A Q P'
A Q P'
Q O B
+q
(1) $\frac{F}{3}$ (2) 3F (3) 9F (4) 27F



Ans. (4)

Sol. Electric field of equitorial plane of dipole

$$= -\frac{K\vec{P}}{r^3}$$

$$\therefore \text{ At P, F} = -\frac{K\vec{P}}{r^3}Q.$$

$$K\vec{P}Q.$$

At P¹, F¹ =
$$-\frac{KPQ}{(r/3)^3} = 27 F$$
.

- 20. Two stars of masses 3×10^{31} kg each, and at distance 2×10^{11} m rotate in a plane about their common centre of mass O. A meteorite passes through O moving perpendicular to the star's rotation plane. In order to escape from the gravitational field of this double star, the minimum speed that meteorite should have at O is : (Take Gravitational constant $G = 6.67 \times 10^{-11}$ Nm² kg⁻²)
 - (1) 1.4×10^5 m/s (2) 24×10^4 m/s (3) 3.8×10^4 m/s (4) 2.8×10^5 m/s

Ans. (4)

By energy convervation between 0 & ∞ .

$$-\frac{GMm}{r} + \frac{-GMm}{r} + \frac{1}{2}mV^2 = 0 + 0$$

[M is mass of star m is mass of meteroite)

$$\Rightarrow v = \sqrt{\frac{4GM}{r}} = 2.8 \times 10^5 \,\text{m/s}$$

21. A closed organ pipe has a fundamental frequency of 1.5 kHz. The number of overtones that can be distinctly heard by a person with this organ pipe will be : (Assume that the highest frequency a person can hear is 20,000 Hz)

(1) 7 (2) 5 (3) 6 (4) 4

Ans. (1)

Ε

Sol. For closed organ pipe, resonate frequency is odd multiple of fundamental frequency.

: (2n + 1) $f_0 \le 20,000$

(
$$f_o$$
 is fundamental frequency = 1.5 KHz)
∴ n = 6

 \therefore Total number of overtone that can be heared is 7. (0 to 6).

22. A rigid massless rod of length 3*l* has two masses attached at each end as shown in the figure. The rod is pivoted at point P on the horizontal axis (see figure). When released from initial horizontal position, its instantaneous angular acceleration will be :



Ans. (3)

$$5M$$
 P Q $2M$

Applying torque equation about point P. $2M_0 (2l) - 5 M_0 gl = I\alpha$

$$I = 2M_0 (2l)^2 + 5M_0 l^2 = 13 M_0 l^2 d$$

$$\therefore \quad \alpha = -\frac{M_0 g \ell}{13M_0 \ell^2} \implies \alpha = -\frac{g}{13\ell}$$

$$\therefore \quad \alpha = \frac{g}{13\ell} \text{ anticlockwise}$$

$$\therefore \alpha = \frac{s}{13\ell}$$
 anticlockwise

23. For the circuit shown below, the current through the Zener diode is :



Assuming zener diode doesnot undergo breakdown, current in circuit = $\frac{120}{15000} = 8 \text{ mA}$ \therefore Voltage drop across diode = 80 V > 50 V. The diode undergo breakdown.



5

collegebatch.com

24. Four equal point charges Q each are placed in the xy plane at (0, 2), (4, 2), (4, -2) and (0, -2). The work required to put a fifth charge Q at the origin of the coordinate system will be :

(1)
$$\frac{Q^2}{2\sqrt{2}\pi\epsilon_0}$$
 (2) $\frac{Q^2}{4\pi\epsilon_0}\left(1+\frac{1}{\sqrt{5}}\right)$
(3) $\frac{Q^2}{4\pi\epsilon_0}\left(1+\frac{1}{\sqrt{3}}\right)$ (4) $\frac{Q^2}{4\pi\epsilon_0}$

Ans. (2)

 $(0,2) \bullet Q$





Potential at origin = $\frac{KQ}{2} + \frac{KQ}{2} + \frac{KQ}{\sqrt{20}} + \frac{KQ}{\sqrt{20}}$ (Potential at $\infty = 0$)

 $= KQ\left(1 + \frac{1}{\sqrt{5}}\right)$

 \therefore Work required to put a fifth charge Q at origin

is equal to $\frac{Q^2}{4\pi\epsilon_0}\left(1+\frac{1}{\sqrt{5}}\right)$

- 25. A cylindrical plastic bottle of negligible mass is filled with 310 ml of water and left floating in a pond with still water. If pressed downward slightly and released, it starts performing simple harmonic motion at angular frequency ω . If the radius of the bottle is 2.5 cm then ω close to : (density of water = 10³ kg / m³) (1) 5.00 rad s⁻¹ (2) 1.25 rad s⁻¹
- (3) 3.75 rad s⁻¹





(4) 2.50 rad s⁻¹

$$B = ma$$

$$a = \left(\frac{\delta Ag}{m}\right)^{x}$$

$$w^{2} = \frac{\delta Ag}{m}$$

$$w = \sqrt{\frac{10^{3} \times \pi (2.5)^{2} \times 10^{-4} \times 10}{310 \times 10^{-6} \times 10^{3}}}$$

$$= \sqrt{63.30} = 7.95$$

26. A parallel plate capacitor having capacitance 12 pF is charged by a battery to a potential difference of 10 V between its plates. The charging battery is now disconnected and a porcelain slab of dielectric constant 6.5 is slipped between the plates the work done by the capacitor on the slab is :

Ans. (3)

Intial energy of capacitor

$$U_{i} = \frac{1}{2} \frac{v^{2}}{c}$$
$$= \frac{1}{2} \times \frac{120 \times 120}{12} = 600 \text{ J}$$

Since battery is disconnected so charge remain same.

Final energy of capacitor

$$U_{f} = \frac{1}{2} \frac{v^{2}}{c}$$
$$= \frac{1}{2} \times \frac{120 \times 120}{12 \times 6.5} = 92$$
$$W + U_{f} = U_{i}$$
$$W = 508 \text{ J}$$

27. Two kg of a monoatomic gas is at a pressure of 4×10^4 N/m². The density of the gas is 8 kg /m³. What is the order of energy of the gas due to its thermal motion ?

(1)
$$10^3 \text{ J}$$
 (2) 10^5 J

(3)
$$10^6 \text{ J}$$
 (4) 10^4 J

Ans. (4)

Thermal energy of N molecule

$$= N\left(\frac{3}{2}kT\right)$$



$$= \frac{N}{N_A} \frac{3}{2} RT$$

$$= \frac{3}{2} (nRT)$$

$$= \frac{3}{2} PV$$

$$= \frac{3}{2} P\left(\frac{m}{8}\right)$$

$$= \frac{3}{2} \times 4 \times 10^4 \times \frac{2}{8}$$

$$= 1.5 \times 10^4$$
order will 10⁴
A particle which is experiencing a

by $\vec{F} = 3\vec{i} - 12\vec{j}$, undergoes a displacement of $\vec{d} = 4\vec{i}$. If the particle had a kinetic energy of 3 J at the beginning of the displacement, what is its kinetic energy at the end of the displacement ? (1) 15 J (2) 10 J (3) 12 J (4) 9 J

force, given

Ans. (1)

28.

Work done = $\vec{F} \cdot \vec{d}$ = 12J work energy theorem $w_{net} = \Delta K.E.$ $12 = K_f - 3$

$$K_f = 15J$$

The Wheater

29. The Wheatstone bridge shown in Fig. here, gets balanced when the carbon resistor used as R_1 has the colour code (Orange, Red, Brown). The resistors R_2 and R_4 are 80 Ω and 40 Ω , respectively.

Assuming that the colour code for the carbon resistors gives their accurate values, the colour code for the carbon resistor, used as R_3 , would be :



- (1) Red, Green, Brown
- (2) Brown, Blue, Brown
- (3) Grey, Black, Brown
- (4) Brown, Blue, Black

Ans. (2)

$$R_1 = 32 \times 10 = 320$$

for wheat stone bridge
 $\Rightarrow \frac{R_1}{R_3} = \frac{R_2}{R_4}$
 $\frac{320}{R_3} = \frac{80}{40}$
 $R_3 = 160$
Brown Blue Brown

30. Two identical spherical balls of mass M and radius R each are stuck on two ends of a rod of length 2R and mass M (see figure). The moment of inertia of the system about the axis passing perpendicularly through the centre of the rod is :



Ans. (3)

For Ball

using parallel axis theorem.

$$I_{\text{ball}} = \frac{2}{5} \text{MR}^2 + \text{M}(2\text{R})^2$$
$$= \frac{22}{5} \text{MR}^2$$

2 Balls so $\frac{44}{5}$ MR²

Irod = for rod
$$\frac{M(2R)^2}{R} = \frac{MR^2}{3}$$

 $I_{system} = I_{Ball} + I_{rod}$
 $= \frac{44}{5}MR^2 + \frac{MR^2}{3}$

$$=\frac{137}{15} \mathrm{MR}^2$$

collegebatch.com

E

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Thrusday 10th JANUARY, 2019) TIME : 02 : 30 PM To 05 : 30 PM CHEMISTRY

1. An ideal gas undergoes isothermal compression from 5 m³ against a constant external pressure of 4 Nm⁻². Heat released in this process is used to increase the temperature of 1 mole of Al. If molar heat capacity of Al is 24 J mol⁻¹ K⁻¹, the temperature of Al increases by :

(1)
$$\frac{3}{2}$$
K (2) $\frac{2}{3}$ K (3) 1 K (4) 2 K

Ans. (2)

Sol. Work done on isothermal irreversible for ideal gas

$$= -P_{ext} (V_2 - V_1)$$

= -4 N/m² (1m³ - 5m³)
= 16 Nm
Isothermal process for ideal gas
 $\Delta U = 0$
q = -w
= -16 Nm
= - 16 J
Heat used to increase temperature of Al
q = n C_m ΔT

$$16 J = 1 \times 24 \frac{J}{\text{mol. K}} \times \Delta T$$

$$\Delta T = \frac{2}{3}K$$

2. The 71^{st} electron of an element X with an atomic number of 71 enters into the orbital : (1) 4f (2) 6p (3) 6s (4) 5d

Ans. (1)

- 3. The number of 2-centre-2-electron and 3centre-2-electron bonds in B_2H_6 , respectively, are :
 - (1) 2 and 4 (2) 2 and 1
 - (3) 2 and 2 (4) 4 and 2

Ans. (4)

4. The amount of sugar (C₁₂H₂₂O₁₁) required to prepare 2 L of its 0.1 M aqueous solution is : (1) 68.4 g (2) 17.1 g (3) 34.2 g (4)136.8 g
Ans (1)

Ans. (1)

Ε

Sol. Molarity =
$$\frac{(n)_{solute}}{V_{solution} (in lit)}$$

 $0.1 = \frac{WL/342}{2}$ wt (C₁₂H₂₂O₁₁) = 68.4 gram 5. Among the following reactions of hydrogen with halogens, the one that requires a catalyst is :

(1)
$$H_2 + I_2 \rightarrow 2HI$$
 (2) $H_2 + F_2 \rightarrow 2HF$
(3) $H_2 + Cl_2 \rightarrow 2HCI$ (4) $H_2 + Br_2 \rightarrow 2HBr$

Ans. (1)

- 6. Sodium metal on dissolution in liquid ammonia gives a deep blue solution due to the formation of:
 - (1) sodium ion-ammonia complex
 - (2) sodamide
 - (3) sodium-ammonia complex
 - (4) ammoniated electrons

Ans. (4)

7. What will be the major product in the following mononitation reaction ?











Ans. (3)



In the cell $Pt(s)|H_2(g, 1bar|HCl(aq)|Ag(s)|Pt(s)$ 8. the cell potential is 0.92 when a 10⁻⁶ molal HCl solution is used. THe standard electrode potential of (AgCl/Ag,Cl-) electrode is : $\left\{ \text{given}, \frac{2.303\text{RT}}{\text{F}} = 0.06\text{Vat}298\text{K} \right\}$ (1) 0.20 V (2) 0.76 V (3) 0.40 V (4) 0.94 V Ans. (1) $Pt(s)|H_2(g, 1bar)|HCl(aq)|AgCl(s)|Ag(s)|Pt(s)$ Sol. 10^{-6} m $H_2 \longrightarrow 2H^+ + 2e \times 1$ Anode: **Cathode :** $e^{-} + AgCl(s) \longrightarrow Ag(s) + Cl^{-}(aq)$ $\times 2$ $H_2(g)l + AgCl(s) \longrightarrow 2H^+ +$ $2Ag(s) + 2Cl^{-}(aq)$ $E_{cell} = E_{cell}^{0} - \frac{0.06}{2} \log_{10} \left((H^{+})^{2} \cdot (Cl^{-})^{2} \right)$ $.925 = \left(E_{H_2/H^+}^0 + E_{AgCl/Ag,Cl^-}^0 \right) - \frac{0.06}{2} \log_{10}$ $((10^{-6})^2 (10^{-6})^2)$ $.92 = 0 + E_{AgCl/Ag,Cl^{-}}^{0} - 0.03 \log_{10}(10^{-6})^{4}$ E^{0}_{AgCl} / Ag, $Cl^{-} = .92 + .03 \times -24 = 0.2 V$ 9. The major product of the following recation is: CH₃N NaBH₄ OH $(1) CH_3N$ OH (2) CH₃N. OH $(3) CH_3N$

OH

- 10. The pair that contains two P-H bonds in each of the oxoacids is :
 (1) H₃PO₂ nad H₄P₂O₅
 (2) H₄P₂O₅ and H₄P₂O₆
 (3) H₃PO₃ and H₃PO₂
 (4) H₄P₂O₅ nad H₃PO₃
- Ans. (1)
- 11. The major product of the following reaction is:



Ans. (4)

12. The difference in the number of unpaired electrons of a metal ion in its high-spin and low-spin octahedral complexes is two. The metal ion is :

(4) Ni^{2+}

(1)
$$Fe^{2+}$$
 (2) Co^{2+} (3) Mn^{2+}

Ans. (2)

13. A compound of formula A_2B_3 has the hcp lattice. Which atom forms the hcp lattice and what fraction of tetrahedral voids is occupied by the other atoms :

2



(4) CH₃N

Ans. (3)

E

Sol. A_2B_3 has HCP lattice

If A form HCP, then $\frac{3}{4}^{\text{th}}$ of THV must occupied by B to form A_2B_3

If B form HCP, then $\frac{1}{3}^{\text{th}}$ of THV must occupied by A to form A_2B_3

14. The reaction that is NOT involved in the ozone layer depletion mechanism is the stratosphere is:

(1) HOCl(g)
$$\xrightarrow{h\upsilon} OH(g) + Cl(g)$$

- (2) $CF_2Cl_2(g) \xrightarrow{uv} Cl(g) + CF_2Cl(g)$
- (3) $CH_4 + 2O_3 \rightarrow 3CH_2 = O + 3H_2OP$
- (4) $\dot{ClO(g)} + O(g) \rightarrow \dot{Cl}(g) + O_2(g)$

Ans. (3)

- 15. The process with negative entropy change is :
 - (1) Dissolution of iodine in water
 - (2) Synthesis of ammonia from N_2 and H_2
 - (3) Dissolution of $CaSO_4(s)$ to CaO(s) and $SO_3(g)$
 - (4) Subimation of dry ice

Ans. (2)

Ε

Sol. $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$; $\Delta n_g < 0$

16. The major product of the following reaction is:



Ans. (3)

- 17. A reaction of cobalt(III) chloride and ethylenediamine in a 1 : 2 mole ratio generates two isomeric products A (violet coloured) B (green coloured). A can show optial actively, B is optically inactive. What type of isomers does A and B represent ?
 - (1) Geometrical isomers
 - (2) Ionisation isomers]
 - (3) Coordination isomers
 - (4) Linkage isomers

Ans. (1)



18. The major product obtained in the following reaction is :



Ans. (4)

- **19.** Which of the following tests cannot be used for identifying amino acids ?
 - (1) Biuret test (2) Xanthoproteic test
 - (3) Barfoed test (4) Ninhydrin test
- Ans. (3)
- **20.** What is the IUPAC name of the following compound ?



- (1) 3-Bromo-1, 2-dimethylbut-1-ene]
- (2) 4-Bromo-3-methylpent-2-ene
- (3) 2-Bromo-3-methylpent-3-ene
- (4) 3-Bromo-3-methyl-1, 2-dimethylprop-1-ene
- Ans. (2)
- **21.** Which is the most suitable reagent for the following transformation ?

$$\begin{array}{c} & & & & & \\ & H \\ CH_3-CH=CH-CH_2-CH-CH_3 \longrightarrow \\ & & CH_3-CH=CH-CH_2CO_2H \\ (1) \text{ alkaline KMnO}_4 \quad (2) \text{ I}_2/\text{NaOH} \\ (3) \text{ Tollen's reagent} \quad (4) \text{ CrO}_2/\text{CS}_2 \\ \textbf{Ans. (2)} \end{array}$$

22.	The correct match between	n item T' and item 'II' is :	
	Item 'I'	Item 'II'	
	(compound)	(reagent)	
	(A) Lysine	(P) 1-naphthol	
	(B) Furfural	(Q) ninhydrin	
	(C) Benzyl alcohol	(R) KMnO ₄	
	(D) Styrene	(S) Ceric ammonium	
		nitrate	
	(1) (A) \rightarrow (Q), (B) \rightarrow (P),	$(C)\rightarrow(S), (D)\rightarrow(R)$	
	(2) (A) \rightarrow (Q), (B) \rightarrow (R),	$(C)\rightarrow(S), (D)\rightarrow(P)$	
	$(3) (A) \rightarrow (Q), (B) \rightarrow (P),$	$(C) \rightarrow (R), (D) \rightarrow (S)$	
	$(4) (A) \rightarrow (R), (B) \rightarrow (P),$	$(C) \rightarrow (Q), (D) \rightarrow (S)$	
Ans.		·.1	
23.	In the reaction of oxalate w	rith permaganate in acidic	
	medium, the number of	electrons involved in	
	producing one molecule (1) 10 (2) 2	(2) 1 (4) 5	
Ang	(1) 10 (2) 2	(3) 1 (4) 3	
Ans.	(3)		
	$2 \stackrel{+7}{M} nO_4 + 5C_2O_4^{2-} + 16$	$H^+ \longrightarrow 2 \stackrel{+2}{M} n^{2+}$	
Sol.	7 2 7	$+10CO_{2} + 8H_{2}O_{3}$	
	$10 e^{-}$ trans for 10 mol	$\frac{1}{1000} + \frac{1}{2} + \frac{1}{100} + \frac{1}{2} + \frac{1}{100} + \frac{1}{2} + \frac{1}{100} + \frac{1}{100}$	
	molecule of CO transf	fer of e^- is '1'	
24.	5.1g NH ₄ SH is introduc	ed in 3.0 L evacuated	
	flask at 327°C. 30%	of the solid NH ₄ SH	
	decomposed to NH ₂ and	H_2S as gases. The K.	
	of the reaction at 327° C is (R = 0.082 L atm		
	$mol^{-1}K^{-1}$, Molar mass of S = 32 g mol ^{/01} , molar		
	mass of N = 14g mol ^{-1})	
	(1) $1 \times 10^{-4} \text{ atm}^2$	(2) $4.9 \times 10^{-3} \text{ atm}^2$	
	(3) 0.242 atm^2	(4) $0.242 \times 10^{-4} \text{ atm}^2$	
Ans.	(3)		
	NH SH(α) \longrightarrow NH (α) + H S(α)		
	$NH_4SH(s) \Longrightarrow NH_3(g)$	$(H_2S(g))$	
a .	$n = \frac{5.1}{1} = 1$ mole 0	0	
Sol.	51	Ū	
	$.1(-1-\alpha)$ $.1\alpha$.1α	
	$\alpha = 30\% = .3$		
	so number of moles at	equilibrium	
	$.1(13)$ $.1 \times .22$	$3 .1 \times .3$	
	= .0/ $= .03$		
	Now use $\mathbf{P}\mathbf{v} = \mathbf{IIKI}$ at $\mathbf{P} \times 3 \text{ lit} = (03 \pm 0)^2$	$(4) \times 082 \times 600$	
	$P_{total} = .984 \text{ atm}$	5) ~ .002 ~ 000	
	At equilibrium		
	D		
	$P_{\rm NH_3} = P_{\rm H_2S} = \frac{r_{\rm total}}{2} = .49$	2	
	So $k_n = P_{NH_2} \cdot P_{H_2S} = (.4)$	92) (.492)	
	p525		
	$k_{p} = .242 \text{ atm}^{2}$		



- 25. The electrolytes usually used in the electroplating of gold and silver, respectively, are :
 (1) [Au(OH)₄]⁻ and [Ag(OH)₂]⁻
 (2) [Au(CN)₂]⁻ and [Ag CI₂]⁻
 (3) [Au(NH₃)₂]⁺ and [Ag(CN)₂]⁻
 - (4) $[Au(CN)_2]^-$ and $[Ag(CN)_2]^-$

Ans. (4)

26. Elevation in the boiling point for 1 molal solution of glucose is 2 K. The depression in the freezing point of 2 molal solutions of glucose in the same solvent is 2 K. The relation between K_b and K_f is:

(1)
$$K_b = 0.5 K_f$$
 (2) $K_b = 2 K_f$
(3) $K_b = 1.5 K_f$ (4) $K_b = K_f$

Ans. (2)

Sol. Ans.(2)

 $\frac{\Delta T_{b}}{\Delta T_{f}} = \frac{i.m \times k_{b}}{i \times m \times k_{f}}$

$$\frac{2}{2} = \frac{1 \times 1 \times k_{b}}{1 \times 2 \times k_{c}}$$

 $k_{\rm b} = 2k_{\rm f}$

27. An aromatic compound 'A' having molecular formula $C_7H_6O_2$ on treating with aqueous ammonia and heating forms compound 'B'. The compound 'B' on reaction with molecular bromine and potassium hydroxide provides compound 'C' having molecular formula C_6H_7N . The structure of 'A' is :







Ans. (3)

28. The ground state energy of hydrogen atom is -13.6 eV. The energy of second excited state He⁺ ion in eV is : (1) -6.04 (2) -27.2 (3) -54.4 (4) -3.4
Ans. (1)

Sol.
$$(E)_{n^{\text{th}}} = (E_{\text{GND}})_{\text{H}} \cdot \frac{Z^2}{n^2}$$

$$E_{3^{rd}}(He^+) = (-13.6 \text{ eV}) \cdot \frac{2^2}{3^2} = -6.04 \text{ eV}$$

29. For an elementary chemical reaction,

A₂
$$\xrightarrow{k_1} 2A$$
, the expression for $\frac{d[A]}{dt}$ is :
(1) $2k_1[A_2]-k_{-1}[A]^2$ (2) $k_1[A_2]-k_{-1}[A]^2$
(3) $2k_1[A_2]-2k_{-1}[A]^2$ (4) $k_1[A_2]+k_{-1}[A]^2$
Ans. (3)
Sol. Ans.(3)
 $A_2 \xrightarrow{K_1} 2A$

$$\frac{d[A]}{dt} = 2k_1[A_2] - 2k_{-1}[A]^2$$

K 1

30. Haemoglobin and gold sol are examples of : (1) negatively charged sols

- (2) positively charged sols]
- (3) negatively and positively charged sols, respectively
- (4) positively and negatively charged sols, respectively

Ans. (4)

Sol. Ans.(4)

Haemoglobin \longrightarrow positive sol Ag - sol \longrightarrow negative sol

Ε



TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Friday 11th JANUARY, 2019) TIME : 9 : 30 AM To 12 : 30 PM M ATHEM ATICS

	$\begin{pmatrix} 0 & 2a & r \end{pmatrix}$	3.	The outcome of each of 30 items was observed;
1.	Let $A = \begin{bmatrix} 0 & -q & 1 \\ p & q & -r \\ p & -q & r \end{bmatrix}$. It $AA^{T} = I_{3}$, then $ p $		10 items gave an outcome $\frac{1}{2}$ – d each, 10 items
	(p q 1) is:		gave outcome $\frac{1}{2}$ each and the remaining
	(1) $\frac{1}{\sqrt{2}}$		10 items gave outcome $\frac{1}{2}$ + d each. If the
	(2) $\frac{1}{\sqrt{5}}$		variance of this outcome data is $\frac{4}{3}$ then $ d $
	(3) $\frac{1}{\sqrt{6}}$		equals :-
	(4) $\frac{1}{\sqrt{2}}$		(1) 2 (2) $\frac{\sqrt{5}}{2}$ (3) $\frac{2}{3}$ (4) $\sqrt{2}$
Ans. Sol.	(1) A is orthogonal matrix	Ans. Sol.	(4) Variance is independent of origin. So we shift
	$\Rightarrow 0^2 + p^2 + p^2 = 1 \Rightarrow \mathbf{p} = \frac{1}{\sqrt{2}}$		the given data by $\frac{1}{2}$.
2.	The area (in sq. units) of the region bounded by the curve $x^2 = 4y$ and the straight line		so, $\frac{10d^2 + 10 \times 0^2 + 10d^2}{30} - (0)^2 = \frac{4}{3}$
	x = 4y - 2 :-		$\Rightarrow d^2 = 2 \Rightarrow d - \sqrt{2}$
	(1) $\frac{5}{4}$	4.	The sum of an infinite geometric series with positive terms is 3 and the sum of the cubes of
	(2) $\frac{9}{8}$		its terms is $\frac{27}{19}$. Then the common ratio of this
	(3) $\frac{3}{4}$		series is :
	$(1) \frac{7}{2}$		(1) $\frac{4}{9}$ (2) $\frac{2}{9}$
Ans.	(4) 8 (2)		(3) $\frac{2}{3}$ (4) $\frac{1}{3}$
		Ans.	(3)
Sol.		Sol.	$\frac{a}{1-r} = 3 \qquad \dots (1)$
	$x = 4y - 2 & x^2 = 4y$ $\Rightarrow x^2 = x + 2 \Rightarrow x^2 - x - 2 = 0$		$\frac{a^3}{1-r^3} = \frac{27}{19} \Rightarrow \frac{27(1-r)^3}{1-r^3} = \frac{27}{19}$
	x = 2, -1		$\Rightarrow 6r^2 - 13r + 6 = 0$
	So, $\int_{-1}^{2} \left(\frac{x+2}{4} - \frac{x^2}{4} \right) dx = \frac{9}{8}$		\Rightarrow r = $\frac{2}{3}$ as $ r < 1$
			1



Е

5.	Let $\vec{a} = \hat{i} + 2\hat{j} + 4\hat{k}$, $\vec{b} = \hat{i} + \lambda\hat{j} + 4\hat{k}$ and		
	$\vec{c} = 2\hat{i} + 4\hat{j} + (\lambda^2 - 1)\hat{k}$ be coplanar vectors.		
	Then the non-zero vector $\vec{a} \times \vec{c}$ is :		
	(1) $-14\hat{i} - 5\hat{j}$ (2) $-10\hat{i} - 5\hat{j}$		
	$(3) -10\hat{i} + 5\hat{j} \qquad (4) -14\hat{i} + 5\hat{j}$		
Ans.	(3)		
Sol.	$\begin{bmatrix} \vec{a} & \vec{b} & \vec{c} \end{bmatrix} = 0$		
	$\Rightarrow \begin{vmatrix} 1 & 2 & 4 \\ 1 & \lambda & 4 \\ 2 & 4 & \lambda^2 - 1 \end{vmatrix} = 0$		
	$\Rightarrow \lambda^{3} - 2\lambda^{2} - 9\lambda + 18 = 0$ $\Rightarrow \lambda^{2}(\lambda - 2) - 9(\lambda - 2) = 0$ $\Rightarrow (\lambda - 3)(\lambda + 3)(\lambda - 2) = 0$ $\Rightarrow \lambda = 2, 3, -3$ So, $\lambda = 2$ (as \vec{a} is parallel to \vec{c} for $\lambda = \pm 3$)		
	Hence $\vec{a} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 4 \\ 2 & 4 & 3 \end{vmatrix}$		
	$=-10\hat{i}+5\hat{j}$		
6.	Let $\left(-2 - \frac{1}{3}i\right)^3 = \frac{x + iy}{27}(i = \sqrt{-1})$, where x		
	and y are real numbers, then $y - x$ equals :		
	$\begin{array}{cccccccccccccccccccccccccccccccccccc$		
Ans.	(4)		
Sol.	$\left(-2 - \frac{i}{3}\right)^3 = -\frac{(6+i)^3}{27}$		
	$=\frac{-198-107i}{27}=\frac{x+iy}{27}$		
	Hence, $y - x = 198 - 107 = 91$		

7. Let $f(x) = \begin{cases} -1, -2 \le x < 0 \\ x^2 - 1, \ 0 \le x \le 2 \end{cases}$ and g(x) = |f(x)| + f(|x|). Then, in the interval (-2, 2), g is :-(1) differentiable at all points (2) not differentiable at two points (3) not continuous (4) not differentiable at one point Ans. (4) Sol. $|f(\mathbf{x})| = \begin{cases} 1 & , -2 \le \mathbf{x} < 0 \\ 1 - \mathbf{x}^2 & , 0 \le \mathbf{x} < 1 \\ \mathbf{x}^2 - 1 & , 1 \le \mathbf{x} \le 2 \end{cases}$ and $f(|\mathbf{x}|) = \mathbf{x}^2 - 1$, $\mathbf{x} \in [-2, 2]$ Hence $g(x) = \begin{cases} x^2 & , x \in [-2,0) \\ 0 & , x \in [0,1) \\ 2(x^2 - 1) & , x \in [1,2] \end{cases}$ It is not differentiable at x = 1Let $f: R \to R$ be defined by $f(x) = \frac{x}{1 + x^2}$, $x \in R$. Then the range of f is : (1) $(-1, 1) - \{0\}$ (2) $\left[-\frac{1}{2}, \frac{1}{2}\right]$ (3) $R = \left[-\frac{1}{2}, \frac{1}{2} \right]$ (4) R = [-1, 1]Ans. (2) **Sol.** f(0) = 0 & f(x) is odd. Further, if x > 0 then $f(\mathbf{x}) = \frac{1}{\mathbf{x} + \frac{1}{x}} \in \left(0, \frac{1}{2}\right]$

Hence, $f(x) \in \left| -\frac{1}{2}, \frac{1}{2} \right|$

8.

2



- 9. The sum of the real values of x for which the middle term in the binomial expansion of $\left(\frac{x^3}{3} + \frac{3}{x}\right)^8$ equals 5670 is : (1) 6 (2) 8 (3) 0 (4) 4 Ans. (3) Sol. $T_5 = {}^8C_4 \frac{x^{12}}{81} \times \frac{81}{x^4} = 5670$ $\Rightarrow 70x^8 = 5670$ $\Rightarrow x = \pm \sqrt{3}$ 10. The value of r for which ${}^{20}C_r {}^{20}C_0 + {}^{20}C_{r-1} {}^{20}C_1 + {}^{20}C_{r-2} {}^{20}C_2 + \dots {}^{20}C_0 {}^{20}C_r$
 - is maximum, is (1) 20 (2) 15 (3) 11 (4) 10
- Ans. (1)
- Sol. Given sum = coefficient of x^r in the expansion of $(1 + x)^{20}(1 + x)^{20}$, which is equal to ${}^{40}C_r$ It is maximum when r = 20
- 11. Let a_1, a_2, \dots, a_{10} be a G.P. If $\frac{a_3}{a_1} = 25$, then
 - $\frac{a_9}{a_5} \text{ equals :}$ (1) 2(5²)
 (2) 4(5²)
 (3) 5⁴
 (4) 5³
- Ans. (3)
- **Sol.** a_1, a_2, \dots, a_{10} are in G.P., Let the common ratio be r

$$\frac{a_3}{a_1} = 25 \implies \frac{a_1 r^2}{a_1} = 25 \implies r^2 = 25$$
$$\frac{a_9}{a_1} = \frac{a_1 r^8}{a_1} = r^4 = 5^4$$

12. If
$$\int \frac{\sqrt{1-x^2}}{x^4} dx = A(x) (\sqrt{1-x^2})^m + C$$
, for

a suitable chosen integer m and a function A(x), where C is a constant of integration then $(A(x))^m$ equals :

(1)
$$\frac{-1}{3x^3}$$
 (2) $\frac{-1}{27x^9}$

(3)
$$\frac{1}{9x^4}$$
 (4) $\frac{1}{27x^6}$

Ans. (2)

Sol.
$$\int \frac{\sqrt{1-x^2}}{x^4} dx = A(x) \left(\sqrt{1-x^2}\right)^m + C$$
$$\int \frac{|x|}{\sqrt{\frac{1}{x^2}-1}} dx,$$
$$Put \frac{1}{x^2} - 1 = t \Rightarrow \frac{dt}{dx} = \frac{-2}{x^3}$$
$$Case-1 \ x \ge 0$$
$$-\frac{1}{2} \int \sqrt{t} \ dt \Rightarrow -\frac{t^{3/2}}{3} + C$$
$$\Rightarrow -\frac{1}{3} \left(\frac{1}{x^2} - 1\right)^{3/2}$$
$$\Rightarrow \frac{\left(\sqrt{1-x^2}\right)^3}{-3x^2} + C$$
$$A(x) = -\frac{1}{3x^3} \ and \ m = 3$$
$$(A(x))^m = \left(-\frac{1}{3x^3}\right)^3 = -\frac{1}{27x^9}$$
$$Case-II \ x \le 0$$
$$We \ get \ \frac{\left(\sqrt{1-x^2}\right)^3}{-3x^3} + C$$
$$A(x) = -\frac{1}{-3x^3}, \ m = 3$$
$$(A(x))^m = \frac{-1}{27x^9}$$

Е



 $a_5 a_1 r^4$

13. In a triangle, the sum of lengths of two sides is x and the product of the lengths of the same two sides is y. If $x^2 - c^2 = y$, where c is the length of the third side of the triangle, then the circumradius of the triangle is :

(1)
$$\frac{y}{\sqrt{3}}$$
 (2) $\frac{c}{\sqrt{3}}$ (3) $\frac{c}{3}$ (4) $\frac{3}{2}y$

Ans. (2)

Sol. Given a + b = x and ab = yIf $x^2 - c^2 = y \implies (a + b)^2 - c^2 = ab$ $\Rightarrow a^2 + b^2 - c^2 = -ab$ $\Rightarrow \frac{a^2 + b^2 - c^2}{2ab} = -\frac{1}{2}$ $\Rightarrow \cos C = -\frac{1}{2}$ $\Rightarrow \angle C = \frac{2\pi}{3}$ $R = \frac{c}{2 \sin C} = \frac{c}{\sqrt{3}}$

The value of the integral $\int \frac{\sin^2 x}{\left\lceil \frac{x}{\pi} \right\rceil + \frac{1}{2}} dx$ 14.

> (where [x] denotes the greatest integer less than 20 Cr or equal to x) is :

(1) 4(2) $4 - \sin 4$ (3) sin 4 (4) 0

Ans. (4)

Sol.
$$I = \int_{-2}^{2} \frac{\sin^{2} x}{\left[\frac{x}{\pi}\right] + \frac{1}{2}} dx$$
$$I = \int_{0}^{2} \left(\frac{\sin^{2} x}{\left[\frac{x}{\pi}\right] + \frac{1}{2}} + \frac{\sin^{2}(-x)}{\left[-\frac{x}{\pi}\right] + \frac{1}{2}} \right) dx$$
$$\left(\left[\frac{x}{\pi}\right] + \left[-\frac{x}{\pi}\right] = -1 \text{ as } x \neq n\pi \right)$$
$$I = \int_{0}^{2} \left(\frac{\sin^{2} x}{\left[\frac{x}{\pi}\right] + \frac{1}{2}} + \frac{\sin^{2} x}{-1 - \left[\frac{x}{\pi}\right] + \frac{1}{2}} \right) dx = 0$$

15. If the system of linear equations $2\mathbf{x} + 2\mathbf{y} + 3\mathbf{z} = \mathbf{a}$ 3x - y + 5z = bx - 3y + 2z = cwhere a, b, c are non-zero real numbers, has more then one solution, then : (1) b - c - a = 0(2) a + b + c = 0(4) b - c + a = 0(3) b + c - a = 0Ans. (1) **Sol.** $P_1: 2x + 2y + 3z = a$ $\mathbf{P}_2: \mathbf{3x} - \mathbf{y} + \mathbf{5z} = \mathbf{b}$ $P_3: x - 3y + 2z = c$ We find $\mathbf{P}_1 + \mathbf{P}_3 = \mathbf{P}_2 \Longrightarrow \mathbf{a} + \mathbf{c} = \mathbf{b}$ A square is inscribed in he circle 16.

 $x^2 + y^2 - 6x + 8y - 103 = 0$ with its sides parallel to the corrdinate axes. Then the distance of the vertex of this square which is nearest to the origin is :-

(1) 13 (2)
$$\sqrt{137}$$

(3) 6 (4)
$$\sqrt{41}$$

Ans. (4)

Sol.
$$R = \sqrt{9 + 16 + 103} = 8\sqrt{2}$$

 $OA = 13$
 $OB = \sqrt{265}$
 $OC = \sqrt{137}$
 $OD = \sqrt{41}$
 $(-5,-12)$
 $(-5,-12)$
 $(-5,-12)$
 $(-5,-12)$
 $(-5,-12)$

17. Let $f_k(x) = \frac{1}{k}(\sin^k x + \cos^k x)$ for k = 1, 2,

3, Then for all $x \in R$, the value of $f_4(x) - f_6(x)$ is equal to :-

(1)
$$\frac{5}{12}$$
 (2) $\frac{-1}{12}$ (3) $\frac{1}{4}$ (4) $\frac{1}{12}$

Ans. (4)

Sol. $f_4(x) - f_6(x)$ $=\frac{1}{4}(\sin^{4}x + \cos^{4}x) - \frac{1}{6}(\sin^{6}x + \cos^{6}x)$ $=\frac{1}{4}\left(1-\frac{1}{2}\sin^2 2x\right)-\frac{1}{6}\left(1-\frac{3}{4}\sin^2 2x\right)=\frac{1}{12}$



18. Let [x] denote the greatest integer less than or equal to x. Then :-

$$\lim_{x\to 0}\frac{\tan(\pi\sin^2 x) + \left(|x| - \sin(x[x])\right)^2}{x^2}$$

(1) equals π

- (2) equals 0
- (3) equals $\pi + 1$
- (4) does not exist

Ans. (4)

Sol. R.H.L. = $\lim_{x \to 0^+} \frac{\tan(\pi \sin^2 x) + (|x| - \sin(x[x]))^2}{x^2}$ (as $x \to 0^+ \Rightarrow [x] = 0$) = $\lim_{x \to 0^+} \frac{\tan(\pi \sin^2 x) + x^2}{x^2}$ = $\lim_{x \to 0^+} \frac{\tan(\pi \sin^2 x)}{(\pi \sin^2 x)} + 1 = \pi + 1$ L.H.L. = $\lim_{x \to 0^-} \frac{\tan(\pi \sin^2 x) + (-x + \sin x)^2}{x^2}$ (as $x \to 0^- \Rightarrow [x] = -1$) $\lim_{x \to 0^+} \frac{\tan(\pi \sin^2 x)}{\pi \sin^2 x} \cdot \frac{\pi \sin^2 x}{x^2} + (-1 + \frac{\sin x}{x})^2 \Rightarrow \pi$ R.H.L. \neq L.H.L.

19. The direction ratios of normal to the plane through the points (0, -1, 0) and (0, 0, 1) and making an anlge π/4 with the plane y-z+5=0 are:

(1) $2\sqrt{3}$, 1, -1 (2) 2, $\sqrt{2}$, $-\sqrt{2}$ (3) 2, -1, 1

(4) $\sqrt{2}$, 1, -1

Е

Ans. (2, 4)

Sol. Let the equation of plane be a(x - 0) + b(y + 1) + c(z - 0) = 0It passes through (0,0,1) then $b + c = 0 \qquad \dots(1)$ Now $\cos \frac{\pi}{4} = \frac{a(0) + b(1) + c(-1)}{\sqrt{2}\sqrt{a^2 + b^2 + c^2}}$ $\Rightarrow a^2 = -2bc \text{ and } b = -c$ we get $a^2 = 2c^2$ $\Rightarrow a = \pm\sqrt{2} c$ $\Rightarrow \text{ direction ratio } (a, b, c) = (\sqrt{2}, -1, 1) \text{ or } (\sqrt{2}, 1, -1)$ 20. If $x \log_e(\log_e x) - x^2 + y^2 = 4(y > 0)$, then dy/dx at x = e is equal to :

(1)
$$\frac{e}{\sqrt{4 + e^2}}$$

(2) $\frac{(1+2e)}{2\sqrt{4 + e^2}}$
(3) $\frac{(2e-1)}{2\sqrt{4 + e^2}}$
(4) $\frac{(1+2e)}{\sqrt{4 + e^2}}$

Ans. (3)

Sol. Differentiating with respect to x,

$$x.\frac{1}{\ln x} \cdot \frac{1}{x} + \ln(\ln x) - 2x + 2y \cdot \frac{dy}{dx} = 0$$

at x = e we get
$$1 - 2e + 2y \frac{dy}{dx} = 0 \implies \frac{dy}{dx} = \frac{2e - 1}{2y}$$
$$\implies \frac{dy}{dx} = \frac{2e - 1}{2\sqrt{4 + e^2}} \text{ as } y(e) = \sqrt{4 + e^2}$$



21. The straight line x + 2y = 1 meets the coordinate axes at A and B. A circle is drawn through A, B and the origin. Then the sum of perpendicular distances from A and B on the tangent to the circle at the origin is :

(1)
$$\frac{\sqrt{5}}{4}$$

(2)
$$\frac{\sqrt{5}}{2}$$

- (3) $2\sqrt{5}$
- (4) $4\sqrt{5}$
- Ans. (2)



Equation of circle

$$(x - 1)(x - 0) + (y - 0)\left(y - \frac{1}{2}\right) = 0$$

 $\Rightarrow x^2 + y^2 - x - \frac{y}{2} = 0$

Equation of tangent of origin is 2x + y = 0

$$\ell_1 + \ell_2 = \frac{2}{\sqrt{5}} + \frac{1}{2\sqrt{5}}$$
$$= \frac{4+1}{2\sqrt{5}} = \frac{\sqrt{5}}{2}$$

- If q is false and $p \land q \leftrightarrow r$ is true, then which 22. one of the following statements is a tautology?
 - (1) $(p \lor r) \to (p \land r)$
 - (2) p ∨ r
 - (3) $p \wedge r$

$$(4)(p \land r) \to (p \lor r)$$

Ans. (4)

Sol. Given q is F and $(p \land q) \leftrightarrow r$ is T $\Rightarrow p \land q \text{ is } F$ which implies that r is F \Rightarrow q is F and r is F \Rightarrow (p \land r) is always F \Rightarrow (p \land r) \rightarrow (p \lor r) is tautology.

23. If
$$y(x)$$
 is the solution of the differential equation

$$\frac{dy}{dx} + \left(\frac{2x+1}{x}\right)y = e^{-2x}, x > 0,$$
where $y(1) = \frac{1}{2}e^{-2}$, then :
(1) $y(x)$ is decreasing in (0, 1)
(2) $y(x)$ is decreasing in $\left(\frac{1}{2}, 1\right)$
(3) $y(\log_{e} 2) = \frac{\log_{e} 2}{4}$
(4) $y(\log_{e} 2) = \log_{e} 4$
Ans. (2)
Sol. $\frac{dy}{dx} + \left(\frac{2x+1}{x}\right)y = e^{-2x}$
I.F. $= e^{\int \left(\frac{2x+1}{x}\right)dx} = e^{\int \left(\frac{2+1}{x}\right)dx}} = e^{2x+t/nx} = e^{2x}.x$
So, $y(xe^{2x}) = \int e^{-2x}.xe^{2x} + C$
 $\Rightarrow xye^{2x} = \int x \, dx + C$
 $\Rightarrow 2xye^{2x} = x^{2} + 2C$
It passess through $\left(1, \frac{1}{2}e^{-2}\right)$ we get $C = 0$
 $y = \frac{xe^{-2x}}{2}$
 $\Rightarrow \frac{dy}{dx} = \frac{1}{2}e^{-2x}(-2x+1)$
 $\Rightarrow f(x)$ is decreasing in $\left(\frac{1}{2}, 1\right)$
 $y(\log_{e} 2) = \frac{(\log_{e} 2)e^{-2(\log_{e} 2)}}{2}$
 $= \frac{1}{8}\log_{e} 2$

Sol.





- 24. The maximum value of the function $f(x) = 3x^3 - 18x^2 + 27x - 40$ on the set $S = \{x \in R : x^2 + 30 \le 11x\}$ is : (1) 122(2) - 222(3) - 122(4) 222 Ans. (1)**Sol.** S = {x \in R, x² + 30 - 11x \leq 0} $= \{ x \in \mathbb{R}, 5 \le x \le 6 \}$ Now $f(x) = 3x^3 - 18x^2 + 27x - 40$ $\Rightarrow f'(x) = 9(x - 1)(x - 3),$ which is positive in [5, 6] \Rightarrow f(x) increasing in [5, 6] Hence maximum value = f(6) = 122If one real root of the quadratic equation 25. $81x^2 + kx + 256 = 0$ is cube of the other root, then a value of k is (1) - 81(2) 100(3) - 300(4) 144Ans. (3) **Sol.** $81x^2 + kx + 256 = 0$; $x = \alpha, \alpha^3$ $\Rightarrow \alpha^4 = \frac{256}{81} \Rightarrow \alpha = \pm \frac{4}{3}$ Now $-\frac{k}{81} = \alpha + \alpha^3 = \pm \frac{100}{27}$ \Rightarrow k = ±300 26. Two circles with equal radii are intersecting at
- **26.** Two circles with equal radii are intersecting at the points (0, 1) and (0, -1). The tangent at the point (0, 1) to one of the circles passes through the centre of the other circle. Then the distance between the centres of these circles is :
 - (1) 1 (2) $\sqrt{2}$
 - (3) $2\sqrt{2}$
- Ans. (4)

Ε

Sol. In **AAPO**





(4) 2

So distance between centres = $\sqrt{2} r = 2$

- 27. Equation of a common tangent to the parabola y² = 4x and the hyperbole xy = 2 is :
 (1) x + 2y + 4 = 0
 (2) x 2y + 4 = 0
 (3) x + y + 1 = 0
 (4) 4x + 2y + 1 = 0
 Ans. (1)
- Sol. Let the equation of tangent to parabola

$$y^2 = 4x$$
 be $y = mx + \frac{1}{m}$

It is also a tangent to hyperbola xy = 2

$$\Rightarrow x\left(mx + \frac{1}{m}\right) = 2$$
$$\Rightarrow x^{2}m + \frac{x}{m} - 2 = 0$$

$$D = 0 \implies m = -\frac{1}{2}$$

So tangent is 2y + x + 4 = 0

28. The plane containing the line $\frac{x-3}{2} = \frac{y+2}{-1} = \frac{z-1}{3}$

and also containing its projection on the plane 2x + 3y - z = 5, contains which one of the following points ?

(1) (2, 0, -2) (2) (-2, 2, 2)(3) (0, -2, 2) (4) (2, 2, 0)

Ans. (1)

Sol. The normal vector of required plane

$$= (2\hat{i} - \hat{j} + 3\hat{k}) \times (2\hat{i} + 3\hat{j} - \hat{k})$$
$$= -8\hat{i} + 8\hat{i} + 8\hat{k}$$

So, direction ratio of normal is (-1, 1, 1) So required plane is -(x - 3) + (y + 2) + (z - 1) = 0 $\Rightarrow -x + y + z + 4 = 0$

Which is satisfied by (2, 0, -2)



29. If tangents are drawn to the ellipse $x^2 + 2y^2 = 2$ at all points on the ellipse other than its four vertices then the mid points of the tangents intercepted betwen the coordinate axes lie on the curve :

(1)
$$\frac{x^2}{2} + \frac{y^2}{4} = 1$$
 (2) $\frac{x^2}{4} + \frac{y^2}{2} = 1$
(3) $\frac{1}{2x^2} + \frac{1}{4y^2} = 1$ (4) $\frac{1}{4x^2} + \frac{1}{2y^2} = 1$

Ans. (3)

Sol. Equation of general tangent on ellipse

$$\frac{x}{a \sec \theta} + \frac{y}{b \csc \theta} = 1$$
$$a = \sqrt{2}, b = 1$$
$$\Rightarrow \frac{x}{\sqrt{2} \sec \theta} + \frac{y}{\csc \theta} = 1$$
Let the midpoint be (h, k)

$$h = \frac{\sqrt{2} \sec \theta}{2} \implies \cos \theta = \frac{1}{\sqrt{2}h}$$

and $k = \frac{\csc \theta}{2} \implies \sin \theta = \frac{1}{2k}$
 $\therefore \sin^2 \theta + \cos^2 \theta = 1$
 $\implies \frac{1}{2h^2} + \frac{1}{4k^2} = 1$
 $\implies \frac{1}{2x^2} + \frac{1}{4y^2} = 1$

30. Two integers are selected at random from the set {1, 2,..., 11}. Given that the sum of selected numbers is even, the conditional probability that both the numbers are even is :

(1)
$$\frac{2}{5}$$

(2) $\frac{1}{2}$
(3) $\frac{3}{5}$
(4) $\frac{7}{10}$

Ans. (1)

Sol. Since sum of two numbers is even so either both are odd or both are even. Hence number of elements in reduced samples space

$$= {}^{5}C_{2} + {}^{6}C_{2}$$

so required probability =
$$\frac{{}^{5}C_{2}}{{}^{5}C_{2} + {}^{6}C_{2}}$$

8



TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Friday 11th JANUARY, 2019) TIME : 9 : 30 AM To 12 : 30 PM PHYSICS

1. A body is projected at t = 0 with a velocity 10 ms⁻¹ at an angle of 60° with the horizontal. The radius of curvature of its trajectory at t = 1s is R. Neglecting air resistance and taking acceleration due to gravity g = 10 ms⁻², the value of R is :

(1) 2.5 m	(2) 10.3 m
(3) 2.8 m	(4) 5.1 m

Ans. (3)



- $v_{x} = 10\cos 60^{\circ} = 5 \text{ m/s}$ $v_{y} = 10\cos 30^{\circ} = 5\sqrt{3} \text{ m/s}$ velocity after t = 1 sec. $v_{x} = 5 \text{ m/s}$ $v_{y} = \left| (5\sqrt{3} 10) \right| \text{m/s} = 10 5\sqrt{3}$ $a_{n} = \frac{v^{2}}{R} \Rightarrow R = \frac{v_{x}^{2} + v_{y}^{2}}{a_{n}} = \frac{25 + 100 + 75 100\sqrt{3}}{10\cos\theta}$ $\tan\theta = \frac{10 5\sqrt{3}}{5} = 2 \sqrt{3} \Rightarrow \theta = 15^{\circ}$ $R = \frac{100(2 \sqrt{3})}{10\cos 15} = 2.8\text{m}$
- 2. A particle is moving along a circular path with a constant speed of 10 ms^{-1} . What is the magnitude of the change is velocity of the particle, when it moves through an angle of 60° around the centre of the circle?
- (1) zero (2) 10 m/s (3) $10\sqrt{3}$ m/s (4) $10\sqrt{2}$ m/s Ans. (2)

Sol. Sol. $\overline{\nabla_2}$ $\overline{\nabla_1}$ $\overline{\nabla_2}$ $\overline{\nabla_1}$ $(\pi - \theta)$ $\sqrt{-\overline{\nabla_1}}$ $|\Delta \overline{\nabla}| = \sqrt{\overline{\nabla_1^2 + \overline{\nabla_2^2 + 2\overline{\nabla_1 \nabla_2}}}$ $= 2v \sin \frac{\theta}{2}$ since $[|\overline{\nabla_1}| = |\overline{\nabla_2}|]$ $= (2 \times 10) \times \sin(30^\circ)$ = 10 m/s3. A hydrogen atom, initially in the ground state

3. A hydrogen atom, initially in the ground state is excited by absorbing a photon of wavelength 980Å. The radius of the atom in the excited state, it terms of Bohr radius a_0 , will be : $(h_c = 12500 \text{ eV} - \text{Å})$

(1) $9a_0$ (2) $25a_0$ (3) $4a_0$ (4) $16a_0$ Ans. (4)

Sol. Energy of photon = $\frac{12500}{980}$ =12.75 eV \therefore Electron will excite to n= 4 Since 'R' \propto n²

 \therefore Radius of atom will be $16a_0$

A liquid of density ρ is coming out of a hose pipe of radius a with horizontal speed v and hits a mesh. 50% of the liquid passes through the mesh unaffected. 25% looses all of its momentum and 25% comes back with the same speed. The resultant pressure on the mesh will be :

(1)
$$pv^2$$
 (2) $\frac{3}{4}pv^2$

(3)
$$\frac{1}{2}pv^2$$
 (4) $\frac{1}{4}pv^2$

Ans. (2)

1



E

Sol. Momentum per second carried by liquid per second is ρav^2

net force due to reflected liquid = $2 \times \left[\frac{1}{4}\rho av^2\right]$

net force due to stopped liquid =
$$\frac{1}{4} \rho a v^2$$

- Total force = $\frac{3}{4}\rho av^2$ net pressure = $\frac{3}{4}\rho v^2$
- 5. An electromagnetic wave of intensity 50 Wm⁻² enters in a medium of refractive index 'n' without any loss. The ratio of the magnitudes of electrric fields, and the ratio of the magnitudes of magnetic fields of the wave before and after entering into the medium are respectively, given by :

$$(1) \left(\frac{1}{\sqrt{n}}, \frac{1}{\sqrt{n}}\right)$$
$$(2) \left(\sqrt{n}, \frac{1}{\sqrt{n}}\right)$$
$$(3) \left(\sqrt{n}, \sqrt{n}\right)$$
$$(4) \left(\frac{1}{\sqrt{n}}, \sqrt{n}\right)$$

Ans. (2)

Sol.
$$C = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$$

 $V = \frac{1}{\sqrt{k \epsilon_0 \mu_0}}$ [For transparent medium $\mu_r \approx \mu_0$]
 $\therefore \quad \frac{C}{V} = \sqrt{k} = n$
 $\frac{1}{2} \epsilon_0 E_0^2 C$ = intensity = $\frac{1}{2} \epsilon_0 k E^2 v$
 $\therefore \quad E_0^2 C = k E^2 v$

$$\Rightarrow \frac{E_0^2}{E^2} = \frac{kV}{C} = \frac{n^2}{n} \Rightarrow \frac{E_0}{E} = \sqrt{n}$$

similarly

$$\frac{B_0^2 C}{2\mu_0} = \frac{B^2 v}{2\mu_0} \Longrightarrow \frac{B_0}{B} = \frac{1}{\sqrt{n}}$$

6. An amplitude modulated signal is given by V(t) = 10[1 + 0.3cos(2.2 × 10⁴)]sin(5.5 × 10⁵t). Here t is in seconds. The sideband frequencies (in kHz) are, [Given π = 22/7]
(1) 1785 and 1715
(2) 892.5 and 857.5
(3) 89.25 and 85.75
(4) 178.5 smf 171.5

Ans. (3)

Sol.
$$V(t)=10+\frac{3}{2}[2\cos A \sin B]$$
$$=10+\frac{3}{2}[\sin(A+B)-\sin(A-B)]$$
$$=10+\frac{3}{2}[\sin(57.2\times10^{4}t)-\sin(52.8\times10^{4}t)]$$
$$\omega_{1} = 57.2\times10^{4} = 2\pi f_{1}$$
$$f_{1} = \frac{57.2\times10^{4}}{2\times\left(\frac{22}{7}\right)} = 9.1\times10^{4}$$
$$\approx 91 \text{ KHz}$$
$$f_{2} = \frac{52.8\times10^{4}}{2\times\left(\frac{22}{7}\right)}$$
$$\approx 84 \text{ KHz}$$
$$\lim_{f_{c}-f_{w}} \int_{f_{c}}^{t} \int_{f_{c}+f_{w}}^{t}$$
Side band frequency are

$$f_{1} = f_{c} - f_{w} = \frac{52.8 \times 10^{4}}{2\pi} \approx 85.00 \text{ kHz}$$
$$f_{2} = f_{c} + f_{w} = \frac{57.2 \times 10^{4}}{2\pi} \approx 90.00 \text{ kHz}$$

2



7. The force of interaction between two atoms is

given by
$$F = \alpha \beta \exp\left(-\frac{x^2}{\alpha kt}\right)$$
; where x is the

distance, k is the Boltzmann constant and T is temperature and α and β are two constants. The dimension of β is :

- (1) $M^2L^2T^{-2}$
- (2) $M^{2}LT^{-4}$
- (3) $M^0L^2T^{-4}$
- (4) MLT⁻²
- Ans. (2)

Sol. $F = \alpha \beta e^{\left(\frac{-x^2}{\alpha KT}\right)}$

$$\begin{bmatrix} x^{2} \\ \alpha KT \end{bmatrix} = M^{\circ}L^{\circ}T^{\circ}$$
$$\frac{L^{2}}{[\alpha]ML^{2}T^{-2}} = M^{\circ}L^{\circ}T^{\circ}$$
$$\Rightarrow [\alpha] = M^{-1}T^{2}$$
$$[F] = [\alpha] [\beta]$$
$$MLT^{-2} = M^{-1}T^{2}[\beta]$$
$$\Rightarrow [\beta] = M^{2}LT^{-4}$$

8. The charges Q + q and +q are placed at the vertices of a right-angle isosceles triangle as shown below. The net electrostatic energy of the configuration is zero, it the value of Q is:



(1)
$$\frac{-\sqrt{2}q}{\sqrt{2}+1}$$
 (2) $-2q$

(3)
$$\frac{-q}{1+\sqrt{2}}$$
 (4) +q

Ans. (1)

Ε

Sol. U = K
$$\left[\frac{q^2}{a} + \frac{Qq}{a} + \frac{Qq}{a\sqrt{2}}\right] = 0$$

9. In the circuit shown,



the switch S_1 is closed at time t = 0 and the switch S_2 is kept open. At some later time (t_0) , the switch S_1 is opened and S_2 is closed. The behavious of the current I as a function of time 't' is given by :





Sol. From time t = 0 to $t = t_0$, growth of current takes place and after that decay of current takes place.



most appropriate is (2)



10. Equation of travelling wave on a stretched string of linear density 5 g/m is $y = 0.03 \sin(450 t - 9x)$ where distance and time are measured is SI units. The tension in the string is : (2) 12.5 N (3) 7.5 N (1) 10 N (4) 5 N

Ans. (2)

Sol. $y = 0.03 \sin(450 t - 9x)$

$$v = \frac{\omega}{k} = \frac{450}{9} = 50 \text{ m/s}$$
$$v = \sqrt{\frac{T}{\mu}} \Rightarrow \frac{T}{\mu} = 2500$$
$$\Rightarrow T = 2500 \times 5 \times 10^{-3}$$
$$= 12.5 \text{ N}$$

11. An equilateral triangle ABC is cut from a thin solid sheet of wood. (see figure) D, E and F are the mid-points of its sides as shown and G is the centre of the triangle. The moment of inertia of the triangle about an axis passing through G and perpendicular to the plane of the triangle is I₀. It the smaller triangle DEF is removed from ABC, the moment of inertia of the remaining figure about the same axis is I. Then:





Ans. (4)

Sol. Suppose M is mass and a is side of larger triangle, then
$$\frac{M}{4}$$
 and $\frac{a}{2}$ will be mass and side length of smaller triangle.

$$\frac{I_{removed}}{I_{original}} = \frac{\frac{M}{4}}{M} \cdot \frac{\left(\frac{a}{2}\right)^2}{\left(a\right)^2}$$

$$I_{\text{removed}} = \frac{I_0}{16}$$

So, $I = I_0 - \frac{I_0}{16} = \frac{15I_0}{16}$

12. There are two long co-axial solenoids of same length l. the inner and outer coils have radii r_1 and r_2 and number of turns per unit length n_1 and n₂ respectively. The rate of mutual inductance to the self-inductance of the inner-coil is :

(1)
$$\frac{\mathbf{n}_2}{\mathbf{n}_1} \cdot \frac{\mathbf{r}_2^2}{\mathbf{r}_1^2}$$
 (2) $\frac{\mathbf{n}_2}{\mathbf{n}_1} \cdot \frac{\mathbf{r}_1}{\mathbf{r}_2}$ (3) $\frac{\mathbf{n}_1}{\mathbf{n}_2}$ (4) $\frac{\mathbf{n}_2}{\mathbf{n}_1}$

Ans. (4)

Sol.
$$M = \mu_0 n_1 n_2 \pi r_1^2$$
$$L = \mu_0 n_1^2 \pi r_1^2$$
$$\Rightarrow \frac{M}{L} = \frac{n_2}{n_1}$$

A rigid diatomic ideal gas undergoes an 13. adiabatic process at room temperature,. The relation between temperature and volume of this process is $TV^x = constant$, then x is :

(1)
$$\frac{5}{3}$$
 (2) $\frac{2}{5}$ (3) $\frac{2}{3}$ (4) $\frac{3}{5}$

Ans. (2)

Sol. For adiabatic process : $TV^{\gamma-1} = constant$

For diatomic process : $\gamma - 1 = \frac{7}{5} - 1$

$$\therefore x = \frac{2}{5}$$

14. The gas mixture constists of 3 moles of oxygen and 5 moles of argon at temperature T. Considering only translational and rotational modes, the total inernal energy of the system is: (1) 12 RT (2) 20 RT (3) 15 RT (4) 4 RT)

Sol.
$$U = \frac{f_1}{2}n_1RT + \frac{f_2}{2}n_2RT$$
$$= \frac{5}{2}(3RT) + \frac{3}{2} \times 5RT$$
$$U = 15RT$$

collegebatch.com

15. In a Young's double slit experiment, the path different, at a certain point on the screen,

between two interfering waves is $\frac{1}{9}$ th of

wavelength. The ratio of the intensity at this point to that at the centre of a brigth fringe is close to :

(1) 0.94 (2) 0.74 (3) 0.85 (4) 0.80 **Ans. (3)**

Sol. $\Delta x = \frac{\lambda}{8}$

$$\Delta \phi = \frac{(2\pi)}{\lambda} \frac{\lambda}{8} = \frac{\pi}{4}$$
$$I = I_0 \cos^2\left(\frac{\pi}{8}\right)$$
$$\frac{I}{I_0} = \cos^2\left(\frac{\pi}{8}\right)$$

16. If the deBronglie wavelenght of an electron is equal to 10⁻³ times the wavelength of a photon of frequency 6 × 10¹⁴ Hz, then the speed of electron is equal to :
(Speed of light = 3 × 10⁸ m/s Planck's constant = 6.63 × 10⁻³⁴ J.s Mass of electron = 9.1 × 10⁻³¹ kg)
(1) 1.45 × 10⁶ m/s
(2) 1.7 × 10⁶ m/s
(3) 1.8 × 10⁶ m/s
(4) 1.1 × 10⁶ m/s

Ans. (1)

Ε

Sol.
$$\frac{h}{mv} = 10^{-3} \left(\frac{3 \times 10^8}{6 \times 10^{14}} \right)$$

$$v = \frac{6.63 \times 10^{-34} \times 6 \times 10^{14}}{9.1 \times 10^{-31} \times 3 \times 10^5}$$

v = 1.45 × 10⁶ m/s

17. A slob is subjected to two forces \vec{F}_1 and \vec{F}_2 of same magnitude F as shown in the figure. Force \vec{F}_2 is in XY-plane while force F_1 acts along z-axis at the point $(2\vec{i}+3\vec{j})$. The moment of these forces about point O will be :



Torque for F₂ force $\vec{F}_2 = F\hat{k}$ $\vec{\tau}_2 = 2\hat{i} + 3\hat{j}$ $\vec{\tau}_{F_2} = \vec{\tau}_2 \times \vec{F}_2 = 3F\hat{i} + 2F(-\hat{j})$ $\vec{\tau}_{net} = \vec{\tau}_{F_1} + \vec{\tau}_{F_2}$ $= 3F\hat{i} + 2F(-\hat{j}) + 3F(\hat{k})$

- **18.** A satellite is revolving in a circular orbit at a height h from the earth surface, such that h << R where R is the radius of the earth. Assuming that the effect of earth's atmosphere can be neglected the minimum increase in the speed requried so that the satellite could escapte from the gravitational field of earth is :
 - (1) $\sqrt{\mathrm{gR}}\left(\sqrt{2}-1\right)$

(2)
$$\sqrt{2gR}$$

(3)
$$\sqrt{\mathrm{gR}}$$

(4) $\sqrt{\frac{gR}{2}}$



Ans. (1)

Sol.
$$v_0 = \sqrt{g(R+h)} \approx \sqrt{gR}$$

 $v_e = \sqrt{2g(R+h)} \approx \sqrt{2gR}$
 $\Delta v = v_e - v_0 = (\sqrt{2} - 1)\sqrt{gR}$

19. In an experiment electrons are accelerated, from rest, by applying a voltage of 500 V. Calculate the radius of the path if a magnetic field 100 mT is then applied. [Charge of the electron = 1.6×10^{-19} C Mass of the electron = 9.1×10^{-31} kg] (1) 7.5×10^{-4} m (2) 7.5×10^{-3} m (4) 7.5×10^{-2} m (3) 7.5 m Ans. (1)

Sol.
$$r = \frac{\sqrt{2mk}}{eB} = \frac{\sqrt{2me\Delta v}}{eB}$$

 $r = \frac{\sqrt{\frac{2m}{e}.\Delta v}}{B} = \frac{\sqrt{\frac{2 \times 9.1 \times 10^{-31}}{1.6 \times 10^{-19}}}(500)}{100 \times 10^{-3}}$
 $r = \frac{\sqrt{\frac{9.1}{0.16} \times 10^{-10}}}{10^{-1}} = \frac{3}{.4} \times 10^{-4} = 7.5 \times 10^{-4}$

20. A particle undergoing simple harmonic motion has time dependent displacement given by $x(t) = A \sin \frac{\pi t}{90}$. The ratio of kinetic to potential energy of this particle at t = 210 s will be :

(1) 2 (2)
$$\frac{1}{9}$$
 (3) 3 (4) 1

Ans. (3)

Sol.
$$k = \frac{1}{2}m\omega^2 A^2 \cos^2 \omega t$$

 $U = \frac{1}{2}m\omega^2 A^2 \sin^2 \omega t$
 $\frac{k}{U} = \cot^2 \omega t = \cot^2 \frac{\pi}{90}(210) = \frac{1}{3}$
Hence ratio is 3 (most appropriate)

21. Ice at -20° C os added tp 50 g of water at 40° C. When the temperature of the mixture reaches 0°C, it is found that 20 g of ice is still unmelted. The amount of ice added to the water was close to (Specific heat of water = $4.2 \text{ J/g/}^{\circ}\text{C}$) Specific heat of Ice = $2.1 \text{ J/g/}^{\circ}\text{C}$ Heat of fusion of water at $0^{\circ}C = 334 \text{ J/g}$) (2) 40 g (1) 50 g (3) 60 g (4) 100 g

Ans. (2)

- Sol. Let amount of ice is m gm. According to principal of calorimeter heat taken by ice = heat given by water $\therefore 20 \times 2.1 \times m + (m - 20) \times 334$ $= 50 \times 4.2 \times 40$ 376 m = 8400 + 6680m = 40.1
- \therefore correct answer is (2) 22. In the figure shown below, the charge on the left plate of the 10 μ F capacitor is -30 μ C. ?The charge on the right plate of the 6 μ F capacitor is :



Ans. (4)



6µF & 4µF are in parallel & total charge on this combination is 30 µC

 \therefore Charge on 6µF capacitor = $\frac{6}{6+4} \times 30$

 $= 18 \ \mu C$ Since charge is asked on right plate therefore is $+18\mu$ C Correct answer is (4)

6



23. In the given circuit the current through Zener Diode is close to :





Ε

- Sol. Since voltage across zener diode must be less than 10V therefore it will not work in breakdown region, & its resistance will be infinite & current through it = 0
 - \therefore correct answer is (4)
- 24. The variation of refractive index of a crown glass thin prism with wavelength of the incident light is shown. Which of the following graphs is the correct one, if D_m is the angle of minimum deviation?



collegebatch.com



Ans. (2)

- Sol. Since $D_m = (\mu 1)A$ & on increasing the wavelength, μ decreases & hence D_m decreases. Therefore correct answer is (2)
- 25. The resistance of the meter bridge AB is given figure is 4 Ω . With a cell of emf $\varepsilon = 0.5$ V and rheostat resistance $R_h = 2 \Omega$ the null point is obtained at some point J. When the cell is replaced by another one of emf $\varepsilon = \varepsilon_2$ the same null point J is found for $R_h = 6 \Omega$. The emf ε_2 is;



Sol. Potential gradient with $R_h = 2\Omega$
is
$$\left(\frac{6}{2+4}\right) \times \frac{4}{L} = \frac{dV}{dL}$$
; L = 100 cm

Let null point be at $\ell\ cm$

thus
$$\varepsilon_1 = 0.5 \text{V} = \left(\frac{6}{2+4}\right) \times \frac{4}{L} \times \ell$$
 ...(1)

Now with $R_h = 6\Omega$ new potential gradient is

$$\left(\frac{6}{4+6}\right) \times \frac{4}{L}$$
 and at null point

$$\left(\frac{6}{4+6}\right)\left(\frac{4}{L}\right) \times \ell = \varepsilon_2 \qquad \dots (2)$$

dividing equation (1) by (2) we get

$$\frac{0.5}{\varepsilon_2} = \frac{10}{6} \text{ thus } \varepsilon_2 = 0.3$$

26. The given graph shows variation (with distance r from centre) of :



- (1) Potential of a uniformly charged sphere
- (2) Potential of a uniformly charged spherical shell
- (3) Electric field of uniformly charged spherical shell

(4) Electric field of uniformly charged sphere **Ans.** (2)

27. Two equal resistance when connected in series to a battery, consume electric power of 60 W. If these resistances are now connected in parallel combination to the same battery, the electric power consumed will be :

(1) 60 W	(2) 240 W
(3) 30 W	(4) 120 W

Ans. (2)

Sol. In series condition, equivalent resistance is 2R

thus power consumed is
$$60W = \frac{\epsilon^2}{2R}$$

In parallel condition, equivalent resistance is R/ 2 thus new power is

$$P' = \frac{\epsilon^2}{(R/2)}$$

or P' = 4P = 240W

28. An object is at a distacen of 20 m from a convex lens of focal length 0.3 m. The lens forms an image of the object. If the object moves away from the lens at a speed of 5 m/s, the speed and direction of the image will be :
(1) 0.92 × 10⁻³ m/s away from the lens
(2) 2.26 × 10⁻³ m/s away from the lens
(3) 1.16 × 10⁻³ m/s towards the lens
(4) 3.22 × 10⁻³ m/s towards the lens

Ans. (3)

Sol. From lens equation

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$$

$$\frac{1}{v} - \frac{1}{(-20)} = \frac{1}{(.3)} = \frac{10}{3}$$

$$\frac{1}{v} = \frac{10}{3} - \frac{1}{20}$$

$$\frac{1}{v} = \frac{197}{60}; v = \frac{60}{197}$$

$$m = \left(\frac{v}{u}\right) = \frac{\left(\frac{60}{197}\right)}{20}$$

velocity of image wrt. to lens is given by $v_{I/L} = m^2 v_{O/L}$

direction of velocity of image is same as that of object

 $v_{O/L} = 5 m/s$



$$\mathbf{v}_{\rm I/L} = \left(\frac{60 \times 1}{197 \times 20}\right)^2 (5)$$

= 1.16×10^{-3} m/s towards the lens

- 29. A body of mass 1 kg falls freely from a height of 100 m on a platform of mass 3 kg which is mounted on a spring having spring constant $k = 1.25 \times 10^6$ N/m. The body sticks to the platform and the spring's maximum compression is found to be x. Given that g = 10 ms⁻², the value of x will be close to : (1) 4 cm
 - (2) 8 cm
 - (3) 80 cm
 - (4) 40 cm

Ans. (1)

Sol. Velocity of 1 kg block just before it collides with 3kg block = $\sqrt{2gh} = \sqrt{2000}$ m/s

Applying momentum conversation just before and just after collision.

$$1 \times \sqrt{2000} = 4v \implies v = \frac{\sqrt{2000}}{4} \text{ m/s}$$



initial compression of spring $1.25 \times 10^{6} x_{0} = 30 \Rightarrow x_{0} \approx 0$ applying work energy theorem, $W_{g} + W_{sp} = \Delta KE$ $\Rightarrow 40 \times x + \frac{1}{2} \times 1.25 \times 10^{6} (0^{2} - x^{2})$

$$=0-\frac{1}{2}\times4\times v^{2}$$

solving $x \approx 4$ cm

30. In a Wheatstone bridge (see fig.), Resistances P and Q are approximately equal. When $R = 400 \Omega$, the bridge is equal. When $R = 400 \Omega$, the bridge is balanced. On inter-changing P and Q, the value of R, for balance, is 405 Ω . The value of X is close to :



Ans.	(4)	
	(3) 401.5 ohm	(4) 402.5 ohm
	(1) 403.5 ohm	(2) 404.5 ohm



Ε

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Friday 11th JANUARY, 2019) TIME : 9 : 30 AM To 12 : 30 PM CHEMISTRY

1. For the cell $Zn(s) | Zn^{2+}(aq) || M^{x+}(aq) | M(s)$, different half cells and their standard electrode potentials are given below :

$M^{x+}(aa/M(s))$	Au ³⁺ (aq)/	Ag ⁺ (aq)/	Fe ³⁺ (aq)/	Fe ²⁺ (aq)/
wi (aq/ wi(3)	Au(s)	Ag(s)	$\mathrm{Fe}^{2+}(\mathrm{aq})$	Fe(s)
$E^{o}_{M^{x+}/M^{(v)}}$	1.40	0.80	0.77	-0.44

If $E_{Zn^{2+}/Zn}^{o} = -0.76 V$, which cathode will give a

mximum value of E_{cell}^{o} per electron transferred ?

(1) Fe^{3+} / Fe^{2+}	(2) Ag ⁺ / Ag
(3) Au ³⁺ / Au	(4) Fe ²⁺ / Fe

Ans. (2)

2. The correct match between items-I and II is :

Item-I	Item-II
(Mixture)	(Separation method)
(A) H_2O : Sugar	(P) Sublimation
(B) H_2O : Aniline	(Q) Recrystallization
(C) H_2O : Toluene	(R) Steam distillation
	(S) Differential
	extraction
(1) A-Q, B-R, C-S	(2) A-R, B-P, C-S
(3) A-S, B-R, C-P	(4) A-Q, B-R, C-P

- Ans. (1)
- Sol. (Mixture) (Seperation method) H_2O : Sugar \Rightarrow Recrystallization H_2O : Aniline \Rightarrow Steam distillation H_2O : Toluene \Rightarrow Differential extraction
- **3.** If a reaction follows the Arrhenius equation, the

plot lnk vs $\frac{1}{(RT)}$ gives straight line with a

gradient (-y) unit. The energy required to activate the reactant is :

(1) y unit	(2) –y unit
(3) yR unit	(4) y/R unit

Ans. (1)

- 4. The concentration of dissolved oxygen (DO) in cold water can go upto :
 - (1) 10 ppm (2) 14 ppm
 - (3) 16 ppm (4) 8 ppm

Ans. (1)

- **Sol.** In cold water, dissolved oxygen (DO) can reach a concentration upto 10 ppm
- 5. The major product of the following reaction is:



Ans. (2) Sol.



- 6. Th correct statements among (a) to (d) regarding H₂ as a fuel are :
 - (a) It produces less pollutant than petrol
 - (b) A cylinder of compressed dihydrogen weighs
 ~ 30 times more than a petrol tank producing the same amount of energy
 - (c) Dihydrogen is stored in tanks of metal alloys like NaNi₅
 - (d) On combustion, values of energy released per gram of liquid dihydrogen and LPG are 50 and 142 kJ, respectively
 - (1) b and d only (2) a, b and c only
 - (3) b, c and d only (4) a and c only

Ans. (2)

Sol. Option (a), (b) & (c) are correct answer (NCERT THEORY BASED)

Ε



JEE (Main) Examination-2019/Morning Session/11-01-2019

7. The major poduct of the following reaction is:



Ans. (1)





HO

8. The element that usually does not show variable oxidation states is :

> (1) V (2) Ti (3) Sc (4) Cu

Ans. (3)

Sol. Usally Sc(Scandium) does not show variable oxidation states.

Most common oxidation states of :

(i) Sc : +3

- (ii) V : +2, +3, +4, +5
- (iii) Ti : +2, +3, +4
- (iv) Cu : +1, +2
- 9. An organic compound is estimated through Dumus method and was found to evolve 6 moles of CO₂. 4 moles of H₂O and 1 mole of nitrogen gas. The formula of the compound is :

(1)
$$C_{12}H_8N$$
 (2) $C_{12}H_8N_2$
(3) C_6H_8N (4) $C_6H_8N_2$

Ans. (4)

2

Sol.
$$[C_xH_yN_z] \xrightarrow{Duma} 6CO_2 + 4H_2O + N_2$$

Hence, $C_cH_0N_2$

10. The major product of the following reaction is :



Ans. (2)



11. Among the following compound which one is found in RNA?



Ans. (3)

Sol. For the given structure 'uracil' is found in RNA

JH

Н

collegebatch.com

Е

Which compound(s) out of the following is/are not 12. aromatic ?



Ans. (2)

Sol.	out of the given options only	$\overset{\clubsuit}{\bigtriangleup}$	is aromatic.
------	-------------------------------	---------------------------------------	--------------

Hence (B), (C) and (D) are not aromatic

13. The correct match between Item(I) and Item(II) is :

Item-I	Item-II
(A) Nortehindrone	(P) Anti-biotic
(B) Ofloxacin	(Q) Anti-fertility
(C) Equanil	(R) Hypertension
	(S) Analgesics
(1) A-R, B-P, C-S	(2) A-Q, B-P, C-R
(3) A-R, B-P, C-R	(4) A-Q, B-R, C-S

Ans. (2)

- **Sol.** (A) Norethindrone Antifertility
 - (B) Ofloaxacin Anti-Biotic
 - (C) Equanil Hypertension (traiquilizer)
- 14. Heat treatment of muscular pain involves radiation of wavelength of about 900 nm. Which spectral line of H-atom is suitable for this purpose ?

 $[R_{\rm H} = 1 \times 10^5 \text{ cm}^{-1}, \text{ h} = 6.6 \times 10^{-34} \text{ Js},$ $c = 3 \times 10^8 \text{ ms}^{-1}$]

- (1) Paschen, $5 \rightarrow 3$ (2) Paschen, $\infty \rightarrow 3$
- (3) Lyman, $\infty \rightarrow 1$ (4) Balmer, $\infty \rightarrow 2$

Ans. (2)

Ε

15. Consider the reaction, $N_2(g) + 3H_2(g) \longrightarrow 2NH_3(g)$

The equilibrium constant of the above reaction is K_{P} . If pure ammonia is left to dissociate, the partial pressure of ammonia at equilibrium is given by (Assume that $P_{NH_2} \ll P_{total}$ at equilibrium)

(1)
$$\frac{3^{\frac{3}{2}} K_{p}^{\frac{1}{2}} P^{2}}{4}$$
 (2) $\frac{3^{\frac{3}{2}} K_{p}^{\frac{1}{2}} P^{2}}{16}$
(3) $\frac{K_{p}^{\frac{1}{2}} P^{2}}{16}$ (4) $\frac{K_{p}^{\frac{1}{2}} P^{2}}{4}$

Ans. (2)

16. Match the ores(Column A) with the metals (column B) :

	Column-A	Column-B		
	Ores	Metals		
	(I) Siderite	(a) Zinc		
	(II) Kaolinite	(b) Copper		
	(III) Malachite	(c) Iron		
	(IV) Calamine	(d) Aluminium		
	(1) I-b ; II-c ; III-d ; IV	-a		
	(2) I-c ; II-d ; III-a ; IV	-b		
	(3) I-c ; II-d ; III-b ; IV	-a		
	(4) I-a ; II-b ; III-c ; IV	-d		
Ans.	(3)			
Sol.	Siderite : FeCO ₃			
	Kaolinite : $Al_2(OH)_4Si_2O_5$			
	Malachite : $Cu(OH)_2$. $CuCO_3$			
	Calamine : ZnCO ₃			
17.	The correct order of the a	tomic radii of C, Cs, Al		
	and S is :			
	(1) S < C < Al < Cs	(2) S < C < Cs < Al		
	(3) C < S < Cs < Al	(4) C < S < Al < Cs		
Ans.	(4)			
	In a period	AR↓		
Sal	In a group (AR-Ator	nic radius)		
501.	\checkmark			
	$AR\uparrow$			
	Atomic radii order : C <	S < Al < Cs		
	Atomic radius of C: 170) pm		
	Atomic radius of S: 180	pm		

Atomic radius of Al: 184 pm

Atomic radius of Cs : 300 pm

JEE (Main) Examination-2019/Morning Session/11-01-2019

 $\label{eq:linear} \begin{array}{ll} \textbf{18.} & \mbox{Match the metals (Column I) with the coordination} \\ & \mbox{compound(s) / enzyme(s) (Column II)} \end{array}$

Column-I	Column-II
Metals	Coordination compound(s) / Enzyme(s)
(A)Co	(i) Wilkinson catalyst
(B)Zn	(ii) Chlorophyll
(C) Rh	(iii) Vitamin B ₁₂
(D) Mg	(iv) Carbonic anhydrase
(1) A-ii ; B-i ; C-i	v ; D-iii

- (2) A-iii ; B-iv ; C-i ; D-ii
- (3) A-iv ; B-iii ; C-i ; D-ii
- (4) A-i ; B-ii ; C-iii ; D-iv

Ans. (2)

- **Sol.** (i) Wilkinson catalyst : $RhCl(PPh_3)_3$
 - (ii) Chlorophyll : C₅₅H₇₂O₅N₄Mg
 - (iii) Vitamin B₁₂(also known as

cyanocobalamin) contain cobalt.

- (iv) Carbonic anhydrase contains a zinc ion.
- **19.** A 10 mg effervescent tablet contianing sodium bicarbonate and oxalic acid releases 0.25 ml of CO_2 at T = 298.15 K and p = 1 bar. If molar volume of CO_2 is 25.0 L under such condition, what is the percentage of sodium bicarbonate in each tablet ? [Molar mass of NaHCO₃ = 84 g mol⁻¹]

(1) 16.8	(2) 8.4
(3) 0.84	(4) 33.6

Ans. (1)

20. The major product of the following reaction is :



Ans. (1)

Sol.



21. Two blocks of the same metal having same mass and at temperature T_1 and T_2 , respectively. are brought in contact with each other and allowed to attain thermal equilibrium at constant pressure. The change in entropy, ΔS , for this process is :

(1)
$$2C_{P} \ln\left(\frac{T_{1}+T_{2}}{4T_{1}T_{2}}\right)$$
 (2) $2C_{P} \ln\left[\frac{(T_{1}+T_{2})^{\frac{1}{2}}}{T_{1}T_{2}}\right]$
(3) $C_{P} \ln\left[\frac{(T_{1}+T_{2})^{2}}{4TT_{2}}\right]$ (4) $2C_{P} \ln\left[\frac{T_{1}+T_{2}}{2TT_{2}}\right]$

Ans. (3)

4



E

- **22.** The chloride that CANNOT get hydrolysed is :
 - (1) SiCl_4 (2) SnCl_4
 - $(3) \operatorname{PbCl}_4 \qquad (4) \operatorname{CCl}_4$
- Ans. (4)
- **Sol.** CCl_4 cannot get hydrolyzed due to the absence of vacant orbital at carbon atom.
- 23. For the chemical reaction X → Y, the standard reaction Gibbs energy depends on temperature T (in K) as :

$$\Delta_r G^o \text{ (in kJ mol^{-1})} = 120 - \frac{3}{8} T$$

The major component of the reaction mixture at T is :

- (1) X if T = 315 K
- (2) X if T = 350 K
- (3) Y if T = 300 K
- (4) Y if T = 280 K

Ans. (1)

- 24. The freezing point of a diluted milk sample is found to be -0.2°C, while it should have been -0.5°C for pure milk. How much water has been added to pure milk to make the diluted sample ?
 - (1) 2 cups of water to 3 cups of pure milk
 - (2) 1 cup of water to 3 cups of pure milk
 - (3) 3 cups of water to 2 cups of pure milk
 - (4) 1 cup of water to 2 cups of pure milk

Ans. (3)

- 25. A solid having density of 9×10^3 kg m⁻³ forms face centred cubic crystals of edge length $200\sqrt{2}$ pm. What is the molar mass of the
 - solid ?
 - (Avogadro constant \cong 6 × 10²³ mol⁻¹, $\pi \cong$ 3)
 - (1) $0.0216 \text{ kg mol}^{-1}$ (2) $0.0305 \text{ kg mol}^{-1}$
 - (3) $0.4320 \text{ kg mol}^{-1}$ (4) $0.0432 \text{ kg mol}^{-1}$

Ans. (2)

Ε

26. The polymer obtained from the following reactions is :



Ans. (2) Sol.



- **27.** An example of solid sol is :
 - (1) Butter (2) Gem stones
 - (3) Paint (4) Hair cream
- Ans. (2)
- **28.** Peoxyacetyl nitrate (PAN), an eye irritant is produced by :
 - (1) Acid rain (2) Photochemical smog
 - (3) Classical smog (4) Organic waste

Ans. (2)

- **Sol.** Photochemical smog produce chemicals such as formaldehyde, acrolein and peroxyacetyl nitrate (PAN).
- 29. NaH is an example of :
 (1) Electron-rich hydride (2) Molecular hydride
 (3) Saline hydride (4) Metallic hydride
- Ans. (3)
- **Sol.** NaH is an example of ionic hydride which is also known as saline hydride.
- **30.** The amphoteric hydroxide is : (1) $Ca(OH)_2$ (2) $Be(OH)_2$
 - (3) $Sr(OH)_2$ (4) $Mg(OH)_2$

Ans. (2)

Sol. $Be(OH)_2$ is amphoteric in nature while rest all alkaline earth metal hydroxide are basic in nature.



TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Friday 11th JANUARY, 2019) TIME : 2 : 30 PM To 5 : 30 PM MATHEMATICS

1. If the point $(2, \alpha, \beta)$ lies on the plane which passes through the points (3, 4, 2) and (7, 0, 6) and is perpendicular to the plane 2x - 5y = 15, then $2\alpha - 3\beta$ is equal to :-

(1) 5 (2) 17 (3) 12 (4) 7

Ans. (4)

Sol. Normal vector of plane

$$= \begin{vmatrix} i & j & k \\ 2 & -5 & 0 \\ 4 & -4 & 4 \end{vmatrix} = -4 \left(5\hat{i} + 2\hat{j} - 3\hat{k} \right)$$

equation of plane is 5(x-7)+2y-3(z-6) = 05x + 2y - 3z = 17

2. Let α and β be the roots of the quadratic equation $x^2 \sin \theta - x \ (\sin \theta \cos \theta + 1) + \cos \theta = 0$

$$(0 < \theta < 45^{\circ})$$
, and $\alpha < \beta$. Then $\sum_{n=0}^{\infty} \left(\alpha^{n} + \frac{(-1)^{n}}{\beta^{n}} \right)$

is equal to :-

(1)
$$\frac{1}{1-\cos\theta} + \frac{1}{1+\sin\theta}$$

(2)
$$\frac{1}{1+\cos\theta} + \frac{1}{1-\sin\theta}$$

(3)
$$\frac{1}{1-\cos\theta} - \frac{1}{1+\sin\theta}$$

(4)
$$\frac{1}{1+\cos\theta} - \frac{1}{1-\sin\theta}$$

Ans. (1)

Sol. D = $(1 + \sin\theta \cos\theta)^2 - 4\sin\theta\cos\theta = (1 - \sin\theta \cos\theta)^2$ \Rightarrow roots are β = cosec θ and α = cos θ

$$\Rightarrow \sum_{n=0}^{\infty} \left(\alpha^n + \left(-\frac{1}{\beta} \right)^n \right) = \sum_{n=0}^{\infty} \left(\cos \theta \right)^n + \sum_{n=0}^n \left(-\sin \theta \right)^n$$
$$= \frac{1}{1 - \cos \theta} + \frac{1}{1 + \sin \theta}$$

Let K be the set of all real values of x where the function f(x) = sin |x| - |x| + 2(x - π) cos |x| is not differentiable. Then the set K is equal to:(1) {π}
(2) {0}

(3) ϕ (an empty set) (4) {0, π }

Ans. (3)

Ε

- Sol. $f(x) = \sin|x| |x| + 2(x \pi) \cos x$ $\therefore \sin|x| - |x|$ is differentiable function at x=0 $\therefore k = \phi$
- 4. Let the length of the latus rectum of an ellipse with its major axis along x-axis and centre at the origin, be 8. If the distance between the foci of this ellipse is equal to the length of its minor axis, then which one of the following points lies on it ?

(1)
$$(4\sqrt{3}, 2\sqrt{3})$$
 (2) $(4\sqrt{3}, 2\sqrt{2})$
(3) $(4\sqrt{2}, 2\sqrt{2})$ (4) $(4\sqrt{2}, 2\sqrt{3})$

Ans. (2)

Sol.
$$\frac{2b^2}{a} = 8$$
 and $2ae = 2b$

$$\Rightarrow \frac{b}{a} = e \text{ and } 1 - e^2 = e^2 \Rightarrow e = \frac{1}{\sqrt{2}}$$
$$\Rightarrow b = 4\sqrt{2} \text{ and } a = 8$$

so equation of ellipse is
$$\frac{x^2}{64} + \frac{y^2}{32} = 1$$

5. If the area of the triangle whose one vertex is at the vertex of the parabola, $y^2 + 4(x - a^2) = 0$ and the other two vertices are the points of intersection of the parabola and y-axis, is 250 sq. units, then a value of 'a' is :-

(1)
$$5\sqrt{5}$$
 (2) $(10)^{2/3}$ (3) $5(2^{1/3})$ (4) 5

Ans. (4)

Sol. Vertex is $(a^2,0)$

$$y^2 = -(x - a^2)$$
 and $x = 0 \Longrightarrow (0, \pm 2a)$

Area of triangle is $=\frac{1}{2}.4a.(a^2)=250$

$$\Rightarrow$$
 a³ = 125 or a = 5

1

Collegebatch.com

JEE (Main) Examination-2019/Evening Session/11-01-2019

6. The integral
$$\int_{\pi/6}^{\pi/4} \frac{dx}{\sin 2x (\tan^5 x + \cot^5 x)}$$
 equals :
(1) $\frac{1}{10} \left(\frac{\pi}{4} - \tan^{-1} \left(\frac{1}{9\sqrt{3}} \right) \right)$
(2) $\frac{1}{5} \left(\frac{\pi}{4} - \tan^{-1} \left(\frac{1}{3\sqrt{3}} \right) \right)$
(3) $\frac{\pi}{10}$
(4) $\frac{1}{20} \tan^{-1} \left(\frac{1}{9\sqrt{3}} \right)$
Ans. (1)
Sol. $I = \int_{\pi/6}^{\pi/4} \frac{dx}{\sin 2x (\tan^5 x + \cot^5 x)}$

$$I = \frac{1}{2} \int_{\pi/6}^{\pi/4} \frac{\tan^4 x \sec^2 x \, dx}{\left(1 + \tan^{10} x\right)} \quad \text{Put } \tan^5 x = t$$

$$I = \frac{1}{10} \int_{\left(\frac{1}{\sqrt{3}}\right)^5}^{1} \frac{dt}{1 + t^2} = \frac{1}{10} \left(\frac{\pi}{4} - \tan^{-1} \frac{1}{9\sqrt{3}}\right)$$
7. Let $(x + 10)^{50} + (x - 10)^{50} = a_0 + a_1 x + a_2 x^2 - a_1 x + a_2 x^2 - a_2 x^{50}$ for all $x \in \mathbb{R}$ then $\frac{a_2}{2}$ is equal to:

+ $a_{50} x^{50}$, for all $x \in \mathbb{R}$, then $\overline{a_0}$ is equal to:-(1) 12.50 (2) 12.00 (3) 12.75 (4) 12.25

Ans. (4)

Sol.
$$(10 + x)^{50} + (10 - x)^{50}$$

 $\Rightarrow a_2 = 2.50C_2 + 10^{48}, a_0 = 2.10^{50}$
 $\frac{a_2}{a_0} = \frac{{}^{50}C_2}{10^2} = 12.25$

8. Let a function $f : (0, \infty) \to (0, \infty)$ be defined by $f(x) = \left| 1 - \frac{1}{x} \right|$. Then f is :-(1) Injective only (2) Not injective but it is surjective

- (3) Both injective as well as surjective
- (4) Neither injective nor surjective

Ans. (Bonus)





Remark : If co-domain is $[0,\infty)$, then f(x) will be surjective

9. Let S = {1, 2,, 20}. A subset B of S is said to be "nice", if the sum of the elements of B is 203. Then the probability that a randomly chosen subset of S is "nice" is :-

(1)
$$\frac{6}{2^{20}}$$
 (2) $\frac{5}{2^{20}}$ (3) $\frac{4}{2^{20}}$ (4) $\frac{7}{2^{20}}$

Ans. (2) Sol. 7,

> 1,6 2,5

3,4 1,2,4

$$\mathbf{P} = \frac{5}{2^{20}}$$

10. Two lines $\frac{x-3}{1} = \frac{y+1}{3} = \frac{z-6}{-1}$ and $\frac{x+5}{7} = \frac{y-2}{-6} = \frac{z-3}{4}$ intersect at the point R. The reflection of R in the xy-plane has coordinates :-(1) (2, 4, 7) (2) (-2, 4, 7) (3) (2, -4, -7) (4) (2, -4, 7) **Ans. (3) Sol.** Point on L₁ (λ + 3, 3 λ - 1, - λ + 6) Point on L₂ (7 μ - 5, -6 μ + 2, 4 μ + 3) $\Rightarrow \lambda$ + 3 = 7 μ - 5 ...(i) $3\lambda - 1 = -6\mu + 2$...(ii) $\Rightarrow \lambda = -1, \mu = 1$

Reflection is
$$(2,-4,-7)$$

2



11. The number of functions f from {1, 2, 3, ..., 20} onto {1, 2, 3,, 20} such that f(k) is a multiple of 3, whenever k is a multiple of 4, is :- (1) (15)! × 6!
(2) 5⁶ × 15
(3) 5! × 6!
(4) 6⁵ × (15)!

Ans. (1)

- Sol. f(k) = 3m (3,6,9,12,15,18)for k = 4,8,12,16,20 6.5.4.3.2 ways For rest numbers 15! ways Total ways = 6!(15!)
- **12.** Contrapositive of the statement "If two numbers are not equal, then their squares are not equal." is :-
 - (1) If the squares of two numbers are equal, then the numbers are equal.
 - (2) If the squares of two numbers are equal, then the numbers are not equal.
 - (3) If the squares of two numbers are not equal, then the numbers are equal.
 - (4) If the squares of two numbers are not equal, then the numbers are not equal.

Ans. (1)

- **Sol.** Contrapositive of $p \rightarrow q$ is $\neg q \rightarrow \neg p$
- **13.** The solution of the differential equation,

$$\frac{dy}{dx} = (x - y)^{2}, \text{ when } y(1) = 1, \text{ is :}$$
(1) $\log_{e} \left| \frac{2 - y}{2 - x} \right| = 2(y - 1)$
(2) $\log_{e} \left| \frac{2 - x}{2 - y} \right| = x - y$
(3) $-\log_{e} \left| \frac{1 + x - y}{1 - x + y} \right| = x + y - 2$
(4) $-\log_{e} \left| \frac{1 - x + y}{1 + x - y} \right| = 2(x - 1)$
(4)

Ans. (4)

Ε

Sol.
$$x - y = t \Rightarrow \frac{dy}{dx} = 1 - \frac{dt}{dx}$$

 $\Rightarrow 1 - \frac{dt}{dx} = t^2 \Rightarrow \int \frac{dt}{1 - t^2} = \int 1 dx$
 $\Rightarrow \frac{1}{2} \ell n \left(\frac{1 + t}{1 - t} \right) = x + \lambda$

$$\Rightarrow \frac{1}{2} \ell n \left(\frac{1+x-y}{1-x+y} \right) = x + \lambda \quad \text{given} \quad y(1) = 1$$
$$\Rightarrow \frac{1}{2} \ell n (1) = 1 + \lambda \Rightarrow \lambda = -1$$
$$\Rightarrow \ell n \left(\frac{1+x-y}{1-x+y} \right) = 2(x-1)$$
$$\Rightarrow -\ell n \left(\frac{1-x+y}{1+x-y} \right) = 2(x-1)$$

14. Let A and B be two invertible matrices of order 3 × 3. If det(ABA^T) = 8 and det(AB⁻¹) = 8, then det (BA⁻¹ B^T) is equal to :-

(1) 16 (2)
$$\frac{1}{16}$$
 (3) $\frac{1}{4}$ (4) 1

Ans. (2)

Sol.
$$|A|^2 \cdot |B| = 8$$
 and $\frac{|A|}{|B|} = 8 \implies |A| = 4$ and $|B| = \frac{1}{2}$
 $\therefore \det(BA^{-1} \cdot B^T) = \frac{1}{4} \times \frac{1}{4} = \frac{1}{16}$

15. If $\int \frac{x+1}{\sqrt{2x-1}} dx = f(x)\sqrt{2x-1} + C$, where C is a constant of integration, then f(x) is equal to :-

(1)
$$\frac{1}{3}(x+4)$$
 (2) $\frac{1}{3}(x+1)$
(3) $\frac{2}{3}(x+2)$ (4) $\frac{2}{3}(x-4)$

Ans. (1)

Sol.
$$\sqrt{2x-1} = t \Longrightarrow 2x-1 = t^2 \Longrightarrow 2dx = 2t.dt$$

$$\int \frac{x+1}{\sqrt{2x-1}} dx = \int \frac{\frac{t^2+1}{2}+1}{t} t dt = \int \frac{t^2+3}{2} dt$$
$$= \frac{1}{2} \left(\frac{t^3}{3} + 3t \right) = \frac{t}{6} \left(t^2 + 9 \right) + c$$
$$= \sqrt{2x-1} \left(\frac{2x-1+9}{6} \right) + c = \sqrt{2x-1} \left(\frac{x+4}{3} \right) + c$$
$$\Rightarrow f(x) = \frac{x+4}{3}$$



JEE (Main) Examination-2019/Evening Session/11-01-2019

16. A bag contains 30 white balls and 10 red balls. 16 balls are drawn one by one randomly from the bag with replacement. If X be the number of white balls

drawn, the $\left(\frac{\text{mean of } X}{\text{standard deviation of } X}\right)$ is equal to:-

(2) $\frac{4\sqrt{3}}{3}$ (3) $4\sqrt{3}$ (4) $3\sqrt{2}$

Ans. (3)

(1) 4

- **Sol.** p (probability of getting white ball) = $\frac{30}{40}$
 - $q = \frac{1}{4}$ and n = 16mean = $np = 16 \cdot \frac{3}{4} = 12$

and standard diviation

$$=\sqrt{npq}=\sqrt{16.\frac{3}{4}.\frac{1}{4}}=\sqrt{3}$$

- 17. If in a parallelogram ABDC, the coordinates of A, B and C are respectively (1, 2), (3, 4) and (2, 5), then the equation of the diagonal AD is:-(1) 5x + 3y - 11 = 0 (2) 3x - 5y + 7 = 0(3) 3x + 5y - 13 = 0 (4) 5x - 3y + 1 = 0
- Ans. (4)
- Sol. co-ordinates of point D are (4,7) \Rightarrow line AD is 5x - 3y + 1 = 0
- **18.** If a hyperbola has length of its conjugate axis equal to 5 and the distance between its foci is 13, then the eccentricity of the hyperbola is :-

(1) 2 (2) $\frac{13}{6}$ (3) $\frac{13}{8}$ (4) $\frac{13}{12}$

Ans. (4)

Sol. 2b = 5 and 2ae = 13

$$b^{2} = a^{2}(e^{2} - 1) \Rightarrow \frac{25}{4} = \frac{169}{4} - a^{2}$$
$$\Rightarrow a = 6 \Rightarrow e = \frac{13}{12}$$

19. The area (in sq. units) in the first quadrant bounded by the parabola, $y = x^2 + 1$, the tangent to it at the point (2, 5) and the coordinate axes is :-

(1)
$$\frac{14}{3}$$
 (2) $\frac{187}{24}$ (3) $\frac{37}{24}$ (4) $\frac{8}{3}$
Ans. (3)



Area =
$$\int_{0}^{2} (x^{2} + 1) dx - \frac{1}{2} (\frac{5}{4}) (5) = \frac{37}{24}$$

20. Let $\sqrt{3}\hat{i} + \hat{j}$, $\hat{i} + \sqrt{3}\hat{j}$ and $\beta\hat{i} + (1 - \beta)\hat{j}$ respectively be the position vectors of the points A, B and C with respect to the origin O. If the distance of C from the bisector of the acute angle between OA and OB is $\frac{3}{\sqrt{2}}$, then the sum of all possible values

$$\sqrt{2}$$
 f β is :-

Ans. (2)

0

Sol. Angle bisector is x - y = 0

$$\Rightarrow \frac{|\beta - (1 - \beta)|}{\sqrt{2}} = \frac{3}{\sqrt{2}}$$

$$\Rightarrow |2\beta - 1| = 3$$

$$\Rightarrow \beta = 2 \text{ or } -1$$

21. If $\begin{vmatrix} a - b - c & 2a & 2a \\ 2b & b - c - a & 2b \\ 2c & 2c & c - a - b \end{vmatrix}$

$$= (a + b + c) (x + a + b + c)^2, x \neq 0 \text{ and}$$

$$a + b + c \neq 0, \text{ then } x \text{ is equal to } :-$$

(1) -(a + b + c) (2) 2(a + b + c)
(3) abc (4) -2(a + b + c)
(3) abc (4) -2(a + b + c)
(3) abc (4) -2(a + b + c)
Ans. (4)
Sol. $\begin{vmatrix} a - b - c & 2a & 2a \\ 2b & b - c - a & 2b \\ 2c & 2c & c - a - b \end{vmatrix}$

$$R_1 \rightarrow R_1 + R_2 + R_3$$

$$= \begin{vmatrix} a + b + c & a + b + c & a + b + c \\ 2b & b - c - a & 2b \\ 2c & 2c & c - a - b \end{vmatrix}$$

4

$$\begin{aligned} &= (a+b+c) \begin{vmatrix} 1 & 0 & 0 \\ 2b & -(a+b+c) & 0 \\ 2c & 2c & c-a-b \end{vmatrix} \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 24 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 24 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 24 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 24 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 24 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 24 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 24 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 24 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 25 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 26 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 26 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 26 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 26 \\ &= (a+b+c)(a+b+c)^2 \\ &\Rightarrow x = -2(a+b+c) \end{aligned} 26 \\ &= (a+b+c)(a+b+c) \\ &= (a+b+c)(a+b+c)(a+b+c) \\ &= (a+b+c)(a+b+c)(a+b+c)(a+b+c) \\ &= (a+b+c)(a+b+c)(a+b+c)(a+b+c) \\ &= (a+b+c)(a+b+c)(a+b+c)(a+b+c)(a+b+c) \\ &= (a+b+c)$$

$$g^{2} - c = 4a^{2} \Rightarrow c = (g^{2} - 4a^{2})$$
Putting in equation (1)

$$\Rightarrow 4b^{2} + 4f + g^{2} - 4a^{2} = 0$$

$$\Rightarrow x^{2} + 4y + 4(b^{2} - a^{2}) = 0, \text{ it represent a parabola.}$$
24. If 19th term of a non-zero A.P. is zero, then its
(49th term) : (29th term) is :-
(1) 3 : 1 (2) 4 : 1 (3) 2 : 1 (4) 1 : 3
Ans. (1)
Sol. $a + 18d = 0$...(1)
 $\frac{a + 48d}{a + 28d} = \frac{-18d + 48d}{-18d + 28d} = \frac{3}{1}$
25. Let $f(x) = \frac{x}{\sqrt{a^{2} + x^{2}}} - \frac{d - x}{\sqrt{b^{2} + (d - x)^{2}}}, x \in \mathbb{R},$
where a, b and d are non-zero real constants.
Then :-
(1) f is a decreasing function of x
(2) f is neither increasing nor decreasing
function of x
(3) f is not a continuous function of x
(4) f is an increasing function of x
Ans. (4)
Sol. $f(x) = \frac{x}{\sqrt{a^{2} + x^{2}}} - \frac{d - x}{\sqrt{b^{2} + (d - x)^{2}}}$
 $f'(x) = \frac{a^{2}}{(a^{2} + x^{2})^{3/2}} + \frac{b^{2}}{(b^{2} + (d - x)^{2})^{3/2}} > 0 \forall x \in \mathbb{R}$
 $f(x)$ is an increasing function.
26. Let z be a complex number such that

10. Let z be a complex number such that
$$|z| + z = 3 + i$$
 (where $i = \sqrt{-1}$). Then $|z|$ is equal to :-

(1)
$$\frac{5}{4}$$
 (2) $\frac{\sqrt{41}}{4}$ (3) $\frac{\sqrt{34}}{3}$ (4) $\frac{5}{3}$

Ans. (4)

bl.
$$|z| + z = 3 + i$$

 $z = 3 - |z| + i$
Let $3 - |z| = a \Rightarrow |z| = (3 - a)$
 $\Rightarrow z = a + i \Rightarrow |z| = \sqrt{a^2 + 1}$
 $\Rightarrow 9 + a^2 - 6a = a^2 + 1 \Rightarrow a = \frac{8}{6} = \frac{4}{3}$
 $\Rightarrow |z| = 3 - \frac{4}{3} = \frac{5}{3}$

5



JEE (Main) Examination-2019/Evening Session/11-01-2019

27.	All x satisfying the inequality	29	
	$(\cot^{-1} x)^2 - 7 (\cot^{-1} x) + 10 > 0$, lie in the interval:-		
	$(1) (-\infty, \cot 5) \cup (\cot 4, \cot 2)$		
	(2) (cot 5, cot 4)		
	(3) (cot 2, ∞)		
	(4) $(-\infty, \cot 5) \cup (\cot 2, \infty)$		
Ans.	(4)		
Sol.	$\cot^{-1}x > 5$, $\cot^{-1}x < 2$	A	
	\Rightarrow x < cot5, x > cot2		
28.	Given $\frac{b+c}{11} = \frac{c+a}{12} = \frac{a+b}{13}$ for a $\triangle ABC$ with	So	
	usual notation. If $\frac{\cos A}{\alpha} = \frac{\cos B}{\beta} = \frac{\cos C}{\gamma}$, then		
	the ordered triad (α , β , γ) has a value :-	30	
	(1) (3, 4, 5) (2) (19, 7, 25)		
	(3) (7, 19, 25) (4) (5, 12, 13)		
28.	Ans. (3)		
Sol.	$b + c = 11\lambda, c + a = 12\lambda, a + b = 13\lambda$		
	\Rightarrow a = 7 λ , b = 6 λ , c = 5 λ		
	(using cosine formula)	30	
	$\cos A = \frac{1}{5}, \cos B = \frac{19}{35}, \cos C = \frac{5}{7}$		

29. Let x, y be positive real numbers and m, n positive integers. The maximum value of the expression $\frac{x^{m}y^{n}}{(1+x^{2m})(1+y^{2n})}$ is :(1) $\frac{1}{2}$ (2) $\frac{1}{4}$ (3) $\frac{m+n}{6mn}$ (4) 1

Ans. (2)

Sol.
$$\frac{x^m y^n}{(1+x^{2m})(1+y^{2n})} = \frac{1}{\left(x^m + \frac{1}{x^m}\right)\left(y^n + \frac{1}{y^n}\right)} \le \frac{1}{4}$$

using $AM \ge GM$

30.
$$\lim_{x \to 0} \frac{x \cot(4x)}{\sin^2 x \cot^2(2x)}$$
 is equal to :-
(1) 2 (2) 0 (3) 4 (4) 1

Sol.
$$\lim_{x \to 0} \frac{x \tan^2 2x}{\tan 4x \sin^2 x} = \lim_{x \to 0} \frac{x \left(\frac{\tan^2 2x}{4x^2}\right) 4x^2}{\left(\frac{\tan 4x}{4x}\right) 4x \left(\frac{\sin^2 x}{x^2}\right) x^2} = 1$$

6



 $\alpha:\beta:\gamma \Rightarrow 7:19:25$

Е

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Friday 11th JANUARY, 2019) TIME : 02 : 30 PM To 05 : 30 PM PHYSICS

1. A paramagnetic substance in the form of a cube with sides 1 cm has a magnetic dipole moment of 20×10^{-6} J/T when a magnetic intensity of 60×10^{3} A/m is applied. Its magnetic susceptibility is :-(1) 2.3 × 10⁻² (2) 3.3 × 10⁻²

$$(3) \ 3.3 \times 10^{-4} \qquad (2) \ 5.5 \times 10^{-2} (4) \ 4.3 \times 10^{-2}$$

Sol. $\chi = \frac{1}{H}$

 $I = \frac{Magnetic moment}{Volume}$

$$I = \frac{20 \times 10^{-6}}{10^{-6}} = 20 \text{ N/m}^2$$
$$\chi = \frac{20}{60 \times 10^{+3}} = \frac{1}{3} \times 10^{-3}$$
$$= 0.33 \times 10^{-3} = 3.3 \times 10^{-4}$$

2. A particle of mass m is moving in a straight line with momentum p. Starting at time t = 0, a force F = kt acts in the same direction on the moving particle during time interval T so that its momentum changes from p to 3p. Here k is a constant. The value of T is :-

(1)
$$2\sqrt{\frac{p}{k}}$$
 (2) $\sqrt{\frac{2p}{k}}$ (3) $\sqrt{\frac{2k}{p}}$ (4) $2\sqrt{\frac{k}{p}}$

Ans. (1)

E

Sol.
$$\frac{dp}{dt} = F = kt$$

 $\int_{P}^{3P} dP = \int_{O}^{T} kt dt$
 $2p = \frac{KT^{2}}{2}$
 $T = 2\sqrt{\frac{P}{K}}$

3. Seven capacitors, each of capacitance 2 μ F, are to be connected in a configuration to obtain an

effective capacitance of $\left(\frac{6}{13}\right)\mu F$. Which of

the combinations, shown in figures below, will achieve the desired value ?





Ans. (4)

Sol.
$$C_{eq} = \frac{6}{13} \mu F$$

Therefore three capacitors most be in parallel to get 6 in

$$\frac{1}{C_{eq}} = \frac{1}{3C} + \frac{1}{C} + \frac{1}{C} + \frac{1}{C} + \frac{1}{C}$$

$$C_{eq} = \frac{3C}{13} = \frac{6}{13} \mu F$$





JEE (Main) Examination-2019/Evening Session/11-01-2019

- 4. An electric field of 1000 V/m is applied to an electric dipole at angle of 45° . The value of electric dipole moment is 10^{-29} C.m. What is the potential energy of the electric dipole ? (1) - 9 × 10^{-20} J (2) - 7 × 10^{-27} J (3) - 10 × 10^{-29} J (4) - 20 × 10^{-18} J Ans. (2)
- Sol. $U = -\vec{P}.\vec{E}$ = -PE cos θ = -(10⁻²⁹) (10³) cos 45° = - 0.707 × 10⁻²⁶ J = -7 × 10⁻²⁷ J.
- 5. A simple pendulum of length 1 m is oscillating with an angular frequency 10 rad/s. The support of the pendulum starts oscillating up and down with a small angular frequency of 1 rad/s and an amplitude of 10^{-2} m. The relative change in the angular frequency of the pendulum is best given by :-
 - (1) 10^{-3} rad/s
 - (2) 10^{-1} rad/s
 - (3) 1 rad/s
 - (4) 10^{-5} rad/s

Ans. (1)

Sol. Angular frequency of pendulum

$$\omega = \sqrt{\frac{g_{\rm eff}}{\ell}}$$

 $\therefore \quad \frac{\Delta\omega}{\omega} = \frac{1}{2} \frac{\Delta g_{\text{eff}}}{g_{\text{eff}}}$ $\Delta\omega = \frac{1}{2} \frac{\Delta g}{g} \times \omega$

$$[\omega_s = angular frequency of support]$$

$$\Delta \omega = \frac{1}{2} \times \frac{2A\omega_s^2}{100} \times 100$$

- $\Delta \omega = 10^{-3}$ rad/sec.
- 6. Two rods A and B of identical dimensions are at temperature 30°C. If A is heated upto 180°C and B upto T°C, then the new lengths are the same. If the ratio of the coefficients of linear expansion of A and B is 4 : 3, then the value of T is :-

2

Ans. (2)

- Sol. $\Delta \ell_1 = \Delta \ell_2$ $\ell \alpha_1 \Delta T_1 = \ell \alpha_2 \Delta T_2$ $\frac{\alpha_1}{\alpha_2} = \frac{\Delta T_1}{\Delta T_2}$ $\frac{4}{3} = \frac{T - 30}{180 - 30}$ $\overline{T = 230^{\circ}C}$
- 7. In a double-slit experiment, green light (5303 Å) falls on a double slit having a separation of 19.44 μ m and a width of 4.05 μ m. The number of bright fringes between the first and the second diffraction minima is :-

(1) 09 (2) 10

(3) 04 (4) 05

Ans. (4)

Sol. For diffraction

location of 1st minime

$$y_1 = \frac{D\lambda}{a} = 0.2469 D\lambda$$
 Q

location of 2nd minima

$$y_2 = \frac{2D\lambda}{a} = 0.4938D\lambda$$
 $P = \int y_1$

Now for interference

Path difference at P.

$$\frac{\mathrm{dy}}{\mathrm{D}} = 4.8\lambda$$

path difference at Q

$$\frac{\mathrm{dy}}{\mathrm{D}} = 9.6 \,\lambda$$

So orders of maxima in between P & Q is 5, 6, 7, 8, 9 So 5 bright fringes all present between P & Q.

2



8. An amplitude modulated signal is plotted below :-



Which one of the following best describes the above signal ?

- (1) $(9 + \sin (2.5\pi \times 10^5 \text{ t})) \sin (2\pi \times 10^4 \text{ t})\text{V}$
- (2) $(9 + \sin (4\pi \times 10^4 \text{ t})) \sin (5\pi \times 10^5 \text{ t})\text{V}$
- (3) $(1 + 9\sin (2\pi \times 10^4 \text{ t})) \sin (2.5\pi \times 10^5 \text{ t})\text{V}$
- (4) $(9 + \sin (2\pi \times 10^4 \text{ t})) \sin (2.5\pi \times 10^5 \text{t})\text{V}$

Ans. (4)

- Sol. Analysis of graph says
 - (1) Amplitude varies as 8 10 V or 9 ± 1
 - (2) Two time period as

100 µs (signal wave) & 8 µs (carrier wave)

Hence signal is
$$\left[9\pm1\sin\left(\frac{2\pi t}{T_1}\right)\right]\sin\left(\frac{2\pi t}{T_2}\right)$$

 $= 9 \pm 1\sin((2\pi \times 10^4 t)) \sin (2.5\pi \times 10^5 t)$

9. In the circuit, the potential difference between A and B is :-



(1) 6 V (2) 1 V (3) 3 V Ans. (4)

Sol. Potential difference across AB will be equal to battery equivalent across CD

$$V_{AB} = V_{CD} = \frac{\frac{E_1}{r_1} + \frac{E_2}{r_2} + \frac{E_3}{r_3}}{\frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3}} = \frac{\frac{1}{1} + \frac{2}{1} + \frac{3}{1}}{\frac{1}{1} + \frac{1}{1} + \frac{1}{1}}$$

10. A 27 mW laser beam has a cross-sectional area of 10 mm². The magnitude of the maximum electric field in this electromagnetic wave is given by [Given permittivity of space $\epsilon_0 = 9 \times 10^{-12}$ SI units, Speed of light $c = 3 \times 10^8$ m/s]:-

(1) 1 kV/m (2) 2 kV/m
(3) 1.4 kV/m (4)
$$0.7$$
 kV/m

Ans. (3)

Sol. Intensity of EM wave is given by

$$I = \frac{Power}{Area} = \frac{1}{2} \varepsilon_0 E_0^2 C$$
$$= \frac{27 \times 10^{-3}}{10 \times 10^{-6}} = \frac{1}{2} \times 9 \times 10^{-12} \times E^2 \times 3 \times 10^8$$
$$E = \sqrt{2} \times 10^3 \text{ kv/m}$$
$$= 1.4 \text{ kv/m}$$

11. A pendulum is executing simple harmonic motion and its maximum kinetic energy is K_1 . If the length of the pendulum is doubled and it performs simple harmonic motion with the same amplitude as in the first case, its maximum kinetic energy is K_2 . Then :-

(1)
$$K_2 = \frac{K_1}{4}$$
 (2) $K_2 = \frac{K_1}{2}$

(3)
$$K_2 = 2K_1$$
 (4) $K_2 = K_1$

Ans. (3)

(4) 2 V

Sol. Maximum kinetic energy at lowest point B is given by

 $K = mgl (1 - \cos \theta)$ where θ = angular amp.

 $K_1 = mg_{\ell} (1 - \cos \theta)$ $K_2 = mg(2_{\ell}) (1 - \cos \theta)$ $K_2 = 2K_1.$

12. In a hydrogen like atom, when an electron jumps from the M - shell to the L - shell, the wavelength of emitted radiation is λ . If an electron jumps from N-shell to the L-shell, the wavelength of emitted radiation will be :-

3



 $=\frac{6}{3}=2V$

(1)
$$\frac{27}{20}\lambda$$
 (2) $\frac{16}{25}\lambda$ (3) $\frac{20}{27}\lambda$ (4) $\frac{25}{16}\lambda$

Ans. (3)

Sol. For $M \to L$ steel

for $N \rightarrow L$

$$\frac{1}{\lambda} = K\left(\frac{1}{2^2} - \frac{1}{3^2}\right) = \frac{K \times 5}{36}$$

$$\frac{1}{\lambda'} = K\left(\frac{1}{2^2} - \frac{1}{4^2}\right) = \frac{K \times 3}{16}$$
$$\lambda' = \frac{20}{27}\lambda$$

13. If speed (V), acceleration (A) and force (F) are considered as fundamental units, the dimension of Young's modulus will be :-

(1)
$$V^{-2} A^2 F^2$$
 (2) $V^{-4} A^2 F$
(3) $V^{-4} A^{-2} F$ (4) $V^{-2} A^2 F^{-2}$

Ans. (2)

Sol. $\frac{F}{A} = y \cdot \frac{\Delta \ell}{\ell}$

$$[Y] = \frac{F}{A}$$

Now from dimension

$$F = \frac{ML}{T^2}$$

$$L = \frac{F}{M} \cdot T^2$$

$$L^2 = \frac{F^2}{M^2} \left(\frac{V}{A}\right)^4 \quad \because \quad T = \frac{V}{A}$$

$$L^2 = \frac{F^2}{M^2 A^2} \frac{V^4}{A^2} \qquad F = MA$$

$$L^2 = \frac{V^4}{A^2}$$

$$[Y] = \frac{[F]}{[A]} = F^1 \cdot V^{-4} \cdot A^2$$

A particle moves from the point $(2.0\hat{i} + 4.0\hat{j})$ m, 14. at t = 0, with an initial velocity $(5.0\hat{i} + 4.0\hat{j})$ ms⁻¹. It is acted upon by a constant force which produces constant acceleration а $(4.0\hat{i}+4.0\hat{j})$ ms⁻². What is the distance of the particle from the origin at time 2 s? (1) $20\sqrt{2}$ m (2) $10\sqrt{2}$ m (3) 5 m (4) 15 m Ans. (1) **Sol.** $\vec{S} = (5\hat{i} + 4\hat{j})2 + \frac{1}{2}(4\hat{i} + 4\hat{j})4$ $=10\hat{i} + 8\hat{j} + 8\hat{i} + 8\hat{j}$ $\vec{r}_{f} - \vec{r}_{i} = 18\hat{i} + 16\hat{j}$ $\vec{r}_{\rm f} = 20\hat{i} + 20\hat{j}$

$$|\vec{r}_{f}| = 20\sqrt{2}$$

15. A monochromatic light is incident at a certain angle on an equilateral triangular prism and suffers minimum deviation. If the refractive index of the material of the prism is $\sqrt{3}$, then

the angle of incidence is :-(1) 30° (2) 45° (3) 90° (4) 60°

Ans. (4)

Sol. i = e

$$r_1 = r_2 = \frac{A}{2} = 30^{\circ}$$

by Snell's law

$$1 \times \sin i = \sqrt{3} \times \frac{1}{2} = \frac{\sqrt{3}}{2}$$

16. A galvanometer having a resistance of 20 Ω and 30 divisions on both sides has figure of merit 0.005 ampere/division. The resistance that should be connected in series such that it can be used as a voltmeter upto 15 volt, is :-(1) 80 Ω (2) 120 Ω (3) 125 Ω (4) 100 Ω



Ans. (1)
Sol.
$$R_g = 20\Omega$$

 $N_L = N_R = N = 30$
 $FOM = \frac{I}{\phi} = 0.005 \text{ A/Div.}$
Current sentivity = $CS = \left(\frac{1}{0.005}\right)$
 $Ig_{max} = 0.005 \times 30$
 $= 15 \times 10^{-2} = 0.15$
 $15 = 0.15 [20 + R]$
 $100 = 20 + R$
 $R = 80$

17. The circuit shown below contains two ideal diodes, each with a forward resistance of 50Ω. If the battery voltage is 6 V, the current through the 100 Ω resistance (in Amperes) is :-



(1) 0.027	(2) 0.020
(3) 0.030	(4) 0.036

Ans. (2)

Sol. I = $\frac{6}{300}$ = 0.002 (D₂ is in reverse bias)

18. When 100 g of a liquid A at 100°C is added to 50 g of a liquid B at temperature 75°C, the temperature of the mixture becomes 90°C. The temperature of the mixture, if 100 g of liquid A at 100°C is added to 50 g of liquid B at 50°C, will be :-

(1)
$$80^{\circ}$$
C (2) 60° C
(3) 70° C (4) 85° C

Ans. (1)

Sol. $100 \times S_A \times [100 - 90] = 50 \times S_B \times (90 - 75)$ $2S_A = 1.5 S_B$

Now,
$$100 \times S_A \times [100 - T] = 50 \times S_B (T - 50)$$

 $2 \times \left(\frac{3}{4}\right) (100 - T) = (T - 50)$
 $300 - 3T = 2T - 100$
 $400 = 5T$
 $T = 80$

19. The mass and the diameter of a planet are three times the respective values for the Earth. The period of oscillation of a simple pendulum on the Earth is 2s. The period of oscillation of the same pendulum on the planet would be :-

(1)
$$\frac{2}{\sqrt{3}}$$
 s (2) $2\sqrt{3}$ s

3)
$$\frac{\sqrt{3}}{2}$$
s (4) $\frac{3}{2}$ s

Ans. (2)

(

Sol.
$$\because g = \frac{GM}{R^2}$$

 $\frac{g_p}{g_e} = \frac{M_e}{M_e} \left(\frac{R_e}{R_p}\right)^2 = 3\left(\frac{1}{3}\right)^2 = \frac{1}{3}$
Also $T \propto \frac{1}{\sqrt{g}}$
 $\Rightarrow \frac{T_p}{T_e} = \sqrt{\frac{g_e}{g_p}} = \sqrt{3}$
 $\Rightarrow T = 2\sqrt{3}s$

20. The region between y = 0 and y = d contains a magnetic field $\vec{B} = B\hat{z}$. A particle of mass m and charge q enters the region with a velocity

$$\vec{v} = v\hat{i}$$
. If $d = \frac{mv}{2qB}$, the acceleration of the

charged particle at the point of its emergence at the other side is :-

E



 $S_A = \frac{3}{4}S_B$

(1)
$$\frac{qvB}{m} \left(\frac{\hat{i} + \hat{j}}{\sqrt{2}} \right)$$

(2)
$$\frac{qvB}{m} \left(\frac{1}{2} \hat{i} - \frac{\sqrt{3}}{\sqrt{2}} \hat{j} \right)$$

(3)
$$\frac{qvB}{m} \left(\frac{-\hat{j} + \hat{i}}{\sqrt{2}} \right)$$

(4)
$$\frac{qvB}{m} \left(\frac{\sqrt{3}}{2} \hat{i} + \frac{1}{2} \hat{j} \right)$$

Ans. (BONUS)

21. A thermometer graduated according to a linear scale reads a value x_0 when in contact with boiling water, and $x_0/3$ when in contact with ice.

What is the temperature of an object in 0 °C, if this thermometer in the contact with the object reads $x_0/2$?

- (1) 35 (2) 25
- (3) 60 (4) 40

Ans. (2)



$$\Rightarrow T^{\circ}C = \frac{x_0}{6} \& \left(x_0 - \frac{x_0}{3}\right) = (100 - 0^{\circ}C)$$
$$x_0 = \frac{300}{2}$$

$$\Rightarrow T^{\circ}C = \frac{150}{6} = 25^{\circ}C$$

22. A string is wound around a hollow cylinder of mass 5 kg and radius 0.5 m. If the string is now pulled with a horizontal force of 40 N, and the cylinder is rolling without slipping on a horizontal surface (see figure), then the angular acceleration of the cylinder will be (Neglect the mass and thickness of the string) :-

$$40N$$
12 rad/s²
(2) 16 rad/s²
10 rad/s²
(4) 20 rad/s²

Ans. (2)

(1)

(3)

Sol.
$$\longrightarrow_{f}^{40} \xrightarrow{\alpha} a$$

 $\begin{array}{l} 40 \,+\, f = \,m(R\alpha) \,\,(i) \\ 40 \,\times\, R \,-\, f \,\times\, R \,=\, mR^2\alpha \\ 40 \,-\, f \,=\, mR\alpha \,\,\, \,\, (ii) \\ From \,\, (i) \,\, and \,\, (ii) \end{array}$

$$\alpha = \frac{40}{mR} = 16$$

23. In a process, temperature and volume of one mole of an ideal monoatomic gas are varied according to the relation VT = K, where K is a constant. In this process the temperature of the gas is incressed by ΔT . The amount of heat absorbed by gas is (R is gas constant) :

(1)
$$\frac{1}{2}R\Delta T$$
 (2) $\frac{3}{2}R\Delta T$
(3) $\frac{1}{2}KR\Delta T$ (4) $\frac{2K}{3}\Delta T$
Ans. (1)
Sol. VT = K
 $\Rightarrow V\left(\frac{PV}{nR}\right) = k \Rightarrow PV^2 = K$
 $\therefore C = \frac{R}{1-x} + C_V$ (For polytropic process)
 $C = \frac{R}{1-2} + \frac{3R}{2} = \frac{R}{2}$
 $\therefore \Delta Q = nC \Delta T$



$$=\frac{R}{2} \times \Delta T$$

24. In a photoelectric experiment, the wavelength of the light incident on a metal is changed from 300 nm to 400 nm. The decrease in the stopping

potential is close to :
$$\left(\frac{hc}{e} = 1240 \text{ nm} - \text{V}\right)$$

(1) 0.5 V (2) 1.0 V
(3) 2.0 V (4) 1.5 V

Ans. (2)

Sol. $\frac{hc}{\lambda_1} = \phi + eV_1$ (i)

$$\frac{hc}{\lambda_2} = \phi + eV_2 \qquad \dots \dots (ii)$$

$$hc\left(\frac{1}{\lambda_1} - \frac{1}{\lambda_2}\right) = e(V_1 - V_2)$$

$$\Rightarrow V_1 - V_2 = \frac{hc}{e} \left(\frac{\lambda_2 - \lambda_1}{\lambda_1 - \lambda_2} \right)$$
$$= (1240 \text{nm} - \text{V}) \frac{100 \text{nm}}{300 \text{nm} \times 400 \text{nm}}$$
$$= 1 \text{V}$$

25. A metal ball of mass 0.1 kg is heated upto 500°C and dropped into a vessel of heat capacity 800 JK^{-1} and containing 0.5 kg water. The initial temperature of water and vessel is 30°C. What is the approximate percentage increment in the temperature of the water ? [Specific Heat Capacities of water and metal are, $Jkg^{-1}K^{-1}$ respectively, 4200 and $400 \text{ JKg}^{-1}\text{K}^{-1}$] (1) 30%(2) 20% (3) 25%(4) 15%

Ε

Sol. $0.1 \times 400 \times (500 - T) = 0.5 \times 4200 \times (T - 30)$ + 800 (T - 30) $\Rightarrow 40(500 - T) = (T - 30) (2100 + 800)$ $\Rightarrow 20000 - 40T = 2900 T - 30 \times 2900$ $\Rightarrow 20000 + 30 \times 2900 = T(2940)$ T = 30.4°C

$$\frac{\Delta T}{T} \times 100 = \frac{6.4}{30} \times 100$$

26. The magnitude of torque on a particle of mass 1kg is 2.5 Nm about the origin. If the force acting on it is 1 N, and the distance of the particle from the origin is 5m, the angle between the force and the position vector is (in radians) :-

(1)
$$\frac{\pi}{8}$$
 (2) $\frac{\pi}{6}$ (3) $\frac{\pi}{4}$ (4) $\frac{\pi}{3}$

Ans. (2)

Sol. 2.5 = 1 × 5 sin
$$\theta$$

sin θ = 0.5 = $\frac{1}{2}$
 $\theta = \frac{\pi}{6}$

27. In the experimental set up of metre bridge shown in the figure, the null point is obtained at a distance of 40 cm from A. If a 10 Ω resistor is connected in series with R₁, the null point shifts by 10 cm. The resistance that should be connected in parallel with (R₁ + 10) Ω such that the null point shifts back to its initial position is





Ans. (2)
Sol.
$$\frac{R_1}{R_2} = \frac{2}{3}$$
(i)
 $\frac{R_1 + 10}{R_2} = 1 \implies R_1 + 10 = R_2$ (ii)
 $\frac{2R_2}{3} + 10 = R_2$
 $10 = \frac{R_2}{3} \implies R_2 = 30\Omega$
& $R_1 = 20\Omega$
 $\frac{30 \times R}{30 + R} = \frac{2}{3}$
 $R = 60 \Omega$

28. A circular disc D_1 of mass M and radius R has two identical discs D_2 and D_3 of the same mass M and radius R attached rigidly at its opposite ends (see figure). The moment of inertia of the system about the axis OO', passing through the centre of D_1 , as shown in the figure, will be:-



Ans. (1)

Sol.
$$I = \frac{MR^2}{2} + 2\left(\frac{MR^2}{4} + MR^2\right)$$

= $\frac{MR^2}{2} + \frac{MR^2}{2} + 2MR^2$
= $3 MR^2$

29. A copper wire is wound on a wooden frame, whose shape is that of an equilateral triangle. If the linear dimension of each side of the frame is increased by a factor of 3, keeping the number of turns of the coil per unit length of the frame the same, then the self inductance of the coil :

(1) Decreases by a factor of $9\sqrt{3}$

(2) Increases by a factor of 3

(3) Decreases by a factor of 9

(4) Increases by a factor of 27

Ans. (2)

Sol. Total length L will remain constant

 $\label{eq:L} \begin{array}{ll} L = (3a) \ N & (N = total \ turns) \\ \text{and length of winding} = (d) \ N \end{array}$

(d = diameter of wire)



 $\propto a^2 N \propto a$

self inductance = $\mu_0 n^2 A \ell$

$$= \mu_0 n^2 \left(\frac{\sqrt{3} a^2}{4}\right) dN$$

So self inductance will become 3 times

30. A particle of mass m and charge q is in an electric and magnetic field given by

 $\vec{E} = 2\hat{i} + 3\hat{j} \ ; \ \vec{B} = 4\hat{j} + 6\hat{k} \, .$

The charged particle is shifted from the origin to the point P(x = 1; y = 1) along a straight path. The magnitude of the total work done is :-(1) (0.35)q (2) (0.15)q (3) (2.5)q (4) 5q

Ans. (4)

Sol.
$$\vec{F}_{net} = q\vec{E} + q(\vec{v} \times \vec{B})$$

= $(2q\hat{i} + 3q\hat{j}) + q(\vec{v} \times \vec{B})$
 $W = \vec{F}_{net} \cdot \vec{S}$
= $2q + 3q$
= $5q$

8



TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Friday 11th JANUARY, 2019) TIME : 02 : 30 PM To 05 : 30 PM CHEMISTRY



collegebatch.com

Sol.



6. The number of bridging CO ligand (s) and Co-Co bond (s) in CO₂(CO)g, respectively are :-

- (1) 0 and 2 (2) 2 and 0
- (3) 4 and 0 (4) 2 and 1

Ans. (4)

Sol.



Bridging CO are 2 and Co – Co bond is 1.

7. In the following compound,



the favourable site/s for protonation is/are :-

(1) (b), (c) and (d) (2) (a)

(3) (a) and (e) (4) (a) and (d)

Ans. (1)

- Sol. Localised lone pair e⁻.
- 8. The higher concentration of which gas in air can cause stiffness of flower buds ?

(1) SO_2 (2) NO_2 (3) CO_2 (4) CO

Ans. (1)

Sol. Due to acid rain in plants high concentration of SO_2 makes the flower buds stiff and makes them fall.

9. The correct match between item I and item II is :-

	Item I	Item II					
(A)	Allosteric	(P) Molecule binding					
	effect		to the active site				
			ofenzyme				
(B)	Competitive	(Q)	Molecule crucial				
	inhibitor		for				
			communication in				
		the body					
(C)	Receptor	(R)	Molecule binding				
		to a site other than					
		the active site of					
			enzyme				
(D)	Poison	(S)	Molecule binding				
			to the enzyme				
			covalently				

(1) (A)
$$\rightarrow$$
(P); (B) \rightarrow (R); (C) \rightarrow (S); (D) \rightarrow (Q)
(2) (A) \rightarrow (R); (B) \rightarrow (P); (C) \rightarrow (S); (D) \rightarrow (Q)
(3) (A) \rightarrow (P); (B) \rightarrow (R); (C) \rightarrow (Q); (D) \rightarrow (S)

(4) (A)
$$\rightarrow$$
(B) \cdot (B) \rightarrow (P) \cdot (C) \rightarrow (O) \cdot (D) \rightarrow (S)

Ans. (4)

10. The radius of the largest sphere which fits properly at the centre of the edge of body centred cubic unit cell is : (Edge length is represented by 'a') :-

(1) 0.134 a	(2) 0.027 a
(3) 0.067 a	(4) 0.047 a

Ans. (3)

Sol.

a = 2(R + r)

$$\frac{a}{2} = (R + r) \dots (1)$$

 $a\sqrt{3} = 4R \dots (2)$
Using (1) & (2)

$$\frac{a}{2} = \frac{a\sqrt{3}}{4} = r$$

$$a\left(\frac{2-\sqrt{3}}{4}\right) = 1$$

$$r = 0.067 a$$



2



E

- 11. Among the colloids cheese (C), milk (M) and smoke (S), the correct combination of the dispersed phase and dispersion medium, respectively is :-
 - (1) C : solid in liquid; M : solid in liquid;S : solid in gas
 - (2) C : solid in liquid; M : liquid in liquid; S : gas in solid
 - (3) C : liquid in solid; M : liquid in solid; S : solid in gas
 - (4) C : liquid in solid; M : liquid in liquid;S : solid in gas

Ans. (4)

Sol.

	Dispersed Phase	Dispersion Medium
Cheese	Liquid	Solid
Milk	Liquid	Liquid
Smoke	Solid	Gas

12. The reaction that does NOT define calcination is:-

(1)
$$\operatorname{ZnCO}_3 \xrightarrow{\Delta} \operatorname{ZnO} + \operatorname{CO}_2$$

(2) $\operatorname{Fe}_2\operatorname{O}_3 \cdot \operatorname{XH}_2\operatorname{O} \xrightarrow{\Delta} \operatorname{Fe}_2\operatorname{O}_3 + \operatorname{XH}_2\operatorname{O}$
(3) $\operatorname{CaCO}_3 \cdot \operatorname{MgCO}_3 \xrightarrow{\Delta} \operatorname{CaO} + \operatorname{MgO} + 2 \operatorname{CO}_2$

$$(4) 2 \operatorname{Cu}_2 S + 3 \operatorname{O}_2 \xrightarrow{\Delta} 2 \operatorname{Cu}_2 O + 2 \operatorname{SO}_2$$

Ans. (4)

Sol. Calcination in carried out for carbonates and oxide ores in absence of oxygen. Roasting is carried out mainly for sulphide ores in presence of excess of oxygen.

13. The reaction,

MgO(s) + C(s)→Mg(S) + CO(g), for which $\Delta_r H^o$ = + 491.1 kJ mol⁻¹ and $\Delta_r S^o$ = 198.0 JK⁻¹ mol⁻¹, is not feasible at 298 K. Temperature above which reaction will be feasible is :-

(1) 1	890.0	K	(2)	2480.3	K
(3) 2	2040.5	K	(4)	2380.5	K

Ans. (2)

Ε

Sol.
$$T_{eq} = \frac{\Delta H}{\Delta S}$$
$$= \frac{491.1 \times 1000}{198}$$
$$= 2480.3 \text{ K}$$

14. Given the equilibrium constant : KC of the reaction : $Cu(s) + 2Ag^{+}(aq) \rightarrow Cu^{2+}(aq) + 2Ag(s)$ is 10×10^{15} , calculate the E_{cell}^{0} of this reaction at 298 K

$$\left[2.303 \frac{\text{RT}}{\text{F}} \text{ at } 298 \text{ K} = 0.059 \text{ V}\right]$$
(1) 0.04736 V (2) 0.4736 V
(3) 0.4736 mV (4) 0.04736 mV

Ans. (2)

Sol.
$$E_{cell} = E_{cell}^{\circ} - \frac{0.059}{n} \log Q$$

At equilibrium
$$E_{Cell}^{\circ} = \frac{0.059}{2} \log 10^{16}$$
$$= 0.059 \times 8$$
$$= 0.472 \text{ V}$$

15. The hydride that is NOT electron deficient is:-

(1)
$$B_2H_6$$
 (2) AlH_3

$$(3) \operatorname{SiH}_4 \qquad (4) \operatorname{GaH}_3$$

Ans. (3)

Sol. (1) B_2H_6 : Electron deficient

(3) SiH_4 : Electron precise

(4) GaH₃ : Electron deficient



JEE (Main) Examination-2019/Evening Session/11-01-2019

19. 16. The standard reaction Gibbs energy for a The reaction $2X \rightarrow B$ is a zeroth order reaction. chemical reaction at an absolute temperature T is given by $\Delta_{r}G^{o} = A - Bt$ concentration of 0.2 M will be :-Where A and B are non-zero constants. Which (1) 18.0 h (2) 7.2 h (3) 9.0 h of the following is TRUE about this reaction ? Ans. (1) (1) Exothermic if B < 0Sol. For zero order (2) Exothermic if A > 0 and B < 0 $[A_0] - [A_t] = kt$ (3) Endothermic if A < 0 and B > 0 $0.2 - 0.1 = k \times 6$ (4) Endothermic if A > 0Ans. (4) $k = \frac{1}{60} M/hr$ Sol. Theory 17. K_2 HgI₄ is 40% ionised in aqueous solution. The and $0.5-0.2 = \frac{1}{60} \times t$ value of its van't Hoff factor (i) is :-(2) 2.2(3) 2.0(1) 1.8(4) 1.6t = 18 hrs. Ans. (1) 20. **Sol.** For $K_2[HgI_4]$ i = 1 + 0.4 (3 - 1)= 1.8(1) CH₃COCH₂NHCH₃ (2) CH₃CH₂COCH₂NH₂ 18. The de Broglie wavelength (λ) associated with a photoelectron varies with the frequency (v)(3) CH₃CH₂CH₂CONH₂ of the incident radiation as, $[v_0]$ is thrshold (4) $CH_3CON(CH_3)_2$ frequency] : Ans. (3)

(1)
$$\lambda \propto \frac{1}{(v-v_0)^{\frac{3}{2}}}$$
 (2) $\lambda \propto \frac{1}{(v-v_0)^{\frac{1}{2}}}$
(3) $\lambda \propto \frac{1}{(v-v_0)^{\frac{1}{4}}}$ (4) $\lambda \propto \frac{1}{(v-v_0)}$

Ans. (2)

Sol. For electron

$$\lambda_{DB} = \frac{\lambda}{\sqrt{2mK.E.}} \quad (\text{de broglie wavelength})$$

By photoelectric effect
$$hv = hv_0 + KE$$

$$KE = hv -hv_0$$

$$\lambda_{DB} = \frac{h}{\sqrt{2m \times (hv - hv_0)}}$$

$$\lambda_{DB} \propto \frac{1}{(v - v_0)^{\frac{1}{2}}}$$

If the initial concentration of X is 0.2 M, the half-life is 6 h. When the initial concentration of X is 0.5 M, the time required to reach its final (4) 12.0 h

A compound 'X' on treatment with Br₂/NaOH, provided C₃H₉N, which gives positive carbylamine test. Compound 'X' is :-

Sol.

$$[X] \xrightarrow{Br_2} C_3H_9N \xrightarrow{CHCl_3} CH_3CH_2CH_2-NC$$

Hoff mann's Carbylamine
Bromaide Reaction
degradation

Thus [X] must be aride with oen carbon more than is amine.

Thus [X] is CH₂CH₂CH₂CONH₂

Which of the following compounds will 21. produce a precipitate with AgNO₃?



Ans. (4)



133

Sol.



as it can produce aromatic cation so will produce precipitate with $AgNO_3$.

- **22.** The relative stability of +1 oxidation state of group 13 elements follows the order :-
 - (1) Al < Ga < Tl < In (2) Tl < In < Ga < Al
 - (3) Al < Ga < In < Tl (4) Ga < Al < In < Tl

Ans. (3)

Sol. Due to inert pair effect as we move down the group in 13th group lower oxidation state becomes more stable.

 $Al < Ga < In < T\ell$

23. Which of the following compounds reacts with ethylmagnesium bromide and also decolourizes bromine water solution :-

0

(2)
$$Ch_2 - CO_2CH_3$$





Ans. (4)

Sol.



24. Match the following items in column I with the corresponding items in column II.

	Column I	Column II			
(i)	Na ₂ CO ₃ ·10 H ₂ O	(P) Portland cemen			
			ingredient		
(ii)	$Mg(HCO_3)_2$	(Q) Castner-Keller			
			process		
(iii)	NaOH	(R)	Solvay process		
(iv)	$Ca_3Al_2O_6$	(S)	Temporary		
			hardness		

(1) (i)
$$\rightarrow$$
(C); (ii) \rightarrow (B); (iii) \rightarrow (D); (iv) \rightarrow (A)

(2) (i)
$$\rightarrow$$
(C); (ii) \rightarrow (D); (iii) \rightarrow (B); (iv) \rightarrow (A)

(3) (i)
$$\rightarrow$$
(D); (ii) \rightarrow (A); (iii) \rightarrow (B); (iv) \rightarrow (C)

(4) (i)
$$\rightarrow$$
(B); (ii) \rightarrow (C); (iii) \rightarrow (A); (iv) \rightarrow (D)

Ans. (2)

Sol. $Na_2CO_3.10H_2O \rightarrow Solvay process$

 $Mg(HCO_3)_2 \rightarrow Temporary hardness$

 $NaOH \rightarrow Castner-kellner cell$

 $Ca_3Al_2O_6 \rightarrow Portland cement$

25. 25 ml of the given HCl solution requires 30 mL of 0.1 M sodium carbonate solution. What is the volume of this HCl solution required to titrate 30 mL of 0.2 M aqueous NaOH solution?

(1) 25 mL (2) 50 mL (3) 12.5 mL(4) 75 mL

Sol. HCl with Na_2CO_3

Eq. of HCl = Eq. of Na_2CO_3

$$\frac{25}{1000} \times M \times 1 = \frac{30}{1000} \times 0.1 \times 2$$

$$M = \frac{6}{25}M$$

Eq of HCl = Eq. of NaOH

$$\frac{6}{25} \times 1 \times \frac{V}{1000} = \frac{30}{1000} \times 0.2 \times 1$$

$$V = 25 ml$$

Е



JEE (Main) Examination-2019/Evening Session/11-01-2019

26.	$\underline{A} \xrightarrow{4 \text{ KOH, } O_2} 2\underline{B} + 2 \text{ H}_2\text{O}$ (Green)						
	$3 \xrightarrow{B} \xrightarrow{4 \text{ HCl}} 2 \xrightarrow{C} + \text{MnO}_2 + 2 \text{ H}_2\text{O}$						
	(Purple)						
	$2 \xrightarrow{\text{B}} \xrightarrow{\text{H}_2\text{O}, \text{KI}} 2 \xrightarrow{\text{A}} + 2\text{KOH} + \overrightarrow{\text{D}}$						
	In the above sequence of reactions,						
	\underline{A} and \underline{D} respectively,	are :-					
	(1) KIO_3 and MnO_2	(2) KI and $K_2 MnO_4$					
	(3) MnO_2 and KIO_3	(4) KI and $KMnO_4$					
Ans.	(3)						
Sol.	$MnO_2(A) \xrightarrow{4KOH,O_2} 2$	$K_2MnO_4(B) + 2H_2O$					
	(Green)					
	$3K_2MnO_4(B) \xrightarrow{4HCl} 3K_2MnO_4(B)$	$2KMnO_4(C) + 2H_2O$					
		(Purple)					
	$2KMnO_4(C) \xrightarrow{H_2O, KI} \rightarrow$	$2MnO_2(A) + 2KOH +$					
		$KIO_3(D)$					
	$A \rightarrow MnO_2$ $D \rightarrow KIO$						
27.	The coordination	number of Th in					
	$K_4[Th(C_2O_4]_4(OH_2)_2]$ is	8 :-					
	$\left(C_2 O_4^{2-} = \text{Oxalato}\right)$						
	(1) 6 (2) 10	(3) 14 (4) 8					
Ans.	(2)						
Sol.	$C_2 O_4^{2-}$ (oxalato) : biden	ntate					
	H_2O (aqua) : Monoder	ntate					
28.	The major product obtained in the following						
	reaction is :-						
	O _► OH	Li AlH.					
		(excess)					
	CH ₃						
	$\dot{NO}_2 \ddot{O}$	OU					
	U UH UH						
	(1) $\downarrow \downarrow \downarrow$						
	NO ₂ OH OH	NH ₂ OH OH					
	\int						
	(3) () CH ₃	(4) $()$ CH_3					
	NH ₂ OH	$\gamma \gamma \gamma$ NO ₂ OH					
	2	2					

Ans. (2)

6

Sol.



LiAlH₄ will not affect C=C in this compound.
29. The major product of the following reaction is :-



Ans. (2)

Sol.



30. For the equilibrium,
2H₂O ⇒ H₃O⁺ + OH⁻, the value of ΔG° at 298 K is approximately :(1) -80 kJ mol⁻¹
(2) -100 kJ mol⁻¹
(3) 100 kI mol⁻¹
(4) 80 kI mol⁻¹

Ans. (4)

Sol.

 $2H_2O = H_3O^+ + OH^-$ K = 10⁻¹⁴ $\Delta G^\circ = -RT \ ln \ K$ $= \frac{-8.314}{2} \times 298 \times ln 10^{-14}$

$$= \frac{0.0011}{1000} \times 298 \times \ell n1$$
$$= 80 \text{ KJ/Mole}$$



TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On SATURDAY 12th JANUARY., 2019) TIME : 09 : 30 AM To 12 : 30 PM MATHEMATICS

- 1. For x >1, if $(2x)^{2y} = 4e^{2x-2y}$, then $(1 + \log_e 2x)^2 \frac{dy}{dx}$ is equal to : (1) $\log_{e} 2x$ (2) $\frac{x \log_e 2x + \log_e 2}{x}$ $(3) x \log_{e} 2x$ $(4) \ \frac{x \log_e 2x - \log_e 2}{x}$ Ans. (4) **Sol.** $(2x)^{2y} = 4e^{2x-2y}$ $2y\ell n2x = \ell n4 + 2x - 2y$ $y = \frac{x + \ell n 2}{1 + \ell n 2 x}$ $y' = \frac{(1 + \ell n 2x) - (x + \ell n 2)\frac{1}{x}}{(1 + \ell n 2x)^2}$ $y'(1+\ell n 2x)^2 = \left[\frac{x\ell n 2x - \ell n 2}{x}\right]$ The sum of the distinct real values of μ , for 2. which the vectors, $\mu \hat{i} + \hat{j} + \hat{k}$, $\hat{i} + \mu \hat{j} + \hat{k}$, $\hat{i} + \hat{j} + u\hat{k}$ are co-planer, is : (4) 1 (2) 0 (3) -1 (1) 2Ans. (3) **Sol.** $\begin{vmatrix} \mu & 1 & 1 \\ 1 & \mu & 1 \\ 1 & 1 \end{vmatrix} = 0$ $\mu(\mu^2 - 1) - 1(\mu - 1) + 1(1 - \mu) = 0$ $\mu^3 - \mu - \mu + 1 + 1 \ \mu = 0$ $\mu^3 - 3\mu + 2 = 0$ $\mu^3 - 1 - 3(\mu - 1) = 0$ $\mu = 1, \ \mu^2 + \mu - 2 = 0$ $\mu = 1, \ \mu = -2$ sum of distinct solutions = -1
- Let S be the set of all points in (-π,π) at which the function, f(x) = min {sinx, cosx} is not differentiable. Then S is a subset of which of the following?

(1)
$$\left\{-\frac{3\pi}{4}, -\frac{\pi}{4}, \frac{3\pi}{4}, \frac{\pi}{4}\right\}$$

(2) $\left\{-\frac{3\pi}{4}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{4}\right\}$
(3) $\left\{-\frac{\pi}{2}, -\frac{\pi}{4}, \frac{\pi}{4}, \frac{\pi}{2}\right\}$

$$(4) \left\{-\frac{\pi}{4}, 0, \frac{\pi}{4}\right\}$$

Ans. (1)



- 4. The product of three consecutive terms of a G.P. is 512. If 4 is added to each of the first and the second of these terms, the three terms now from an A.P. Then the sum of the original three terms of the given G.P. is
 - (1) 36 (2) 24
 - (3) 32 (4) 28

Ans. (4)

Sol. Let terms are $\frac{a}{r}$, $a, ar \rightarrow G.P$ $\therefore a^3 = 512 \Rightarrow a = 8$ $\frac{8}{r} + 4, 12, 8r \rightarrow A.P.$ $24 = \frac{8}{r} + 4 + 8r$ $r = 2, r = \frac{1}{2}$ r = 2 (4, 8, 16) $r = \frac{1}{2}(16, 8, 4)$ Sum = 28



JEE (Main) Examination-2019/Morning Session/12-01-2019

5.	The integral $\int \cos(\log_e x) dx$ is equal to :	7.	Let $S = \{1,2,3,, 100\}$. The number of non- empty subsets A of S such that the product of				
	(where C is a constant of integration)		elements in A is even is :-			-	
	-		$(1) \ 2^{50}(2^{50}-1) \qquad (2) \ 2^{100}-1$				
	(1) $\frac{x}{2}[\sin(\log_e x) - \cos(\log_e x)] + C$		(3) 2 ⁵⁰ -1		(4) 2 ⁵⁰ +1		
	-	Ans.	(1)				
	(2) $\frac{x}{2}[\cos(\log_e x) + \sin(\log_e x)] + C$	Sol.	$S = \{1, 2, 3\}$ = Total non	100} empty_subse	ets-subsets v	vith product	
	(3) $x[cos(log_e x) + sin(log_e x)] + C$		of element = 2^{100}	is odd $(1-1)(2^{50}-1)$		F	
	(4) $x[\cos(\log_e x) - \sin(\log_e x)] + C$		$= 2^{100} - 2^{50}(2)$	-2^{50}	I		
Ans.	(2)	o	$= 2^{30}(2$	of the devict	ions of 50 s	harmotiona	
Sol.	$I = \int \cos(\ell n x) dx$	0.	from 30	is 50, then	the mean	of these	
	$I = \cos(\ln x) \cdot x + \int \sin(\ell n x) dx$		(1) 50	(2) 51	(3) 30	(4) 31	
	$\cos(\ell n x)x + [\sin(\ell n x).x - \int \cos(\ell n x)dx]$	Ans.	(4)				
	$I = \frac{x}{2} [\sin(\ell n x) + \cos(\ell n x)] + C$	Sol.	$\sum_{i=1}^{30} (x_i - 30)$ $\Sigma x_i = 50 \times 3$	0 = 50 30 = 50			
6.	Let $S_k = \frac{1+2+3++k}{k}$. If		$\Sigma x_i = 50 + 1$ Mean = \overline{x}	$50 + 30$ $= \frac{\Sigma x_i}{n} = \frac{50 \times 10}{100}$	$\frac{30+50}{50} = 30$	+1 = 31	
	$S_1^2 + S_2^2 + \dots + S_{10}^2 = \frac{5}{12}A$, then A is equal to :	9.	If a variabl two circle	le line, $3x+4$ s $x^2 + y^2 -$	$y - \lambda = 0$ is su $2x - 2y + \lambda$	that the $1 = 0$ and	
Ang	(1) 303 (2) 283 (3) 156 (4) 301 (1)		$x^2+y^2-18x-2y+78 = 0$ are on its opposite side then the set of all values of λ is the interval				
Alls.			(1) [12, 21]	(2) (2, 17)	I	
Sol.	$S_{K} = \frac{K+1}{2}$		(3) (23, 31)	(4) [13, 23	3]	
	2	Ans.	(1)				
	$\Sigma S_k^2 = \frac{3}{12} A$	Sol.	Centre of o	circles are op	pposite side	of line	
	$\sum_{n=1}^{10} (K+1)^2 = 2^2 + 3^2 + + 11^2 = 5$		$(3+4-\lambda)$ $(\lambda - 7)$ (λ	(27 + 4 - 7) (27 + 4 - 7)	() < 0		
	$\sum_{K=1} \left(\frac{2}{2} \right) = \frac{4}{4} = \frac{12}{12} A$		$\lambda \in (7, 31)$)			
	$\frac{11 \times 12 \times 23}{1} - 1 = \frac{5}{2} A$		distance fr	om S ₁			
	6 3		$\left \frac{3+4-\lambda}{5}\right \ge 1 \implies \lambda \in (-\infty, 2] \cup [(12,\infty)]$ distance from S ₂				
	$505 = \frac{5}{3}A$, $A = 303$						





$$\left|\frac{27+4-\lambda}{5}\right| \ge 2 \implies \lambda \in (-\infty, 21] \cup [41, \infty)$$

so $\lambda \in [12, 21]$

10. A ratio of the 5^{th} term from the beginning to the 5^{th} term from the end in the binomial

expansion of
$$\left(2^{\frac{1}{3}} + \frac{1}{2(3)^{\frac{1}{3}}}\right)^{10}$$
 is :
(1) 1 : 4(16)^{\frac{1}{3}} (2) 1 : 2(6)^{\frac{1}{3}}
(3) 2(36)^{\frac{1}{3}} : 1 (4) 4(36)^{\frac{1}{3}} : 1
(4)

Ans. (4)

Sol.
$$\frac{T_5}{T_5^1} = \frac{{}^{10}C_4(2^{1/3})^{10-4} \left(\frac{1}{2(3)^{1/3}}\right)^4}{{}^{10}C_4\left(\frac{1}{2(3^{1/3})}\right)^{10-4}(2^{1/3})^4} = 4.(36)^{1/3}$$

11. let C_1 and C_2 be the centres of the circles $x^2+y^2-2x-2y-2 = 0$ and $x^2+y^2-6x-6y+14 = 0$ respectively. If P and Q are the points of intersection of these circles, then the area (in sq. units) of the quadrilateral PC₁QC₂ is : (1) 8 (2) 6 (3) 9 (4) 4

Ans. (4)



12. In a random experiment, a fair die is rolled until two fours are obtained in succession. The probability that the experiment will end in the fifth throw of the die is equal to :

(1) $\frac{150}{6^5}$ (2) $\frac{175}{6^5}$ (3) $\frac{200}{6^5}$ (4) $\frac{225}{6^5}$

Ans. (2) Sol.

$$\frac{1}{6^2} \left(\frac{5^3}{6^3} + \frac{2C_1 \cdot 5^2}{6^3} \right) = \frac{175}{6^5}$$

13. If the straight line, 2x-3y+17 = 0 is perpendicular to the line passing through the points (7, 17) and (15, β), then β equals :-

(1) -5
(2)
$$-\frac{35}{3}$$

(3) $\frac{35}{3}$
(4) 5
Ans. (4)
Sol. $\frac{17-\beta}{-8} \times \frac{2}{3} = -1$
 $\beta = 5$

14. Let f and g be continuous functions on [0, a] such that f(x) = f(a-x) and g(x)+g(a-x)=4,

then
$$\int_{0}^{a} f(x)g(x)dx$$
 is equal to :-

(1)
$$4\int_{0}^{a} f(x)dx$$
 (2) $2\int_{0}^{a} f(x)dx$
(3) $-3\int_{0}^{a} f(x)dx$ (4) $\int_{0}^{a} f(x)dx$

Ans. (2)

Sol.
$$I = \int_0^a f(x)g(x)dx$$
$$I = \int_0^a f(a-x)g(a-x)dx$$
$$I = \int_0^a f(x)(4-g(x)dx$$
$$I = 4\int_0^a f(x)dx - I$$
$$\Rightarrow I = 2\int_0^a f(x)dx$$

15. The maximum area (in sq. units) of a rectangle having its base on the x-axis and its other two vertices on the parabola, $y = 12-x^2$ such that the rectangle lies inside the parabola, is :-

(1) $20\sqrt{2}$ (2) $18\sqrt{3}$ (3) 32 (4) 36



JEE (Main) Examination-2019/Morning Session/12-01-2019

Ans. (3) Ans. (4) **Sol.** $f(a) = 2a(12 - a)^2$ **Sol.** $\tan^{-1}(2x) + \tan^{-1}(3x) = \pi/4$ $\Rightarrow \frac{5x}{1-6x^2} = 1$ $(a, 12 - a^2)$ $\Rightarrow 6x^2 + 5x - 1 = 0$ (0, 0)(a, 0) $f'(a) = 2(12 - 3a^2)$ maximum at a = 2maximum area = f(2) = 3219. 16. The Boolean expression $((p \land q) \lor (p \lor \sim q)) \land (\sim p \land \sim q)$ is equivalent to: (2) $p \lor (\sim q)$ (1) $p \land (\sim q)$ (3) $(\sim p) \land (\sim q)$ (4) $p \wedge q$ Ans. (3) (1)(1,-3) $\lim_{x \to \pi/4} \frac{\cot^3 x - \tan x}{\cos(x + \pi/4)}$ is : (3) (2, 4)17. Ans. (3) (1) 4 (2) $8\sqrt{2}$ (3) 8 (4) $4\sqrt{2}$ 1. Ans. (3) $\lim_{x \to \pi/4} \frac{\cot^3 x - \tan x}{\cos\left(x + \frac{\pi}{4}\right)}$ Sol. $\lim_{x\to\pi/4}\frac{(1-\tan^4 x)}{\cos(x+\pi/4)}$ 20. $2 \lim_{x \to \pi/4} \frac{(1 - \tan^2 x)}{\cos(x + \pi/4)}$ $\operatorname{R}_{x \to \pi/4} \frac{\cos^2 x - \sin^2 x}{\frac{\cos x - \sin x}{\sqrt{2}}} \frac{1}{\cos^2 x}$ Ans. (2) $4\sqrt{2}\lim_{x\to\pi/4}(\cos x + \sin x) = 8$ Sol. 18. Considering only the principal values of inverse functions, the set $A = \left\{ x \ge 0 : \tan^{-1}(2x) + \tan^{-1}(3x) = \frac{\pi}{4} \right\}$ (1) is an empty set (2) Contains more than two elements (3) Contains two elements (4) is a singleton

x = -1 or $x = \frac{1}{6}$ $x = \frac{1}{\epsilon}$ $\therefore x > 0$ An ordered pair(α,β) for which the system of linear equations $(1+\alpha)x + \beta y + z = 2$ $\alpha x + (1+\beta)y + z = 3$ $\alpha x + \beta y + 2z = 2$ has a unique solution is (2) (-3,1)(4) (-4, 2) Sol. For unique solution

$$\Delta \neq 0 \Longrightarrow \begin{vmatrix} 1+\alpha & \beta & 1 \\ \alpha & 1+\beta & 1 \\ \alpha & \beta & 2 \end{vmatrix} \neq 0$$

$$\begin{vmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ \alpha & \beta & 2 \end{vmatrix} \neq 0 \Longrightarrow \alpha + \beta \neq -2$$

The area (in sq. units) of the region bounded by the parabola, $y = x^2 + 2$ and the lines, y = x + 1, x = 0 and x = 3, is :

(4) $\frac{17}{4}$

(1)
$$\frac{15}{4}$$
 (2) $\frac{15}{2}$ (3) $\frac{21}{2}$



Req. area =
$$\int_{0}^{3} (x^2 + 2) dx - \frac{1}{2} \cdot 5 \cdot 3 = 9 + 6 - \frac{15}{2} = \frac{15}{2}$$

4

collegebatch.com

21. If λ be the ratio of the roots of the quadratic equation in x, $3m^2x^2+m(m-4)x+2 = 0$, then the

least value of m for which $\lambda + \frac{1}{\lambda} = 1$, is :

- (1) $2-\sqrt{3}$ (2) $4-3\sqrt{2}$
- (3) $-2 + \sqrt{2}$ (4) $4 2\sqrt{3}$

Ans. (2)

- **Sol.** $3m^2x^2 + m(m-4)x + 2 = 0$
 - $\lambda + \frac{1}{\lambda} = 1, \ \frac{\alpha}{\beta} + \frac{\beta}{\alpha} = 1, \ \alpha^2 + \beta^2 = \alpha\beta$ $(\alpha + \beta)^2 = 3\alpha\beta$ $\left(-\frac{m(m-4)}{3m^2}\right)^2 = \frac{3(2)}{3m^2}, \ \frac{(m-4)^2}{9m^2} = \frac{6}{3m}$ $(m-4)^2 = 18, \ m = 4 \pm \sqrt{18}, \ 4 \pm 3\sqrt{2}$
- 22. If the vertices of a hyperbola be at (-2, 0) and (2, 0) and one of its foci be at (-3, 0), then which one of the following points does not lie on this hyperbola?
 - (1) $(4,\sqrt{15})$ (2) $(-6,2\sqrt{10})$ (3) $(6,5\sqrt{2})$ (4) $(2\sqrt{6},5)$

Ans. (3)

Sol.
$$\frac{(-2,0)}{(-a,0)} \xrightarrow{(-2,0)} (a,0) \qquad (a,0) \qquad (ae = 3)$$
$$ae = 3, \ e = \frac{3}{2}, \ b^2 = 4\left(\frac{9}{4} - 1\right), \ b^2 = 5$$
$$\frac{x^2}{4} - \frac{y^2}{5} = 1$$

23. If $\frac{z-\alpha}{z+\alpha} (\alpha \in \mathbb{R})$ is a purely imaginary number and |z| = 2, then a value of α is :

(1) 1 (2) 2 (3) $\sqrt{2}$ (4) $\frac{1}{2}$

Ans. (2)

Е

Sol. $\frac{z-\alpha}{z+\alpha} + \frac{\overline{z}-\alpha}{\overline{z}+\alpha} = 0$ $z\overline{z} + z\alpha - \alpha\overline{z} - \alpha^2 + z\overline{z} - z\alpha + \overline{z}\alpha - \alpha^2 = 0$ $|z|^2 = \alpha^2, \quad a = \pm 2$ 24. Let P(4, -4) and Q(9, 6) be two points on the parabola, $y^2= 4x$ and let X be any point on the arc POQ of this parabola, where O is the vertex of this parabola, such that the area of ΔPXQ is maximum. Then this maximum area (in sq. units) is :



25. the perpendicular distance from the origin to the plane containing the two lines, $\frac{x+2}{3} = \frac{y-2}{5} = \frac{z+5}{7}$ and $\frac{x-1}{1} = \frac{y-4}{4} = \frac{z+4}{7}$, is: (1) $\frac{11}{\sqrt{6}}$ (2) $6\sqrt{11}$ (3) 11 (4) $11\sqrt{6}$ Ans. (1) Sol. $\begin{vmatrix} i & j & k \\ 3 & 5 & 7 \end{vmatrix}$

$$|1 \quad 4 \quad 7|$$

$$\hat{i}(35-28) - \hat{j}(21.7) + \hat{k}(12-5)$$

$$7\hat{i} - 14\hat{j} + 7\hat{k}$$

$$\hat{i} - 2\hat{j} + \hat{k}$$

$$1(x + 2) - 2(y - 2) + 1 (z + 15) = 0$$

$$x - 2y + z + 11 = 0$$

$$\frac{11}{\sqrt{4+1+1}} = \frac{11}{\sqrt{6}}$$



JEE (Main) Examination-2019/Morning Session/12-01-2019

The maximum value of $3\cos\theta + 5\sin\left(\theta - \frac{\pi}{6}\right)$ for \int **Sol.** $\frac{dy}{dx} = \frac{y}{x} = \ell n x$ 26. any real value of θ is : any real value of θ is : (1) $\sqrt{19}$ (2) $\frac{\sqrt{79}}{2}$ (3) $\sqrt{31}$ (4) $\sqrt{34}$ $e^{\int \frac{1}{x} dx} = x$ $xy = \int x \ell nx + C$ Ans. (1) $\ell n x \frac{x^2}{2} - \int \frac{1}{x} \cdot \frac{x^2}{2}$ **Sol.** $y = 3\cos\theta + 5\left(\sin\theta\frac{\sqrt{3}}{2} - \cos\theta\frac{1}{2}\right)$ $xy = \frac{x}{2} \ell n x - \frac{x^2}{4} + C$, for $2y(2) = 2\ell n 2 - 1$ $\frac{5\sqrt{3}}{2}\sin\theta + \frac{1}{2}\cos\theta$ $\Rightarrow C = 0$ $y_{max} = \sqrt{\frac{75}{4} + \frac{1}{4}} = \sqrt{19}$ $y = \frac{x}{2} \ell n x - \frac{x}{4}$ 27. A tetrahedron has vertices P(1, 2, 1), $y(e) = \frac{e}{4}$ Q(2, 1, 3), R(-1,1,2) and O(0, 0, 0). The angle between the faces OPQ and PQR is : **29.** Let P = $\begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 9 & 3 & 1 \end{bmatrix}$ and Q = [q_{ij}] be two 3×3 (1) $\cos^{-1}\left(\frac{9}{35}\right)$ (2) $\cos^{-1}\left(\frac{19}{35}\right)$ (3) $\cos^{-1}\left(\frac{17}{21}\right)$ (4) $\cos^{-1}\left(\frac{7}{21}\right)$ matrices such that $Q-P^5 = I_3$. Then $\frac{q_{21}+q_{31}}{q_{32}}$ is Ans. (1) equal to: **Sol.** $\overrightarrow{OP} \times \overrightarrow{OQ} = (\hat{i} + 2\hat{j} + \hat{k}) \times (2\hat{i} + \hat{j} + 3\hat{k})$ (1) 15 (2) 9 (3) 135 (4) 10 $5\hat{i}-\hat{i}-3\hat{k}$ Ans. (4) O(0,0,0)**Sol.** $P = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 9 & 3 & 1 \end{bmatrix}$ **>**R (−1,1,2) (1,2,1) $\mathbf{P}^2 = \begin{vmatrix} 1 & 0 & 0 \\ 3+3 & 1 & 0 \\ 9+9+9 & 3+3 & 1 \end{vmatrix}$ (2.13) $\overrightarrow{PQ} \times \overrightarrow{PR} = (\hat{i} - \hat{j} + 2\hat{k}) \times (-2\hat{i} - \hat{j} + \hat{k})$ $P^{3} = \begin{bmatrix} 1 & 0 & 0 \\ 3+3+3 & 1 & 0 \\ 6 & 3+3+3 & 1 \end{bmatrix}$ $\hat{i} - 5\hat{j} - 3\hat{k}$ $\cos\theta = \frac{5+5+9}{\left(\sqrt{25+9+1}\right)^2} = \frac{19}{35}$ $P^{n} = \begin{vmatrix} 1 & 0 & 0 \\ 3n & 1 & 0 \\ \frac{n(n+1)}{2}3^{2} & 3n & 1 \end{vmatrix}$ Let y = y(x) be the solution of the differential 28. equation, $x\frac{dy}{dx} + y = x \log_e x, (x > 1)$. If $2y(2) = \log_e 4 - 1$, then y(e) is equal to :- $\mathbf{P}^{5} = \begin{vmatrix} 1 & 0 & 0 \\ 5.3 & 1 & 0 \\ 15.9 & 5.3 & 1 \end{vmatrix}$ (1) $\frac{e^2}{4}$ (2) $\frac{e}{4}$ (3) $-\frac{e}{2}$ (4) $-\frac{e^2}{2}$ Ans. (2)

6

collegebatch.com

 $Q = P^5 + I_3$ $\mathbf{Q} = \begin{bmatrix} 2 & 0 & 0\\ 15 & 2 & 0\\ 135 & 15 & 2 \end{bmatrix}$ $\frac{\mathbf{q}_{21} + \mathbf{q}_{31}}{\mathbf{q}_{22}} = \frac{15 + 135}{15} = 10$ q_{32} Aliter $\mathbf{P} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 3 & 0 & 0 \\ 9 & 3 & 0 \end{pmatrix}$ $\mathbf{P} = \mathbf{I} + \mathbf{X}$ $\mathbf{X} = \begin{pmatrix} 0 & 0 & 0 \\ 3 & 0 & 0 \\ 9 & 3 & 0 \end{pmatrix}$ $\mathbf{X}^2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 9 & 0 & 0 \end{pmatrix}$ $X^{3} = 0$ $P^5 = I + 5X + 10X^2$ $Q = P^5 + I = 2I + 5X + 10X^2$ $\mathbf{Q} = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 15 & 0 & 0 \\ 15 & 15 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 90 & 0 & 0 \end{pmatrix}$ $\Rightarrow Q = \begin{pmatrix} 2 & 0 & 0 \\ 15 & 2 & 0 \\ 135 & 15 & 2 \end{pmatrix}$

30. Consider three boxes, each containing 10 balls labelled 1,2,...,10. Suppose one ball is randomly drawn from each of the boxes. Denote by n_i , the label of the ball drawn from the ith box, (i = 1, 2, 3). Then, the number of ways in which the balls can be chosen such that $n_1 < n_2 < n_3$ is : (1) 82 (2) 240 (3) 164 (4) 120

Ans. (4)

Sol. No. of ways = $10C_3 = 120$

7



TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Saturday 12th JANUARY, 2019) TIME : 09 : 30 AM To 12 : 30 PM PHYSICS

1. Two light identical springs of spring constant k are attached horizontally at the two ends of a uniform horizontal rod AB of length ℓ and mass m. The rod is pivoted at its centre 'O' and can rotate freely in horizontal plane. The other ends of the two springs are fixed to rigid supports as shown in figure. The rod is gently pushed through a small angle and released. The frequency of resulting oscillation is:







$$\tau = -2Kx \frac{\ell}{2} \cos \theta$$

$$\Rightarrow \tau = \left(\frac{K\ell^2}{2}\right)\theta = -C\theta$$

$$\Rightarrow f = \frac{1}{2\pi} \sqrt{\frac{C}{I}} = \frac{1}{2\pi} \sqrt{\frac{\frac{K\ell^2}{2}}{\frac{M\ell^2}{12}}}$$

$$\Rightarrow f = \frac{1}{2\pi} \sqrt{\frac{6K}{M}}$$

2. A cylinder of radius R is surrounded by a cylindrical shell of inner radius R and outer radius 2R. The thermal conductivity of the material of the inner cylinder is K_1 and that of the outer cylinder is K_2 . Assuming no loss of heat, the effective thermal conductivity of the system for heat flowing along the length of the cylinder is:

(1)
$$K_1 + K_2$$

(2) $\frac{K_1 + K_2}{2}$
(3) $\frac{2K_1 + 3K_2}{5}$
(4) $\frac{K_1 + 3K_2}{4}$
Ans. (4)



$$= \frac{K_1 + 3K_2}{4}$$

3. A travelling harmonic wave is represented by the equation $y(x, t) = 10^{-3} \sin (50 t + 2x)$, where x and y are in meter and t is in seconds. Which of the following is a correct statement about the wave?

The wave is propagating along the

- (1) negative x-axis with speed $25ms^{-1}$
- (2) The wave is propagating along the positive x-axis with speed 25 ms^{-1}
- (3) The wave is propagating along the positive x-axis with speed 100 ms⁻¹
- (4) The wave is propagating along the negative x-axis with speed 100 ms⁻¹

1



JEE (Main) Examination-2019/Morning Session/12-01-2019

Sol. $y=a \sin(\omega t + kx)$ \Rightarrow wave is moving along -ve x-axis with speed

$$v = \frac{\omega}{K} \Longrightarrow v = \frac{50}{2} = 25 \text{ m/sec.}$$

4. A straight rod of length L extends from x = a to x=L + a. The gravitational force is exerts on a point mass 'm' at x = 0, if the mass per unit length of the rod is A + Bx², is given by:

(1)
$$\operatorname{Gm}\left[A\left(\frac{1}{a+L}-\frac{1}{a}\right)-BL\right]$$

(2) $\operatorname{Gm}\left[A\left(\frac{1}{a}-\frac{1}{a+L}\right)+BL\right]$
(3) $\operatorname{Gm}\left[A\left(\frac{1}{a+L}-\frac{1}{a}\right)+BL\right]$
(4) $\operatorname{Gm}\left[A\left(\frac{1}{a}-\frac{1}{a+L}\right)-BL\right]$

Ans. (2)



dm= (A + Bx²)dx
dF =
$$\frac{GM dm}{x^2}$$

= F = $\int_a^{a+L} \frac{GM}{x^2} (A + Bx^2) dx$
= $GM \left[-\frac{A}{x} + Bx \right]_a^{a+L}$
= $GM \left[A \left(\frac{1}{a} - \frac{1}{a+L} \right) + BL \right]$

5. A light wave is incident normally on a glass slab of refractive index 1.5. If 4% of light gets reflected and the amplitude of the electric field of the incident light is 30V/m, then the amplitude of the electric field for the wave propogating in the glass medium will be:

Ans. (2)

Sol.
$$P_{refracted} = \frac{96}{100} P_{I}$$
$$\Rightarrow K_{2}A_{t}^{2} = \frac{96}{100}K_{1}A_{i}^{2}$$
$$\Rightarrow r_{2}A_{t}^{2} = \frac{96}{100}r_{1}A_{i}^{2}$$
$$\Rightarrow A_{t}^{2} = \frac{96}{100} \times \frac{1}{\frac{3}{2}} \times (30)^{2}$$
$$A_{t4}\sqrt{\frac{64}{100}} \times (30)^{2} = 24$$

6. The output of the given logic circuit is :




7. In the figure shown, after the switch 'S' is turned from position 'A' to position 'B', the energy dissipated in the circuit in terms of capacitance 'C' and total charge 'Q' is:



(1)
$$\frac{3}{8} \frac{Q^2}{C}$$
 (2) $\frac{3}{4} \frac{Q^2}{C}$ (3) $\frac{1}{8} \frac{Q^2}{C}$ (4) $\frac{5}{8} \frac{Q^2}{C}$

Ans. (1)

Sol. $V_i = \frac{1}{2}CE^2$

$$V_{f} = \frac{\left(CE\right)^{2}}{2 \times 4c} = \frac{1}{2} \frac{CE^{2}}{4}$$
$$\Delta E = \frac{1}{2} CE^{2} \times \frac{3}{4} = \frac{3}{8} CE$$

8. A particle of mass m moves in a circular orbit

in a central potential field $U(r) = \frac{1}{2}kr^2$. If Bohr's quantization conditions are applied, radii of possible orbitals and energy levels vary with quantum number n as:

(1) $r_n \propto n^2$, $E_n \propto \frac{1}{n^2}$ (2) $r_n \propto \sqrt{n}$, $E_n \propto \frac{1}{n}$ (3) $r_n \propto n$, $E_n \propto n$ (4) $r_n \propto \sqrt{n}$, $E_n \propto n$ **Ans. (4)**

Sol. $F = \frac{dV}{dr} = kr = \frac{mv^2}{r}$ $mvr = \frac{nh}{2\pi}$

$$r^2 \propto n$$

 $r^{2} \propto \sqrt{n}$ $E = \frac{1}{2}kr^{2} + \frac{1}{2}mv^{2} \propto r^{2}$ $\propto n$

9. Two electric bulbs, rated at (25 W, 220 V) and (100 W, 220 V), are connected in series across a 220 V voltage source. If the 25 W and 100 W bulbs draw powers P_1 and P_2 respectively, then:

(1)
$$P1 = 9 W$$
, $P_2 = 16 W$
(2) $P_2 = 4 W P_2 = 16W$

(3)
$$P_1 = 16 \text{ W}, P_2 = 4 \text{W}$$

(4)
$$P_1$$
 16 W, $P_2 = 9W$

Ans. (3)

Sol.
$$R_1 = \frac{220^2}{25}$$

 $R_2 = \frac{220^2}{100}$
 $L = \frac{220}{R_1 + R_2}$
 $P_1 = i^2 R_1$
 $P_2 = i^2 (R_2 = 4W)$
 $= \frac{220^2}{\left(\frac{220^2}{25} + \frac{220^2}{100}\right)} \times \frac{220^2}{25}$

$$=\frac{400}{25}=16W$$

10. A satellite of mass M is in a circular orbit of radius R about the centre of the earth. A meteorite of the same mass, falling towards the earth, collides with the satellite completely inelastically. The speeds of the satellite and the meteorite are the same, just before the collision. The subsequent motion of the combined body will be :

collegebatch.com

Ε

- (1) in a circular orbit of a different radius
- (2) in the same circular orbit of radius R
- (3) in an elliptical orbit
- (4) such that it escapes to infinity
- Ans. (3)

Sol.
$$mv\hat{i} + mv\hat{j}$$

 $=2m\vec{v}^{1}$

°

 $\vec{v} = \frac{1}{\sqrt{2}} \times \sqrt{\frac{GM}{R}}$

- 11. Let the moment of inertia of a hollow cylinder of length 30 cm (inner radius 10 cm and outer radius 20 cm), about its axis be I. The radius of a thin cylinder of the same mass such that its moment of inertia about its axis is also I, is: (1) 12 cm (2) 18 cm (3) 16 cm (4) 14 cm
 Ans. (3)
- 12. A passenger train of length 60m travels at a speed of 80 km/hr. Another freight train of length 120 m travels at a speed of 30 km/hr. The ratio of times taken by the passenger train to completely cross the freight train when : (i) they are moving in the same direction, and (ii) in the opposite directions is :

(1)
$$\frac{5}{2}$$
 (2) $\frac{25}{11}$ (3) $\frac{3}{2}$ (4) $\frac{11}{5}$

Ans. (4)

- 13. An ideal gas occupies a volume of $2m^3$ at a pressure of 3×10^6 Pa. The energy of the gas is:
 - (1) 3×10^2 (2) 10^8 J (3) $6 \times 10^4 \text{ J}$ (4) $9 \times 10^6 \text{ J}$

Ans. (4)

Sol. Energy =
$$\frac{1}{2}$$
nRT = $\frac{f}{2}$ PV

$$=\frac{1}{2}(3\times10^6)(2)$$

 $= f \times 3 \times 10^6$

Considering gas is monoatomic i.e. f = 3

$$E_{.} = 9 \times 10^{6} J_{.}$$

Option-(4)

14. A 100 V carrier wave is made to vary between 160 V and 40 V by a modulating signal. What is the modulation index?

- Sol. $E_m + E_c = 160$ $E_m + 100 = 160$ $E_m = 60$ $\mu = \frac{E_m}{E_c} = \frac{60}{100}$ $\mu = 0.6$
- 15. The galvanometer deflection, when key K_1 is closed but K_2 is open, equals θ_0 (see figure). On closing K_2 also and adjusting R_2 to 5 Ω , the

deflection in galvanometer becomes
$$\frac{\theta_0}{5}$$
. The

resistance of the galvanometer is, then, given by [Neglect the internal resistance of battery]:



	(1) 12Ω	(2) 25Ω	(3) 5Ω	(4) 22Ω
Ans.	(4)			

4



Sol. case I

Case II

$$i_{g} = \left(\frac{E}{220 + \frac{5R_{g}}{5 + R_{g}}}\right) \times \frac{5}{\left(R_{g} + 5\right)} = \frac{C\theta_{0}}{5} \quad ..(ii)$$

 $i_g = \frac{E}{220 + R_g} = C\theta_0$

..(i)

$$\Rightarrow \frac{5E}{225R_a + 1100} = \frac{C\theta_0}{5} \qquad ..(ii)$$

$$\frac{E}{220 + R_g} = C\theta \qquad \dots (i)$$

$$\Rightarrow \frac{225R_g + 1100}{1100 + 5R_g} = 5$$

$$\Rightarrow 5500 + 25R_g = 225R_g + 1100$$

$$200R_g = 4400$$

$$R_g = 22\Omega$$

Ans. - 4

16. A person standing on an open ground hears the sound of a jet aeroplane, coming from north at an angle 60° with ground level. But he finds the aeroplane right vertically above his position. If υ is the speed of sound, speed of the plane is :

(4) $\frac{\sqrt{3}}{2}v$

(1) $\frac{2\upsilon}{\sqrt{3}}$ (2) υ

(3)
$$\frac{v}{2}$$

Ans. (3)



$$AB = V_{p} \times t$$
$$BC = Vt$$
$$\cos 60^{\circ} = \frac{AB}{BC}$$
$$\frac{1}{2} = \frac{V_{p} \times t}{Vt}$$
$$V_{p} = -\frac{V}{2}$$

17. A proton and an α -particle (with their masses in the ratio of 1:4 and charges in the ratio of 1:2) are accelerated from rest through a potential difference V. If a uniform magnetic field (B) is set up perpendicular to their velocities, the ratio of the radii $r_p : r_\alpha$ of the circular paths described by them will be :

(1)
$$1:\sqrt{2}$$
 (2) $1:2$ (3) $1:3$ (4) $1:\sqrt{3}$

Ans. (1)
Sol.
$$KE = q\Delta V$$

$$r = \frac{\sqrt{2mq\Delta V}}{qB}$$

γq

$$\frac{r_{\rm p}}{r_{\rm \infty}} = \frac{1}{\sqrt{2}}$$

18. A point source of light, S is placed at a distance L in front of the centre of plane mirror of width d which is hanging vertically on a wall. A man walks in front of the mirror along a line parallel to the mirror, at a distance 2L as shown below. The distance over which the man can see the image of the light source in the mirror is :





	(1) 3d	(2) $\frac{d}{2}$
	(3) d	(4) 2d
Ans.	(1)	



3d

19. The least count of the main scale of a screw gauge is 1 mm. The minimum number of divisions on its circular scale required to measure 5µm diameter of wire is :

(1)	50	(2)	100
(3)	200	(4)	500

Ans. (3)

Sol. Least count = $\frac{\text{Pitch}}{\text{Number of division on circular scale}}$

 $5 \times 10^{-6} = \frac{10^{-3}}{N}$ N = 200

20. A simple pendulum, made of a string of length l and a bob of mass m, is released from a small angle θ_0 . It strikes a block of mass M, kept on a horizontal surface at its lowest point of oscillations, elastically. It bounces back and goes up to an angle θ_1 . Then M is given by :

(1)
$$\frac{\mathrm{m}}{2} \left(\frac{\theta_0 - \theta_1}{\theta_0 + \theta_1} \right)$$
 (2) $\frac{\mathrm{m}}{2} \left(\frac{\theta_0 + \theta_1}{\theta_0 - \theta_1} \right)$
(3) $\mathrm{m} \left(\frac{\theta_0 + \theta_1}{\theta_0 - \theta_1} \right)$ (4) $\mathrm{m} \left(\frac{\theta_0 - \theta_1}{\theta_0 + \theta_1} \right)$

Ans. (3)

6

$$\begin{split} & \overbrace{\mathbf{m}}^{\mathbf{h}} \underbrace{\mathbf{h}}_{\mathbf{m}} \\ & \text{Before colision} \\ & \overbrace{\mathbf{m}}^{\mathbf{m}} \underbrace{\mathbf{M}}_{\mathbf{v}_{1}} \underbrace{\mathbf{M}}_{\mathbf{v}_{1}} \underbrace{\mathbf{M}}_{\mathbf{m}} \underbrace{\mathbf{M}}_{\mathbf{v}_{m}} \\ & \mathbf{v} = \sqrt{2g\ell(1 - \cos\theta_{0})} \\ & \mathbf{v}_{1} = \sqrt{2g\ell(1 - \cos\theta_{1})} \\ & \text{By momentum conservation} \\ & \mathbf{m}\sqrt{2g\ell(1 - \cos\theta_{0})} = \mathbf{M}\mathbf{V}_{\mathbf{m}} - \mathbf{m}\sqrt{2gl(1 - \cos\theta_{1})} \\ & \Rightarrow \mathbf{m}\sqrt{2g\ell}\left\{\sqrt{1 - \cos\theta_{0}} + \sqrt{1 - \cos\theta_{1}}\right\} = \mathbf{M}\mathbf{V}_{\mathbf{m}} \\ & \text{and} \quad \mathbf{e} = 1 = \frac{\mathbf{V}_{\mathbf{m}} + \sqrt{2g\ell(1 - \cos\theta_{1})}}{\sqrt{2g\ell(1 - \cos\theta_{0})}} \\ & \sqrt{2g\ell}\left(\sqrt{1 - \cos\theta_{0}} - \sqrt{1 - \cos\theta_{1}}\right) = \mathbf{V}_{\mathbf{m}} \\ & \dots(\mathbf{I}) \\ & \mathbf{m}\sqrt{2g\ell}\left(\sqrt{1 - \cos\theta_{0}} + \sqrt{1 - \cos\theta_{1}}\right) = \mathbf{M}\mathbf{V}_{\mathbf{M}} \\ & \dots(\mathbf{I}) \\ & \text{Dividing} \\ & \frac{\left(\sqrt{1 - \cos\theta_{0}} + \sqrt{1 - \cos\theta_{1}}\right)}{\left(\sqrt{1 - \cos\theta_{0}} - \sqrt{1 - \cos\theta_{1}}\right)} = \frac{\mathbf{M}}{\mathbf{m}} \end{split}$$

Λ

Sol.

By componendo divided

$$\frac{m-M}{m+M} = \frac{\sqrt{1-\cos\theta_1}}{\sqrt{1-\cos\theta_0}} = \frac{\sin\left(\frac{\theta_1}{2}\right)}{\sin\left(\frac{\theta_0}{2}\right)}$$

$$\Rightarrow \frac{\mathbf{M}}{\mathbf{m}} = \frac{\theta_0 - \theta_1}{\theta_0 + \theta_1} \Rightarrow \mathbf{M} = \frac{\theta_0 - \theta_1}{\theta_0 + \theta_1}$$

Collegebatch.com

21. What is the position and nature of image formed by lens combination shown in figure? $(f_1, f_2 \text{ are focal lengths})$



- (1) 70 cm from point B at left; virtual
- (2) 40 cm from point B at right; real
- (3) $\frac{20}{3}$ cm from point B at right, real
- (4) 70 cm from point B at right, real

Ans. (4)

Sol. For first lens

 $\frac{1}{V} - \frac{1}{-20} = \frac{1}{5}$

$$V = \frac{20}{3}$$

For second lens

$$V = \frac{20}{3} - 2 = \frac{14}{3}$$
$$\frac{1}{V} - \frac{1}{\frac{14}{3}} = \frac{1}{-5}$$

V = 70cm

22. In the figure shown, a circuit contains two identical resistors with resistance $R = 5\Omega$ and an inductance with L = 2mH. An ideal battery of 15 V is connected in the circuit. What will be the current through the battery long after the switch is closed?



(1) 6A	(2) 7.5A
(3) 5.5A	(4) 3A

Ans. (1)

Sol. Ideal inductor will behave like zero resistance long time after switch is closed



23. Determine the electric dipole moment of the system of three charges, placed on the vertices of an equilateral triangle, as shown in the figure:



Ans. (3)



$$|\mathbf{P}_1| = \mathbf{q}(\mathbf{d})$$

$$|\mathbf{P}_2| = \mathbf{q}\mathbf{d}$$

|Resultant| = 2 P cos30°

$$2 \operatorname{qd}\left(\frac{\sqrt{3}}{2}\right) = \sqrt{3} \operatorname{qd}$$

24. The position vector of the centre of mass \overrightarrow{r} cm of an symmetric uniform bar of negligible area of cross-section as shown in figure is :



Ans. (1)

I

Sol.
$$2m (L,L) = m\left(2L,\frac{L}{2}\right) + \left(\frac{5L}{2}, 0\right)$$

$$X_{cm} = \frac{2mL + 2mL + \frac{5mL}{2}}{4m} = \frac{13}{8}L$$
$$Y_{cm} = \frac{2m \times L + m \times \left(\frac{L}{2}\right) + m \times 0}{4m} = \frac{5L}{8}$$

25. As shown in the figure, two infinitely long, identical wires are bent by 90° and placed in such a way that the segments LP and QM are along the x-axis, while segments PS and QN are parallel to the y-axis. If OP = OQ = 4 cm, and the magnitude of the magnetic field at O is 10⁻⁴ T, and the two wires carry equal currents (see figure), the magnitude of the current in each wire and the direction of the magnetic field at O will be ($\mu_0 = 4\pi \times 10^{-7}$ NA⁻²) :



- (1) 40 A, perpendicular into the page
- (2) 40 A, perpendicular out of the page
- (3) 20 A, perpendicular out of the page
- (4) 20 A, perpendicular into the page

Ans. (4)

Sol. Magnetic field at 'O' will be done to 'PS' and 'QN' only

i.e. $B_0 = B_{PS} + B_{QN} \rightarrow Both$ inwards Let current in each wire = i

$$\therefore \qquad B_0 = \frac{\mu_0 i}{4\pi d} + \frac{\mu_0 i}{4\pi d}$$

or
$$10^{-4} = \frac{\mu_0 i}{2\pi d} = \frac{2 \times 10^{-7} \times i}{4 \times 10^{-2}}$$

$$\therefore \qquad i = 20 \text{ A}$$

26. In a meter bridge, the wire of length 1 m has a non-uniform cross-section such that, the

variation
$$\frac{dR}{d\ell}$$
 of its resistance R with length ℓ

is $\frac{dR}{d\ell} \propto \frac{1}{\sqrt{\ell}}$. Two equal resistances are

connected as shown in the figure. The galvanometer has zero deflection when the jockey is at point P. What is the length AP?





collegebatch.com

Sol. For the given wire : $dR = C \frac{d\ell}{\sqrt{\ell}}$, where C =

constant.

Let resistance of part AP is R_1 and PB is R_2

 $\frac{\mathbf{R'}}{\mathbf{R'}} = \frac{\mathbf{R_1}}{\mathbf{R_2}}$ or $\mathbf{R_1} = \mathbf{R_2}$ By balanced *.*..

WSB concept.

Now
$$\int d\mathbf{R} = c \int \frac{d\ell}{\sqrt{\ell}}$$

$$\therefore \mathbf{R}_1 = \mathbf{C} \int_0^\ell \ell^{-1/2} d\ell = \mathbf{C} \cdot 2 \cdot \sqrt{\ell}$$

$$\mathbf{R}_2 = \mathbf{C} \int_\ell^1 \ell^{-1/2} d\ell = \mathbf{C} \cdot (2 - 2\sqrt{\ell})$$

Putting $\mathbf{R}_1 = \mathbf{R}_2$

$$\mathbf{C} 2 \sqrt{\ell} = \mathbf{C} \cdot (2 - 2\sqrt{\ell})$$

$$C2\sqrt{\ell} = 0$$

$$\therefore 2\sqrt{\ell} = 1$$

$$\sqrt{\ell} = \frac{1}{2}$$
i.e. $\ell = \frac{1}{4}$ m $\Rightarrow 0.25$ m

27. For the given cyclic process CAB as shown for a gas, the work done is :



Ans. (3)

Е

Sol. Since P-V indicator diagram is given, so work done by gas is area under the cyclic diagram.

$$\therefore \Delta W = \text{Work done by gas} = \frac{1}{2} \times 4 \times 5 \text{ J}$$
$$= 10 \text{ J}$$

An ideal battery of 4 V and resistance R are 28. connected in series in the primary circuit of a potentiometer of length 1 m and resistance 5Ω . The value of R, to give a potential difference of 5 mV across 10 cm of potentiometer wire, is :

(1)
$$490 \Omega$$
 (2) 480Ω
(3) 395Ω (4) 495Ω

Ans. (3)

Sol.
$$\begin{array}{c}
4v & R \\
i & 5\Omega & i \\
1m
\end{array}$$

...

Let current flowing in the wire is i.

$$i = \left(\frac{4}{R+5}\right)A$$

If resistance of 10 m length of wire is x

then x = 0.5
$$\Omega$$
 = 5 × $\frac{0.1}{1}\Omega$
 $\therefore \quad \Delta V = P. d. on wire = i. x$
 $5 \times 10^{-3} = \left(\frac{4}{R+5}\right) \cdot (0.5)$
 $\therefore \quad \frac{4}{R+5} = 10^{-2} \text{ or } R+5 = 400 \Omega$
 $\therefore R = 395 \Omega$

29. A particle A of mass 'm' and charge 'q' is accelerated by a potential difference of 50 V. Another particle B of mass '4 m' and charge 'q' is accelerated by a potential difference of 2500

V. The ratio of de-Broglie wavelengths
$$\frac{\lambda_A}{\lambda_B}$$
 is

close to :

(1) 10.00(2) 14.14 (3) 4.47 (4) 0.07

Ans. (2)

Sol. K.E. acquired by charge = K = qV

$$\lambda = \frac{h}{p} = \frac{h}{\sqrt{2mK}} = \frac{h}{\sqrt{2mqV}}$$
$$\therefore \quad \frac{\lambda_A}{\lambda_B} = \frac{\sqrt{2m_B q_B V_B}}{\sqrt{2m_A q_A V_A}} = \sqrt{\frac{4m.q.2500}{m.q.50}} = 2\sqrt{50}$$
$$= 2 \times 7.07 = 14.14$$



30. There is a uniform spherically symmetric surface charge density at a distance R_0 from the origin. The charge distribution is initially at rest and starts expanding because of mutual repulsion. The figure that represents best the speed V(R(t)) of the distribution as a function of its instantaneous radius R (t) is :



Ans. (1)

Sol. At any instant 't' Total energy of charge distribution is constant

i.e.
$$\frac{1}{2}mV^{2} + \frac{KQ^{2}}{2R} = 0 + \frac{KQ^{2}}{2R_{0}}$$
$$\therefore \quad \frac{1}{2}mV^{2} = \frac{KQ^{2}}{2R_{0}} - \frac{KQ^{2}}{2R}$$
$$\therefore \quad V = \sqrt{\frac{2}{m}\frac{KQ^{2}}{2} \cdot \left(\frac{1}{R_{0}} - \frac{1}{R}\right)}$$
$$\therefore \quad V = \sqrt{\frac{KQ^{2}}{m}\left(\frac{1}{R_{0}} - \frac{1}{R}\right)} = C\sqrt{\frac{1}{R_{0}} - \frac{1}{R}}$$

Also the slope of v-s curve will go on decreasing

 \therefore Graph is correctly shown by option(1)



TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Saturday 12th JANUARY, 2019) TIME : 09 : 30 AM To 12 : 30 PM CHEMISTRY

Iodine reacts with concentrated HNO₃ to yield Y along with other products. The oxidation state of iodine in Y, is :
 (1) 5
 (2) 3
 (3) 1
 (4) 7

(1) 5 Ans. (1)

- Sol. $I_2 + 10HNO_3 \longrightarrow 2HIO_3 + 10NO_2 + 4H_2O$ In HIO₃ oxidation state of iodine is +5.
- 2. The major product of the following reaction is:







Ans. (3)



DIBAL-H will reduce cyanides & esters to aldehydes.

In a chemical reaction, A + 2B → 2C + D, the initial concentration of B was 1.5 times of the concentration of A, but the equilibrium concentrations of A and B were found to be equal. The equilibrium constant(K) for the aforesaid chemical reaction is :

(3) 1

(1) 16 (2) 4

Ans.(2)

Ε

6.1	$A \rightarrow 2P \rightarrow 2C \rightarrow D$		
501.	A + 2B = 2C + D		
	$t=0$ a_0 $1.5a_0$ 0 0		
	$t = t_{eq} a_0 - x 1.5a_0 - 2x 2x x$		
	At equilibrium [A] = [B] $a_0 - x = 1.5a_0 - 2x \implies x = 0.5a_0$ $t = t_{eq} = 0.5a_0 = 0.5a_0 = 0.5a_0$ $K_C = \frac{[C]^2[D]}{[A][B]^2} = \frac{(a_0)^2 (0.5a_0)}{(0.5a_0) (0.5a_0)^2} = 4$		
4.	Two solids dissociate as follows $A(s) \iff B(g) + C(g) ; K_{p_1} = x \text{ atm}^2$ $D(s) \iff C(g) + E(g) ; K_{p_2} = y \text{ atm}^2$ The total pressure when both the solids dissociat simultaneously is :-		
	(1) $x^2 + y^2$ atm (2) $x^2 + y^2$ atm		
	(3) $2(\sqrt{x+y})$ atm (4) $\sqrt{x+y}$ atm		
Ans.	(3)		
Sol.	$A(s) \Longrightarrow B(g) + C(g)$ $K_{p} = x = P_{p} \cdot P_{C}$		
	P_1 P_1 $x=P_1(P_1 + P_2)$ (1)		
	$D(c) \rightarrow C(\alpha) + F(\alpha) K = v = P P$		

$$P_{1} P_{1} x = P_{1}(P_{1} + P_{2}) \dots(1)$$

$$D(s) \rightleftharpoons C(g) + E(g) K_{P_{2}} = y = P_{C} \cdot P_{E}$$

$$P_{2} P_{2} y = (P_{1} + P_{2}) (P_{2}) \dots(2)$$
Adding (1) and (2)
$$x + y = (P_{1} + P_{2})^{2}$$
Now total pressure
$$P_{T} = P_{C} + P_{B} + P_{E}$$

$$= (P_{1} + P_{2}) + P_{1} + P_{2} = 2(P_{1} + P_{2})$$

$$P_{T} = 2(\sqrt{x + y})$$

- 5. Freezing point of a 4% aqueous solution of X is equal to freezing point of 12% aqueous solution of Y. If molecular weight of X is A, then molecular weight of Y is :-
 - (1) A
 - (2) 3A
 - (3) 4A (4) 2A

(4) $\frac{1}{4}$

Sol. For same freezing point, molality of both solution should be same. $m_x = m_v$ $\frac{4\times1000}{12\times1000} = \frac{12\times1000}{12\times1000}$ $96 \times M_{v} = 88 \times M_{v}$ or, $M_y = \frac{96 \times 12}{4 \times 88} M_x = 3.27 \text{ A}$ Closest option is 3A. 6. Poly-β-hydroxybutyrate-co-βhydroxyvalerate(PHBV) is a copolymer of (1) 3-hydroxybutanoic acid and 4-hydroxypentanoic acid (2) 2-hydroxybutanoic acid and 3-hydroxypentanoic acid (3) 3-hydroxybutanoic acid and 2-hydroxypentanoic acid (4) 3-hydroxybutanoic acid and 3-hydroxypentanoic acid

Ans. (4)

- **Sol.** PHBV is a polymer of 3-hydroxybutanoic acid and 3-Hydroxy pentanoic acid.
- 7. Among the following four aromatic compounds, which one will have the lowest melting point ?







Ans. (1) Sol. M.P. of Napthalene $\simeq 80^{\circ}$ C

2

8. CH_3CH_2 -C-CH₃ cannot be prepared by : Ph (1) HCHO + PhCH(CH₃)CH₂MgX (2) PhCOCH₂CH₃ + CH₃MgX (3) PhCOCH₃ + CH₃CH₂MgX

(4)
$$CH_3CH_2COCH_3 + PhMgX$$

Ans. (1)

Sol. (1)
$$H = C = H + Ph = CH = CH_2MgX$$

 $\downarrow CH_3$
 $\downarrow CH_3$
 $Ph = CH = CH_2 = CH_2 = OH$
 CH_3

9. The volume of gas A is twice than that of gas B. The compressibility factor of gas A is thrice than that of gas B at same temperature. The pressures of the gases for equal number of moles are : (1) $2B_{1} = 2B_{2}$

(1)
$$2P_A = 3P_B$$

(2) $P_A = 3P_B$
(3) $P_A = 2P_B$
(4) $3P_A = 2P_B$
Ans. (1)

Sol.
$$V_A = 2V_B$$

 $Z_A = 3Z_B$
 $\frac{P_A V_A}{n_A R T_A} = \frac{3 \cdot P_B \cdot V_B}{n_B \cdot R T_B}$

 $2P_A = 3P_B$

- 10. The element with Z = 120 (not yet discovered) will be an/a :
 - (1) transition metal
 - (2) inner-transition metal
 - (3) alkaline earth metal
 - (4) alkali metal

Ans. (3)



Its general electronic configuration may be represented as [Nobal gas] ns², like other alkaline earth metals.

collegebatch.com

E

- 11. Decomposition of X exhibits a rate constant of 0.05 μg/year. How many years are required for the decomposition of 5 μg of X into 2.5 μg?
 (1) 50 (2) 25 (3) 20 (4) 40
- Ans.(1)
- **Sol.** Rate constant (K) = $0.05 \mu g/year$ means zero order reaction

$$t_{1/2} = \frac{a_0}{2K} = \frac{5\mu g}{2 \times 0.05 \mu g / year} = 50 \text{ year}$$

12. The major product of the following reaction is :



Ans. (4)

Sol.
$$CH_{3}O$$

 Cl_{2}
 MeO
 MeO
 Cl
 $HeAS$
 MeO
 MeO
 Cl
 $HeAS$
 MeO
 MeO
 MeO
 MeO
 Cl
 $HeAS$
 MeO
 Me

13. Given

 $\begin{array}{ccccccc} Gas & H_2 & CH_3 & CO_2 & SO_2 \\ Critical & 33 & 190 & 304 & 630 \\ Temperature/K & & & \end{array}$

On the basis of data given above, predict which of the following gases shows least adsorption on a definite amount of charcoal ?

(1)
$$H_2$$
 (2) CH_4 (3) SO_2
Ans. (1)

Sol. Smaller the value of critical temperature of gas, lesser is the extent of adsorption. so least adsorbed gas is H_2

14. For diatomic ideal gas in a closed system, which of the following plots does not correctly describe the relation between various thermodynamic quantities ?



Ans. (2)

Sol. At higher temperature, rotational degree of freedom becomes active.

$$C_{P} = \frac{7}{2}R$$
 (Independent of P)

$$C_V = \frac{3}{2}R$$
 (Independent of V)

Variation of U vs T is similar as C_V vs T.

15. The standard electrode potential E^{\odot} and its

temeprature coefficient $\left(\frac{dE^{\odot}}{dT}\right)$ for a cell are 2V

and -5×10^{-4} VK⁻¹ at 300 K respectively. The cell reaction is

$$Zn(s) + Cu^{2+} (aq) \rightarrow Zn^{2+}(aq) + Cu(s)$$

The standard reaction enthalpy $(\Delta_r H^{\odot})$ at 300

K in kJ mol⁻¹ is, [Use R = $8jK^{-1}$ mol⁻¹ and F = 96,000 Cmol⁻¹] (1) -412.8 (2) -384.0 (3) 206.4 (4) 192.0

Ans. (1)

(4) CO₂

Sol. Chiefly NO_2 , O_3 and hydrocarbon are responsible for build up smog.





- **16.** The molecule that has minimum/no role in the formation of photochemical smog, is :
 - (1) $CH_2 = O$
 - (2) N_2
 - $(3) O_3$
 - (4) NO
- Ans. (2)
- **Sol.** Chiefly NO_2 , O_3 and hydrocarbon are responsible for build up smog.
- **17.** In the Hall-Heroult process, aluminium is formed at the cathode. The cathode is made out of :
 - (1) Platinum
 - (2) Carbon
 - (3) Pure aluminium
 - (4) Copper

Ans. (2)

- 17. Ans.(2) Carbon
- **Sol.** In the Hall-Heroult process the cathode is made of carbon.
- Water samples with BOD values of 4 ppm and 18 ppm, respectively, are :
 - (1) Highly polluted and Clean
 - (2) Highly polluted and Highly polluted
 - (3) Clean and Highly polluted
 - (4) Clean and Clean

Ans. (3)

- **Sol.** Clean water would have BOD value of less than 5 ppm whereas highly polluted water could have a BOD value of 17 ppm or more.
- **19.** In the following reactions, products A and B are :







- (B)
- 20. What is the work function of the metal if the light of wavelength 4000 Å generates photoelectrons of velocity 6 × 10⁵ ms⁻¹ form it ? (Mass of electron = 9 × 10⁻³¹ kg Velocity of light = 3 × 10⁸ ms⁻¹ Planck's constant = 6.626 × 10⁻³⁴ Js Charge of electron = 1.6 × 10⁻¹⁹ JeV⁻¹) (1) 0.9 eV (2) 4.0 eV (3) 2.1 eV (4) 3.1 eV

Ans. (3)

4



156

E

Sol. $hv = \phi + hv^{\circ}$ $\frac{1}{2}mv^{2} = hc\left(\frac{1}{\lambda} - \frac{1}{\lambda_{0}}\right)$ $hv = \phi + \frac{1}{2}mv^{2}$ $\phi = \frac{6.626 \times 10^{-34} \times 3 \times 10^{8}}{4000 \times 10^{-10}} - \frac{1}{2} \times 9 \times 10^{-31} \times (6 \times 10^{5})^{2}$ $\phi = 3.35 \times 10^{-19} \text{ J} \implies \phi \simeq 2.1 \text{ eV}$ 21. Among the following compounds most basic amino acid is : (1) Lysine (2) Asparagine (3) Serine

(4) Histidine

Ans. (4)

Sol. Histidine

- 22. The metal d-orbitals that are directly facing the ligands in $K_3[Co(CN)_6]$ are :
 - (1) d_{xz} , d_{yz} and d_{z^2}
 - (2) $d_{xy}^{},\,d_{xz}^{}$ and $d_{yz}^{}$
 - (3) d_{xy} and $d_{x^2-y^2}$
 - (4) $d_{x^2-y^2}$ and d_{z^2}
- Ans. (4)
- Sol. $K_3[Co(CN)_6]$

$$\mathrm{Co}^{+3} \rightarrow \mathrm{[Ar]}_{18} \, \mathrm{3d}^6$$



- 23. The hardness of a water sample (in terms of equivalents of CaCO₃) containing 10⁻³ M CaSO₄ is :
 - (molar mass of $CaSO_4 = 136 \text{ g mol}^{-1}$)
 - (1) 100 ppm
 - (2) 50 ppm
 - (3) 10 ppm
 - (4) 90 ppm

Ans. (1)

Ε

Sol. ppm of CaCO₃ $(10^{-3} \times 10^3) \times 100 = 100 \text{ ppm}$ The correct order for acid strength of compounds 24. CH=CH, CH₃-C=CH and CH₂=CH₂ is as follows : (1) $CH \equiv CH > CH_2 = CH_2 > CH_3 - C \equiv CH$ (2) HC = CH > CH₃ – C = CH > CH₂ = CH₂ (3) $CH_3-C \equiv CH > CH_2 = CH_2 > HC \equiv CH$ (4) $CH_3-C \equiv CH > CH \equiv CH > CH_2 = CH_2$ Ans. (2) **Sol.** $CH \equiv CH > CH_3 - C \equiv CH > CH_2 = CH_2$ (Acidic strength order) 25. $Mn_2(CO)_{10}$ is an organometallic compound due to the presence of : (1) Mn - Mn bond (2) Mn - C bond (4) C - O bond (3) Mn - O bond Ans. (2) Sol. Compounds having at least one bond between carbon and metal are known as organometallic compounds.

26. The increasing order of reactivity of the following compounds towards reaction with alkyl halides directly is :







6

collegebatch.com

158

E

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On SATURDAY 12th JANUARY., 2019) TIME : 02 : 30 PM To 05 : 30 PM MATHEMATICS

3.

1. Let Ζ be the If set of integers. $A = \left\{ x \in Z : 2(x+2)(x^2 - 5x + 6) \right\} = 1$ and $B = \{x \in Z: -3 < 2x - 1 < 9\}$, then the number of subsets of the set $A \times B$, is : $(2) 2^{10}$ $(1) 2^{18}$ $(3) 2^{15}$ $(4) 2^{12}$ Ans (3)**Sol.** A = {x \in z : 2^{(x+2)(x²-5x+6)} = 1} $2^{(x+2)(x^2-5x+6)} = 2^0 \Longrightarrow x = -2, 2, 3$ $A = \{-2, 2, 3\}$ $B = \{x \in Z : -3 < 2x - 1 < 9\}$ $B = \{0, 1, 2, 3, 4\}$ $A \times B$ has is 15 elements so number of subsets of $A \times B$ is 2^{15} . If $\sin^4 \alpha + 4\cos^4 \beta + 2 = 4\sqrt{2}\sin \alpha \cos \beta$: 2. $\alpha, \ \beta \in [0, \pi]$, then $\cos(\alpha + \beta) - \cos(\alpha - \beta)$ is equal to : (2) $\sqrt{2}$ (3) -1 (4) $\sqrt{2}$ (1) 0Ans (2)**Sol.** A.M. \geq G.M. $\frac{\sin^4 \alpha + 4\cos^4 \beta + 1 + 1}{4} \ge (\sin^4 \alpha \cdot 4\cos^4 \beta \cdot 1 \cdot 1)^{\frac{1}{4}}$ $\sin^4 \alpha + 4 \cos^2 \beta + 2 \ge 4\sqrt{2} \sin \alpha \cos \beta$ given that $\sin^4 \alpha + 4 \cos^4 \beta + 2 = 4 \sqrt{2} \sin \alpha \cos \beta$ \Rightarrow A.M. = G.M. $\Rightarrow \sin^4 \alpha = 1 = 4 \cos^4 \beta$ $\sin \alpha = 1$, $\cos \beta = \pm \frac{1}{\sqrt{2}}$ $\Rightarrow \sin \beta = \frac{1}{\sqrt{2}}$ as $\beta \in [0, \pi]$ $\cos (\alpha + \beta) - \cos (\alpha - \beta) = -2 \sin \alpha \sin \beta$ $=-\sqrt{2}$

If an angle between the line, $\frac{x+1}{2} = \frac{y-2}{1} = \frac{z-3}{-2}$ and the plane, x - 2y - kz = 3 is $\cos^{-1}\left(\frac{2\sqrt{2}}{3}\right)$,

then a value of k is:

(1)
$$-\frac{5}{3}$$
 (2) $\sqrt{\frac{3}{5}}$ (3) $\sqrt{\frac{5}{3}}$ (4) $-\frac{3}{5}$

Ans (3)

S

Sol. DR's of line are 2, 1, -2

normal vector of plane is $\hat{i} - 2\hat{j} - k\hat{k}$

in
$$\alpha = \frac{(2\hat{i} + \hat{j} - 2\hat{k}).(\hat{i} - 2\hat{j} - k\hat{k})}{3\sqrt{1 + 4 + k^2}}$$

$$(1)^2 + (2)^2 = 1 \Longrightarrow k^2 = \frac{5}{3}$$

4. If a straight line passing thourgh the point P(-3, 4) is such that its intercepted portion between the coordinate axes is bisected at P, then its equation is :

(1)
$$x - y + 7 = 0$$

(2) $3x - 4y + 25 = 0$
(3) $4x + 3y = 0$
(4) $4x - 3y + 24 = 0$
(4)

Ans

1



E

Sol.
$$P(-3, 4)$$
 $B(0, b)$ $A(a, 0)$

Let the line be
$$\frac{x}{a} + \frac{y}{b} = 1$$

$$(-3, 4) = \left(\frac{a}{2}, \frac{b}{2}\right)$$
$$a = -6, b = 8$$

equation of line is 4x - 3y + 24 = 0

5. The integral
$$\int \frac{3x^{13} + 2x^{11}}{(2x^4 + 3x^2 + 1)^4} dx$$
 is equal to :

(where C is a constant of integration)

(1)
$$\frac{x^{4}}{(2x^{4}+3x^{2}+1)^{3}} + C$$

(2)
$$\frac{x^{12}}{6(2x^{4}+3x^{2}+1)^{3}} + C$$

(3)
$$\frac{x^{4}}{6(2x^{4}+3x^{2}+1)^{3}} + C$$

(4)
$$\frac{x^{12}}{(2x^{4}+3x^{2}+1)^{3}} + C$$

Ans (2)

Sol.
$$\int \frac{3x^{13} + 2x^{11}}{(2x^4 + 3x^2 + 1)^4} dx$$
$$\int \frac{\left(\frac{3}{x^3} + \frac{2}{x^5}\right) dx}{\left(2 + \frac{3}{x^2} + \frac{1}{x^4}\right)^4}$$
$$\text{Let} \left(2 + \frac{3}{x^2} + \frac{1}{x^4}\right) = t$$
$$-\frac{1}{2} \int \frac{dt}{t^4} = \frac{1}{6t^3} + C \implies \frac{x^{12}}{6(2x^4 + 3x^2 + 1)^3} + C$$

6. There are m men and two women participating in a chess tournament. Each participant plays two games with every other participant. If the number of games played by the men between themselves exceeds the number of games played between the men and the women by 84, then the value of m is :

(1) 9 (2) 11 (3) 12 (4) 7

Sol. Let m-men, 2-women

$${}^{m}C_{2} \times 2 = {}^{m}C_{1} {}^{2}C_{1} . 2 + 84$$

 ${}^{m^{2}}-5m-84 = 0 \Longrightarrow (m-12) (m+7) =$
 ${}^{m}m = 12$

0

7. If the function f given by $f(x) = x^3 - 3(a - 2)x^2 + 3ax + 7$, for some $a \in \mathbb{R}$ is increasing in (0, 1] and decreasing in [1, 5), then a root of the equation,

$$\frac{f(x) - 14}{(x - 1)^2} = 0(x \neq 1) \text{ is :}$$
(1) 6 (2) 5 (3) 7 (4) -7

Ans (3)

Sol.
$$f(x) = 3x^2 - 6(a - 2)x + 3a$$

 $f(x) \ge 0 \ \forall \ x \in (0, 1]$
 $f(x) \le 0 \ \forall \ x \in [1, 5)$
 $\Rightarrow f(x) = 0 \ at \ x = 1 \Rightarrow a = 5$
 $f(x) - 14 = (x - 1)^2 \ (x - 7)$
 $\frac{f(x) - 14}{(x - 1)^2} = x - 7$

8. Let f be a differentiable function such that f(1) = 2 and f'(x) = f(x) for all $x \in \mathbb{R}$. If h(x) = f(f(x)), then h'(1) is equal to : (1) 4e (2) 4e² (3) 2e (4) 2e²

Sol.
$$\frac{f'(x)}{f(x)} = 1 \ \forall \ x \in R$$

Intergrate & use f(1) = 2 $f(x) = 2e^{x-1} \Rightarrow f'(x) = 2e^{x-1}$ $h(x) = f(f(x)) \Rightarrow h'(x) = f'(f(x)) f'(x)$ h'(1) = f'(f(1)) f'(1) = f(2) f'(1) $= 2e \cdot 2 = 4e$





9. The tangent to the curve $y = x^2 - 5x + 5$, parallel to the line 2y = 4x + 1, also passes through the point.

$(1)\left(\frac{1}{4},\frac{7}{2}\right)$	$(2)\left(\frac{7}{2},\frac{1}{4}\right)$
$(3)\left(-\frac{1}{8},7\right)$	$(4)\left(\frac{1}{8},-7\right)$

Ans (4)

Sol. $y = x^2 - 5x + 5$

$$\frac{dy}{dx} = 2x - 5 = 2 \implies x = \frac{7}{2}$$

at $x = \frac{7}{2}$, $y = \frac{-1}{4}$

4

Equation of tangent at $\left(\frac{7}{2}, \frac{-1}{4}\right)$ is $2x - y - \frac{29}{4} = 0$

Now check options

$$x = \frac{1}{8}, y = -7$$

10. Let S be the set of all real values of λ such that a plane passing through the points $(-\lambda^2, 1, 1)$, $(1, -\lambda^2, 1)$ and $(1, 1, -\lambda^2)$ also passes through the point (-1, -1, 1). Then S is equal to :

(1) $\{\sqrt{3}\}$	(2) $\left\{\sqrt{3}-\sqrt{3}\right\}$
(3) {1, -1}	(4) {3, -3}

Ans (2)

Ε

Sol. All four points are coplaner so

$$\begin{vmatrix} 1 - \lambda^2 & 2 & 0 \\ 2 & -\lambda^2 + 1 & 0 \\ 2 & 2 & -\lambda^2 - 1 \end{vmatrix} = 0$$
$$(\lambda^2 + 1)^2 (3 - \lambda^2) = 0$$
$$\lambda = \pm \sqrt{3}$$

If a circle of radius R passes through the origin O 11. and intersects the coordinate axes at A and B, then the locus of the foot of perpendicular from O on AB is :

(1)
$$(x^2 + y^2)^2 = 4Rx^2y^2$$

(2) $(x^2 + y^2)(x + y) = R^2xy$
(3) $(x^2 + y^2)^3 = 4R^2x^2y^2$
(4) $(x^2 + y^2)^2 = 4R^2x^2y^2$

Ans (3)



Slope of AB =
$$\frac{-1}{1}$$

Equation of AB is $hx + ky = h^2 + k^2$

h

$$A\left(\frac{h^2+k^2}{h},0\right), \ B\left(0,\frac{h^2+k^2}{k}\right)$$

$$AB = 2R$$

$$\Rightarrow (h^2 + k^2)^3 = 4R^2h^2k^2$$

$$\Rightarrow (x^2 + y^2)^3 = 4R^2x^2y^2$$

The equation of a tangent to the parabola, 12. $x^2 = 8y$, which makes an angle θ with the positive direction of x-axis, is :

(1)
$$x = y \cot \theta + 2 \tan \theta$$
 (2) $x = y \cot \theta - 2 \tan \theta$
(3) $y = x \tan \theta - 2 \cot \theta$ (4) $y = x \tan \theta + 2 \cot \theta$

Ans (1) **Sol.** $x^2 = 8y$

$$\Rightarrow \frac{dy}{dx} = \frac{x}{4} = \tan \theta$$

$$\therefore x_1 = 4\tan \theta$$

$$y_1 = 2 \tan^2 \theta$$

Equation of tangent :-

$$y - 2\tan^2\theta = \tan \theta (x - 4\tan \theta)$$

$$\Rightarrow x = y \cot \theta + 2 \tan \theta$$



13. If the angle of elevation of a cloud from a point P which is 25 m above a lake be 30° and the angle of depression of reflection of the cloud in the lake from P be 60°, then the height of the cloud (in meters) from the surface of the lake is :

(1) 42
(2) 50
(3) 45
(4) 60

Ans (2)

15. $\lim_{n \to \infty} \left(\frac{n}{n^2 + 1^2} + \frac{n}{n^2 + 2^2} + \frac{n}{n^2 + 3^2} + \dots + \frac{1}{5n} \right)$ is equal to :

(1)
$$\frac{\pi}{4}$$
 (2) tan⁻¹(2)

(3)
$$\tan^{-1}(3)$$
 (4) $\frac{\pi}{2}$

Ans. (2)

$$\lim_{x \to \infty} \sum_{r=1}^{2n} \frac{n}{n^2 + r^2}$$
$$\lim_{x \to \infty} \sum_{r=1}^{2n} \frac{1}{n\left(1 + \frac{r^2}{n^2}\right)} = \int_{0}^{2} \frac{dx}{1 + x^2} = \tan^{-1} 2$$

16. The set of all values of λ for which the system of linear equations.

$$x - 2y - 2z = \lambda x$$

$$x + 2y + z = \lambda y$$

$$-x - y = \lambda z$$
has a non-trivial solution.
(1) contains more than two elements
(2) is a singleton
(3) is an empty set
(4) contains exactly two elements
Ans. (2)

$$\begin{vmatrix} \lambda - 1 & 2 & 2 \\ 1 & 2 - \lambda & 1 \\ 1 & 1 & 1 \end{vmatrix} = 0 \Longrightarrow (\lambda - 1)^3 = 0 \Longrightarrow \lambda = 1$$

n

17. If ${}^{n}C_{4}$, ${}^{n}C_{5}$ and ${}^{n}C_{6}$ are in A.P., then n can be: (1) 14 (2) 11 (3) 9 (4) 12

Ans. (1)

$$2.nC_5 = nC_4 + nC_6$$

$$|n| |n|$$

$$2. \frac{12}{|5|n-5} = \frac{12}{|4|n-4} + \frac{12}{|6|n-6}$$

$$\frac{2}{5} \cdot \frac{1}{n-5} = \frac{1}{(n-4)(n-5)} + \frac{1}{30}$$

n = 14 satisfying equation.

4



18. Let \vec{a}, \vec{b} and \vec{c} be three unit vectors, out of which vectors \vec{b} and \vec{c} are non-parallel. If α and β are the angles which vector \vec{a} makes with vectors \vec{b} and \vec{c} respectively and $\vec{a} \times (\vec{b} \times \vec{c}) = \frac{1}{2}\vec{b}$, then $|\alpha - \beta|$ is equal to : (1) 60° (2) 30° (3) 90° (4) 45° Ans. (2)

> $(\vec{a}.\vec{c})\vec{b} - (\vec{a}.\vec{b}).\vec{c} = \frac{1}{2}\vec{b}$ $\therefore \vec{b} \ll \vec{c}$ are linearly independent

$$\therefore \quad \vec{a}.\vec{c} = \frac{1}{2} \& \vec{a}.\vec{b} = 0$$

(All given vectors are unit vectors)

$$\therefore \quad \vec{a} \wedge \vec{c} = 60^{\circ} \quad \& \quad \vec{a} \wedge \vec{b} = 90^{\circ}$$
$$\therefore \quad |\alpha - \beta| = 30^{\circ}$$

19. If $A = \begin{bmatrix} 1 & \sin \theta & 1 \\ -\sin \theta & 1 & \sin \theta \\ -1 & -\sin \theta & 1 \end{bmatrix}$; then for all

$$\theta \in \left(\frac{3\pi}{4}, \frac{5\pi}{4}\right)$$
, det(A) lies in the interval :

(1)
$$\left[\frac{5}{2}, 4\right]$$
 (2) $\left(\frac{3}{2}, 3\right]$
(3) $\left(0, \frac{3}{2}\right]$ (4) $\left(1, \frac{5}{2}\right]$

Ans (2)

$$|\mathbf{A}| = \begin{vmatrix} 1 & \sin \theta & 1 \\ -\sin \theta & 1 & \sin \theta \\ -1 & -\sin \theta & 1 \end{vmatrix}$$
$$= 2(1 + \sin^2 \theta)$$
$$\theta \in \left(\frac{3\pi}{4}, \frac{5\pi}{4}\right) \Rightarrow \frac{1}{\sqrt{2}} < \sin \theta < \frac{1}{\sqrt{2}}$$
$$\Rightarrow 0 \le \sin^2 \theta < \frac{1}{2}$$
$$\therefore |\mathbf{A}| \in [2, 3)$$

20.
$$\lim_{x \to 1^{-}} \frac{\sqrt{\pi} - \sqrt{2\sin^{-1}x}}{\sqrt{1-x}} \text{ ie equal to :}$$
(1) $\frac{1}{\sqrt{2\pi}}$ (2) $\sqrt{\frac{\pi}{2}}$ (3) $\sqrt{\frac{2}{\pi}}$ (4) $\sqrt{\pi}$

Ans. (3)

$$\lim_{x \to 1^{-}} \frac{\sqrt{\pi} - \sqrt{2\sin^{-1}x}}{\sqrt{1-x}} \times \frac{\sqrt{\pi} + \sqrt{2\sin^{-1}x}}{\sqrt{\pi} + \sqrt{2\sin^{-1}x}}$$

$$\lim_{x \to 1^{-}} \frac{2\left(\frac{\pi}{2} - \sin^{-1}x\right)}{\sqrt{1 - x}\left(\sqrt{\pi} + \sqrt{2\sin^{-1}x}\right)}$$

$$\lim_{\mathbf{x}\to\mathbf{1}^{-}}\frac{2\cos^{-1}\mathbf{x}}{\sqrt{1-\mathbf{x}}}\cdot\frac{1}{2\sqrt{\pi}}$$

Put $x = \cos\theta$

$$\lim_{\theta \to 0^+} \frac{2\theta}{\sqrt{2}\sin\left(\frac{\theta}{2}\right)} \cdot \frac{1}{2\sqrt{\pi}} = \sqrt{\frac{2}{\pi}}$$

21. The expression $\sim (\sim p \rightarrow q)$ is logically equivalent to : (1) $\sim p^{\wedge} \sim q$ (2) $p^{\wedge} q$ (3) $\sim p^{\wedge} q$ (4) $p^{\wedge} \sim q$

Ans. (1)

р	q	~p	~p → q	~(~p→q)	$(\sim p \land \sim q)$
Т	Т	F	Т	F	F
F	Т	Т	Т	F	F
Т	F	F	Т	F	F
F	F	Т	F	Т	Т

22. The total number of irrational terms in the binomial

expansion of $(7^{1/5} - 3^{1/10})^{60}$ is : (1) 55 (2) 49 (3) 48 (4) 54

Ans. (4)

General term $T_{r+1} = {}^{60}C_r \quad 7^{\frac{60-r}{5}} \quad 3^{\frac{r}{10}}$ \therefore for rational term, r = 0, 10, 20, 30, 40, 50, 60 \Rightarrow no of rational terms = 7 \therefore number of irrational terms = 54

Ε



- 23. The mean and the variance of five observation are 4 and 5.20, respectively. If three of the observations are 3, 4 and 4; then then absolute value of the difference of the other two observations, is :
 - (1) 1 (2) 3 (3) 7 (4) 5

Ans. (3)

mean $\overline{\mathbf{x}} = 4, \sigma^2 = 5.2, n = 5, x_1 = 3, x_2 = 4 = x_3$

$$\sum_{i} x_{i} = 20$$

$$x_{4} + x_{5} = 9 \dots (i)$$

$$\frac{\sum_{i} x_{i}^{2}}{x} - (\overline{x})^{2} = \sigma^{2} \implies \sum_{i} x_{i}^{2} = 106$$

$$x_{4}^{2} + x_{5}^{2} = 65 \dots (ii)$$

Using (i) and (ii) $(x_4 - x_5)^2 = 49$

$$|x_4 - x_5| = 7$$

24. If the sum of the first 15 tems of the

series
$$\left(\frac{3}{4}\right)^3 + \left(1\frac{1}{2}\right)^3 + \left(2\frac{1}{4}\right)^3 + 3^3 + \left(3\frac{3}{4}\right)^3 + \dots$$
 is

(4) 54

equal to 225 k, then k is equal to :

(1) 9 (2) 27 (3) 108 Ans. (2)

$$= \frac{27}{64} \sum_{r=1}^{12} r^{3}$$
$$= \frac{27}{64} \cdot \left[\frac{15(15+1)}{2}\right]^{2}$$
$$= 225 \text{ K (Given in question)}$$
$$K = 27$$

25. Let S and S' be the foci of the ellipse and B be any one of the extremities of its minor axis. If Δ S'BS is a right angled triangle with right angle at B and area (Δ S'BS) = 8 sq. units, then the length of a latus rectum of the ellipse is :

(1) $2\sqrt{2}$ (2) 2 (3) 4 (4) $4\sqrt{2}$

Ans. (3)

 $m_{_{\rm SB}}$. $m_{_{\rm S'B}}$ = -1



$$b^2 = a^2 e^2 \dots (i)$$

$$\frac{1}{2}$$
S'B. SB = 8
S'B. SB = 16
 $a^2e^2 + b^2 = 16$ (ii)

$$b^2 = a^2 (1 - e^2) \dots (iii)$$

using (i).(ii). (iii) $a = 4$

$$b = 2\sqrt{2}$$
$$e = \frac{1}{\sqrt{2}}$$
$$\therefore \ \ell \ (L.R) = \frac{2b^2}{a} = 4 \quad \text{[Ans.3]}$$

26. In a class of 60 students, 40 opted for NCC, 30 opted for NSS and 20 opted for both NCC and NSS. If one of these students is selected at random, then the probability that the student selected has opted neither for NCC nor for NSS is :

(1)
$$\frac{2}{3}$$
 (2) $\frac{1}{6}$ (3) $\frac{1}{3}$ (4) $\frac{5}{6}$

Ans. (2)

A → opted NCC
B → opted NSS
∴ P (nither A nor B) =
$$\frac{10}{60} = z\frac{1}{6}$$

6



27. The number of integral values of m for which the quadratic expression. $(1 + 2m)x^2 - 2(1 + 3m)x + 4(1 + m), x \in \mathbb{R}$, is always positive, is : (1) 8(2) 7(3) 6(4) 3Ans. (2) Exprsssion is always positve it $2m+1 > 0 \Longrightarrow m > -\frac{1}{2}$ & $D < 0 \implies m^2 - 6m - 3 < 0$ $3 - \sqrt{12} < m < 3 + \sqrt{12}$ (iii) .: Common interval is $3 - \sqrt{12} < m < 3 + \sqrt{12}$: Intgral value of m $\{0, 1, 2, 3, 4, 5, 6\}$ In a game, a man wins Rs. 100 if he gets 5 of 6 28. on a throw of a fair die and loses Rs. 50 for getting any other number on the die. If he decides to throw the die either till he gets a five or a six or to a maximum of three throws, then his expected gain/

(1)
$$\frac{400}{3}$$
 gain (2) $\frac{400}{3}$ loss

loss (in rupees) is :

(3) 0 (4)
$$\frac{400}{9}$$
 loss

Ans. (3)

Expected Gain/ Loss = = $w \times 100 + Lw (-50 + 100) + L^2w (-50 - 50 + 100) + L^3 (-150)$

$$= \frac{1}{3} \times 100 + \frac{2}{3} \cdot \frac{1}{3} (50) + \left(\frac{2}{3}\right)^2 \left(\frac{1}{3}\right) (0) + \left(\frac{2}{3}\right)^3 (-150) = 0$$

here w denotes probability that outcome 5 or 6 (

$$w=\frac{2}{6}=\frac{1}{3})$$

here L denotes probability that outcome

1,2,3,4 (L =
$$\frac{4}{6} = \frac{2}{3}$$
)

29. If a cuver passes through the point (1, -2) and has slope of the tangent at any point (x, y) on it as

 $\frac{x^2 - 2y}{x}$, then the curve also passes through the point :

(1)
$$\left(-\sqrt{2},1\right)$$
 (2) $\left(\sqrt{3},0\right)$
(3) $(-1, 2)$ (4) $(3, 0)$

Ans. (2)

$$\frac{dy}{dx} = \frac{x^2 - 2y}{x}$$
 (Given)

$$\frac{dy}{dx} + 2\frac{y}{x} = x$$

I.F = $e^{\int \frac{2}{x} dx} = x^2$

$$y.x^2 = \int x.x^2 dx + C$$

$$=\frac{x^4}{y}+C$$

hence bpasses through $(1, -2) \Rightarrow C = -\frac{9}{4}$

$$yx^2 = \frac{x^4}{4} - \frac{9}{4}$$

Now check option(s) , Which is satisly by option (ii)

30. Let Z_1 and Z_2 be two complex numbers satisfying $|Z_1| = 9$ and $|Z_2-3-4i|=4$. Then the minimum value of $|Z_1-Z_2|$ is :

(1) 0 (2) 1 (3)
$$\sqrt{2}$$
 (4) 2

Ans. (1)

$$\begin{aligned} |z_1| &= 9 \ , \ |z_2 - (3 + 4i)| &= 4 \\ C_1 (0, 0) \text{ radius } r_1 &= 9 \\ C_2 (3, 4), \text{ radius } r_2 &= 4 \\ C_1 C_2 &= |r_1 - r_2| &= 5 \\ \therefore \text{ Circle touches internally} \\ \therefore \ |z_1 - z_2|_{\min} &= 0 \end{aligned}$$

7

collegebatch.com

TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Saturday 12th JANUARY, 2019) TIME : 02 : 30 PM To 05 : 30 PM PHYSICS

1. A load of mass M kg is suspended from a steel wire of length 2 m and radius 1.0 mm in Searle's apparatus experiment. The increase in length produced in the wire is 4.0 mm. Now the load is fully immersed in a liquid of relative density 2. The relative density of the material of load is 8. The new value of increase in length of the steel wire is :

(1) 4.0mm (2) 3.0mm (3) 5.0mm (4) zero





$$\frac{F}{A} = y \cdot \frac{\Delta \ell}{\ell}$$
$$\Delta \ell \propto F \qquad \dots (i)$$
$$T = mg$$

$$T = mg - f_{B} = mg - \frac{m}{\rho_{b}} \cdot \rho_{\ell} \cdot g$$
$$= \left(1 - \frac{\rho_{\ell}}{\rho_{b}}\right) mg$$
$$= \left(1 - \frac{2}{8}\right) mg$$
$$T' = \frac{3}{4} mg$$

From (i)

E

$$\frac{\Delta \ell'}{\Delta \ell} = \frac{T'}{T} = \frac{3}{4}$$
$$\Delta \ell' = \frac{3}{4} \cdot \Delta \ell = 3 \text{ mm}$$

2. Formation of real image using a biconvex lens is shown below :



If the whole set up is immersed in water without disturbing the object and the screen position, what will one observe on the screen ?

- (1) Image disappears
- (2) No change
- (3) Erect real image
- (4) Magnified image

Ans (1)

Sol. From
$$\frac{1}{f} = (\mu_{rel} - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$

Focal length of lens will change hence image disappears from the screen.

3. A vertical closed cylinder is separated into two parts by a frictionless piston of mass m and of negligible thickness. The piston is free to move along the length of the cylinder. The length of the cylinder above the piston is ℓ_1 , and that below the piston is ℓ_2 , such that $\ell_1 > \ell_2$. Each part of the cylinder contains n moles of an ideal gas at equal temperature T. If the piston is stationary, its mass, m, will be given by :

(R is universal gas constant and g is the acceleration due to gravity)

(1)
$$\frac{\mathrm{nRT}}{\mathrm{g}} \left[\frac{1}{\ell_2} + \frac{1}{\ell_1} \right]$$
 (2)
$$\frac{\mathrm{nRT}}{\mathrm{g}} \left[\frac{\ell_1 - \ell_2}{\ell_1 \ell_2} \right]$$

(3)
$$\frac{\mathrm{RT}}{\mathrm{g}} \left[\frac{2\ell_1 + \ell_2}{\ell_1 \ell_2} \right]$$
 (4)
$$\frac{\mathrm{RT}}{\mathrm{ng}} \left[\frac{\ell_1 - 3\ell_2}{\ell_1 \ell_2} \right]$$

Ans (2)





$$P_{2}A = P_{1}A + mg$$

$$\frac{nRT.A}{A\ell_{2}} = \frac{nRT.A}{A\ell_{1}} + mg$$

$$nRT\left(\frac{1}{\ell_{2}} - \frac{1}{\ell_{1}}\right) = mg$$

$$m = \frac{nRT}{g}\left(\frac{\ell_{1} - \ell_{2}}{\ell_{1}.\ell_{2}}\right)$$

4. A simple harmonic motion is represented by: $y = 5(\sin 3\pi t + \sqrt{3} \cos 3\pi t) \text{ cm}$

The amplitude and time period of the motion are:

(1) 5cm,
$$\frac{3}{2}$$
s
(2) 5cm, $\frac{2}{3}$ s
(3) 10cm, $\frac{3}{2}$ s
(4) 10cm, $\frac{2}{3}$ s

Ans. (4)



$$y = 5 \left[\sin(3\pi t) + \sqrt{3}\cos(3\pi t) \right]$$

$$= 10\sin\left(3\pi t + \frac{\pi}{3}\right)$$

Amplitude = 10 cm

$$T = \frac{2\pi}{w} = \frac{2\pi}{3\pi} = \frac{2}{3} \sec \theta$$

5. In the given circuit diagram, the currents, $I_1 = -0.3A$, $I_4 = 0.8 A$ and $I_5 = 0.4 A$, are flowing as shown. The currents I_2 , I_3 and I_6 , respectively, are :



A particle of mass 20 g is released with an initial velocity 5 m/s along the curve from the point A, as shown in the figure. The point A is at height h from point B. The particle slides along the frictionless surface. When the particle reaches point B, its angular momentum about O will be : (Take $g = 10 \text{ m/s}^2$)



6.



Sol. Work Energy Theorem from A to B $mgh = \frac{1}{2}mv_B^2 - \frac{1}{2}mv_A^2$

In the above circuit, $C = \frac{\sqrt{3}}{2} \mu F$, $R_2 = 20\Omega$,

$$L = \frac{\sqrt{3}}{10}$$
 H and $R_1 = 10\Omega$. Current in L-R₁ path

is I_1 and in C-R₂ path it is I_2 . The voltage of A.C source is given by

 $V=200\sqrt{2}\sin(100t)$ volts. The phase difference between I_1 and I_2 is :

 $(4) 60^{\circ}$ $(1) 30^{\circ}$ $(2) 0^{\circ}$ $(3) 90^{\circ}$

Ans. (Bonus)

7.



 $\tan\theta/2 \frac{x_e}{R_e} = \frac{10^3}{\sqrt{3}}$ θ_1 is close to 90 For L-R circuit $x_{L} = w_{L} = 100 \times \frac{\sqrt{3}}{10} = \sqrt{3}$ θ. i $R_1 = 10$ $\tan \theta_2 = \frac{\mathbf{x}_e}{\mathbf{R}}$ $\tan\theta_2 = \sqrt{3}$ $\theta_2 = 60$ So phase difference comes out 90 + 60 = 150. Therefore Ans. is Bonus If R_2 is 20 K Ω then phase difference comes out to be 60+30 $= 90^{\circ}$ A paramagnetic material has 10²⁸ atoms/m³. Its magnetic susceptibility at temperature 350 K is 2.8 $\times 10^{-4}$. Its susceptibility at 300 K is : (1) 3.672×10^{-4} (2) 3.726×10^{-4} (3) 3.267×10^{-4} (4) 2.672×10^{-4} Ans (3) **Sol.** $x \alpha \frac{1}{T_c}$ curie law for paramagnetic substane $\frac{x_1}{x_2} = \frac{T_{C_2}}{T_{C_1}}$ $\frac{2.8 \times 10^{-4}}{x_2} = \frac{300}{350}$

$$x_2 = \frac{2.8 \times 350 \times 10^{-4}}{300}$$
$$= 3.266 \times 10^{-4}$$

8.

3



- 9. A 10 m long horizontal wire extends from North East to South West. It is falling with a speed of 5.0ms^{-1} , at right angles to the horizontal component of the earth's magnetic field, of $0.3 \times 10^{-4} \text{Wb/m}^2$. The value of the induced emf in wire is :
 - (1) 2.5×10^{-3} V (2) 1.1×10^{-3} V (3) 0.3×10^{-3} V (4) 1.5×10^{-3} V

10

Ans (4)

Sol. Induied $emf = Bv\ell$

$$= 0.3 \times 10^{-4} \times 5 \times$$

= 1.5 × 10^{-3} V



In the figure, given that V_{BB} supply can vary from 0 to 5.0 V, $V_{CC} = 5V$, $\beta_{dc} = 200$, $R_B = 100 \text{ k}\Omega$, $R_C = 1 \text{ k}\Omega$ and $V_{BE} = 1.0 \text{ V}$, The minimum base current and the input voltage at which the transistor will go to saturation, will be, respectively :

(1) 20μA and 3.5V
(2) 25μA and 3.5V
(3) 25μA and 2.5V
(4) 20μA and 2.8V

Ans (2)

Sol. At saturation, $V_{CE} = 0$ $V_{CE} = V_{CC} - I_C R_C$ $\Rightarrow I_C = \frac{V_{CC}}{R_C} = 5 \times 10^{-3} \text{ A}$

 I_{C}

Given

$$\beta_{dc} = \frac{1}{I_B}$$

$$I_B = \frac{5 \times 10^{-3}}{200}$$

$$I_B = 25 \ \mu A$$
At input side
$$V_{BB} = I_B R_B + V_{BE}$$

$$= (25 \text{ mA}) (100 \text{ k}\Omega) + 1 \text{ V}$$

$$V_{BB} = 3.5 \ \text{ V}$$

11. In the circuit shown, find C if the effective capacitance of the whole circuit is to be 0.5 μ F. All values in the circuit are in μ F.



(1)
$$\frac{7}{10}\mu F$$
 (2) $\frac{7}{11}\mu F$ (3) $\frac{6}{5}\mu F$ (4) $4\mu F$

Ans (2)



12. Two satellites, A and B, have masses m and 2m respectively. A is in a circular orbit of radius R, and B is in a circular orbit of radius 2R around the earth. The ratio of their kinetic energies, T_A/T_B , is:

(1) 2 (2)
$$\sqrt{\frac{1}{2}}$$
 (3) 1 (4) $\frac{1}{2}$

Ans (3)

Sol. Orbital velocity
$$V = \sqrt{\frac{GMe}{r}}$$

 $T_A = \frac{1}{2}m_A V_A^2$
 $T_B = \frac{1}{2}m_B V_B^2$
 $\Rightarrow \frac{T_A}{T_B} = \frac{m \times \frac{Gm}{R}}{2m \times \frac{Gm}{2R}}$
 $\Rightarrow \frac{T_A}{T_B} = 1$

4



13. The moment of inertia of a solid sphere, about an axis parallel to its diameter and at a distance of x from it, is I(x)'. Which one of the graphs represents the variation of I(x) with x correctly?





Ans. (2)

- **Sol.** $I = \frac{2}{5}mR^2 + mx^2$
- 14. When a certain photosensistive surface is illuminated with monochromatic light of frequency v, the stopping potential for the photo current is $V_0/2$. When the surface is illuminated by monochromatic light of frequency v/2, the stopping potential is – V_0 . The threshold frequency for photoelectric emission is:

(1)
$$\frac{3\nu}{2}$$
 (2) 2ν (3) $\frac{4}{3}\nu$ (4) $\frac{5\nu}{3}$

Ans. (BONUS)

15. A galvanometer, whose resistance is 50 ohm, has 25 divisions in it. When a current of 4×10^{-4} A passes through it, its needle (pointer) deflects by one division. To use this galvanometer as a voltmeter of range 2.5 V, it should be connected to a resistance of:

(1) 6250 ohm	(2) 250 ohm
(3) 200 ohm	(4) 6200 ohm

Ans. (3)

E

Sol.
$$I_{\sigma} = 4 \times 10^{-4} \times 25 = 10^{-2} \text{ A}$$

$$\begin{array}{c} G & 1_{g} \\ 50\Omega & R \\ 2.5V \end{array}$$

16. A long cylindrical vessel is half filled with a liquid. When the vessel is rotated about its own vertical axis, the liquid rises up near the wall. If the radius of vessel is 5 cm and its rotational speed is 2 rotations per second, then the difference in the heights between the centre and the sides, in cm, will be:

(1) 1.2 (2) 0.1 (3) 2.0 (4) 0.4

Ans. (3)

$$y = \frac{\omega^2 x^2}{2g} = \frac{(2 \times 2\pi)^2 \times (0.05)^2}{20} \simeq 2 \text{ cm}$$

17. Two particles A, B are moving on two concentric circles of radii R_1 and R_2 with equal angular speed ω . At t = 0, their positions and direction of motion are shown in the figure :



The relative velocity $\vec{v}_{A} - \vec{v}_{B}$ at $t = \frac{\pi}{2\omega}$ is given by :

(1)
$$-\omega(R_1 + R_2)i$$
 (2) $\omega(R_1 + R_2)i$
(3) $\omega(R_1 - R_2)i$ (4) $\omega(R_2 - R_1)i$

Ans. (4)





18. A plano-convex lens (focal length f_2 , refractive index μ_2 , radius of curvature R) fits exactly into a plano-concave lens (focal length f_1 , refractive index μ_1 , radius of curvature R). Their plane surfaces are parallel to each other. Then, the focal length of the combination will be :

(1)
$$f_1 - f_2$$
 (2) $f_1 + f_2$
(3) $\frac{R}{\mu_2 - \mu_1}$ (4) $\frac{2f_1f_2}{f_1 + f_2}$

Ans. (3)



19. Let ℓ , r, c and v represent inductance, resistance, capacitance and voltage, respectively. The

dimension of $\frac{\ell}{rcv}$ in SI units will be: (1) [LTA] (2) [LA⁻²] (3) [A⁻¹] (4) [LT²] Ans. (3)

Sol. $\left\lfloor \frac{\ell}{r} \right\rfloor = T$ [CV] = AT So, $\left\lfloor \frac{\ell}{rCV} \right\rfloor = \frac{T}{AT} = A^{-1}$

- **20.** In a radioactive decay chain, the initial nucleus is $^{232}_{90}$ Th . At the end there are 6 α -particles and 4 β -particles which are emitted. If the end nucleus, If $^{A}_{7}$ X , A and Z are given by :
 - (1) A = 208; Z = 80
 (2) A = 202; Z = 80
 (3) A = 200; Z = 81
 (4) A = 208; Z = 82

Ans. (4)

Sol.
$${}^{232}_{90}$$
Th $\longrightarrow^{208}_{78}$ Y + ${}^{4}_{2}$ He

$$^{208}_{78}$$
 Y \longrightarrow $^{208}_{82}$ X + 4 β praticle

- 21. The mean intensity of radiation on the surface of the Sun is about 10⁸ W/m². The rms value of the corresponding magnetic field is closest to :
 - (1) 10^2 T (2) 10^{-4} T

(3) 1T (4) 10⁻²T

Ans. (2)

Sol. I =
$$\varepsilon_0 C E_{rms}^2$$

& $E_{rms} = cB_{rms}$
I = $\varepsilon_0 C^3 B_{rms}^2$
 $B_{rms} = \sqrt{\frac{I}{\epsilon_0 C^3}}$
 $B_{rms} \approx 10^{-4}$

22. A resonance tube is old and has jagged end. It is still used in the laboratory to determine velocity of sound in air. A tuning fork of frequency 512 Hz produces first resonance when the tube is filled with water to a mark 11 cm below a reference mark, near the open end of the tube. The experiment is repeated with another fork of frequency 256 Hz which produces first resonance when water reaches a mark 27 cm below the reference mark. The velocity of sound in air, obtained in the experiment, is close to:

(1) 328ms ⁻¹	(2) 322ms ⁻¹
(3) 341ms ⁻¹	(4) 335ms ⁻¹

Ans. (1)

6



- 23. An ideal gas is enclosed in a cylinder at pressure of 2 atm and temperature, 300 K. The mean time between two successive collisions is 6×10^{-8} s. If the pressure is doubled and temperature is increased to 500 K, the mean time between two successive collisions will be close to:
 - (1) 4×10^{-8} s (2) 3×10^{-6} s (3) 2×10^{-7} s (4) 0.5×10^{-8} s
- Ans. (1)

Sol. $t \propto \frac{\text{Volume}}{\text{velocity}}$ volume $\propto \frac{T}{T}$

P
∴ t ∝
$$\frac{\sqrt{T}}{P}$$

 $\frac{t_1}{6 \times 10^{-8}} = \frac{\sqrt{500}}{2P} \times \frac{P}{\sqrt{300}}$
 $t_1 = 3.8 \times 10^{-8}$
 $\approx 4 \times 10^{-8}$

24. The charge on a capacitor plate in a circuit, as a function of time, is shown in the figure:What is the value of current at t = 4 s ?



Ans. (3)

Ε

25. A block kept on a rough inclined plane, as shown in the figure, remains at rest upto a maximum force 2 N down the inclined plane. The maximum external force up the inclined plane that does not move the block is 10 N. The coefficient of static friction betwreen the block and the plane is : [Take $g = 10 \text{ m/s}^2$]



collegebatch.com

Sol. 2 + mg sin30 = µmg cos30°
10 = mgsin 30 + µ mg cos30°
= 2µmg cos30 -2
6 = µmg cos 30
4 = mg sin 30

$$\frac{3}{2} = \mu \times \sqrt{3}$$

µ = $\frac{\sqrt{3}}{2}$

26. An alpha-particle of mass m suffers 1-dimensional elastic coolision with a nucleus at rest of unknown mass. It is scattered directly backwards losing, 64% of its initial kinetic energy. The mass of the nucleus is :-

(1) 4 m (2) 3.5 m (3) 2 m (4) 1.5 m Ans. (1)

$$mv_0 = mv_2 - mv_1$$

 $\frac{1}{2}mV_1^2 = 0.36 \times \frac{1}{2}mV_0^2$
 $v_1 = 0.6v_0$

$$\frac{1}{2}MV_2^2 = 0.64 \times \frac{1}{2}mV_0^2$$
$$V_2 = \sqrt{\frac{m}{M}} \times 0.8V_0$$
$$mV_0 = \sqrt{mM} \times 0.8V_0 - m \times 0.6V_0$$
$$\Rightarrow 1.6m = 0.8\sqrt{mM}$$
$$4m^2 = mM$$

27. A soap bubble, blown by a mechanical pump at the mough of a tube, increases in volume, with time, at a constant rate. The graph that correctly depicts the time dependence of pressure inside the bubble is given by :-



28. To double the coverging range of a TV transmittion tower, its height should be multiplied by :-

(1)
$$\frac{1}{\sqrt{2}}$$
 (2) 4 (3) $\sqrt{2}$ (4) 2

Ans. (2)

29. A parallel plate capacitor with plates of area 1m² each, area t a separation of 0.1 m. If the electric field between the plates is 100 N/C, the magnitude of charge each plate is :-

(Take
$$\varepsilon_0 = 8.85 \times 10^{-12} \frac{\text{C}^2}{\text{N} - \text{m}^2}$$
)
(1) 7.85 × 10⁻¹⁰ C (2) 6.85 × 10⁻¹⁰ C
(3) 9.85 × 10⁻¹⁰ C (4) 8.85 × 10⁻¹⁰ C

Ans. (4)

Sol.
$$E = \frac{\sigma}{\epsilon_0} = \frac{Q}{A \epsilon_0}$$

$$\begin{split} \mathbf{Q} &= \mathbf{AE} \in_0 \\ \mathbf{Q} &= (1)(100)(8.85{\times}10^{-12}) \\ \mathbf{Q} &= 8.85 \times 10^{-10} \mathbf{C} \end{split}$$

30. In a Frank-Hertz experiment, an electron of energy 5.6 eV passes through mercury vapour and emerges with an energy 0.7 eV. The minimum wavelength of photons emitted by mercury atoms is close to :-

(1) 2020 nmY:\node05\JEE-Main 2019-(On line)\12-01-2019\Evening\PDF (2) 220 nm

(4) 1700 nm

(3) 250 nm

Sol.
$$\lambda = \frac{1240}{5.6 - 0.7}$$
 nm



TEST PAPER OF JEE(MAIN) EXAMINATION – 2019 (Held On Saturday 12th JANUARY, 2019) TIME : 02 : 30 PM To 05 : 30 PM CHEMISTRY

1. 8g of NaOH is dissolved in 18g of H_2O . Mole fraction of NaOH in solution and molality (in mol kg⁻¹) of the solutions respectively are :

(1) 0.167, 11.11	(2) 0.2, 22.20

(3) 0.2, 11.11 (4) 0.167,22.20

Ans. (1)

Sol. 8g NaOH, mol of NaOH =
$$\frac{8}{40}$$
 = 0.2mol

18g H₂O, mol of H₂O =
$$\frac{18}{18}$$
 = 1mol

$$\therefore X_{\text{NaOH}} = \frac{0.2}{1.2} = 0.167$$

Molality = $\frac{0.2 \times 1000}{18} = 11.11$ m

- 2. The correct statement(s) among I to III with respect to potassium ions that are abundant within the cell fluids is/are :
 - I. They activate many enzymes
 - II. They participate in the oxidation of glucose to produce ATP
 - III. Along with sodium ions, they are responsible for the transmission of nerve signals
 - (1) I, II and III (2) I and III only
 - (3) III only (4) I and II only
- Ans. (1)
- Sol. All the three statements are correct a/c to NCERT (s-block)
- 3. The magnetic moment of an octahedral homoleptic Mn(II) complex is 5.9 BM. The suitable ligand for this complex is :

(1) CN ⁻	(2) NCS ⁻
(3) CO	(4) ethylenediamine

Ans. (2)

Ε

Sol. $\mu = 5.9 \text{ BM}$ \therefore n (no of unpaired.e⁻) = 5 Cation Mn^{II} - 3d⁵ confn only possible for relatively weak ligand.

∴ NCS⁻

4. The correct structure of histidine in a strongly acidic solution (pH=2) is



Ans. (1)

Sol. Histidine is



Zwitter ionic form

pIn = 7.59

5. The compound that is NOT a common component of photochemical smog is :

(1)
$$O_3$$
 (2) CH_2 =CHCHO

(3)
$$CF_2Cl_2$$
 (4) $H_3C-C-OONO_2$

Ans. (3)

Sol. Freons (CFC's) are not common components of photo chemical smog.



- 6. The upper stratosphere consisting of the ozone layer protects us from the sun's radiation that falls in the wavelength region of : (1) 600-750 nm (2) 0.8-1.5 nm (3) 400-550 nm (4) 200-315 nm Ans. (4) Ozone protects most of the medium Sol. freequnecies ultravoilet light from 200 - 315 nm wave length. 7. The major product of the following reaction is: CH₂CH₃ $H_3C \longrightarrow \overset{\blacksquare}{\underline{C}} \checkmark Cl \qquad \xrightarrow{\text{NaOEt}} \Delta$ **Ē**OOCH₂CH₃ CH₂CH₃ (1) H₃C ► Č **O**CH₂CH₃ **COOCH**₂CH₂ OCH₂CH₃ (2) $H_3CH_2C \frown CO_2CH_2CH_3$ ĒH. CO₂CH₂CH₃ (3) CH₃C=CHCH₃ (4) CH₃CH₂C=CH₂ I CO₂CH₂CH₃ Ans. (3) Sol.
- $H_{3}C \xrightarrow{C} O_{-}CH_{2}-CH_{3} \xrightarrow{C} O_{2}CH_{2}-CH_{3}$ $H_{3}C \xrightarrow{C} O_{-}CH_{2}-CH_{3} \xrightarrow{C} CH_{3}-C=CH_{-}CH_{3}$ $\xrightarrow{C} O_{-}CH_{2}-CH_{3} \xrightarrow{C} O_{-}CH_{2}-CH_{3}$ $\xrightarrow{C} O_{-}CH_{2}-CH_{3} \xrightarrow{C} O_{-}CH_{2}-CH_{3}$ $\xrightarrow{C} O_{-}CH_{2}-CH_{3} \xrightarrow{C} O_{-}CH_{2}-CH_{3}$ $\xrightarrow{C} O_{-}CH_{3}-CH_{3}$ $\xrightarrow{C} O_{-}CH_{3}-CH$
- 8. The increasing order of the reactivity of the following with $LiAlH_4$ is :

$$(A) C_{2}H_{5} NH_{2} (B) C_{2}H_{5} OCH_{3}$$

$$(A) C_{2}H_{5} CI (D) C_{2}H_{5} OCH_{3}$$

$$(C) C_{2}H_{5} CI (D) C_{2}H_{5} OCH_{3}$$

$$(1) (A) < (B) < (D) < (C)$$

$$(2) (A) < (B) < (C) < (D)$$

$$(3) (B) < (A) < (D) < (C)$$

$$(4) (B) < (A) < (C) < (D)$$

- Ans. (1)
- Sol. Rate of nucleophilic \propto Electrophilicity of carbonyl group

$$C_{2}H_{5}-C-\overset{\mathbf{N}}{\underset{+m}{\overset{-}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{-}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+}}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+}}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+}}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+}}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+}}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+}}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+m}}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+m}}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+m}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+m}{\overset{+m}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+m}}} = C_{2}H_{5}-C-\overset{\mathbf{O}}{\underset{+m}{\overset{+m}{\overset{+m}{}}} = C_$$

9. The major product of the following reaction is:



- Ans. (4)
- **Sol.** NaBH₄ can not reduce C=C

but can reduce -C - into OH.



10. Molecules of benzoic acid (C_6H_5COOH) dimerise in benzene. 'w' g of the acid dissolved in 30 g of benzene shows a depression in freezing point equal to 2K. If the percentage association of the acid to form dimer in the solution is 80, then w is :

(Given that $K_f = 5 \text{ K kg mol}^{-1}$, Molar mass of benzoic acid = 122 g mol}^{-1})

(1) 1.8 g (2) 2.4 g (3) 1.0 g (4) 1.5 g Ans. (2)

Sol.

$$2(C_{6}H_{5}COOH) \xrightarrow{C_{6}H_{6}} (C_{6}H_{5}COOH)_{2}$$

$$\Delta_{f}T = i k_{f} m$$

$$2 = 0.6 \times 5 \times \frac{W \times 1000}{122 \times 30}$$

$$(i = 1 - 0.8 + 0.4 = 0.6)$$

$$w = 2.44 g$$
11. Given :

$$(i) C(graphite) + O_{2}(g) \rightarrow CO_{2}(g);$$

$$\Delta rH^{\circ} = x kJ mol^{-1}$$

$$(ii) C(graphite) + \frac{1}{2}O_{2}(g) \rightarrow CO_{2}(g);$$

2



$$\Delta r H^{\circ} = y \ kJ \ mol^{-1}$$
(iii) $CO(g) + \frac{1}{2}O_{2}(g) \rightarrow CO_{2}(g)$;

$$\Delta r H^{\circ} = z \ kJ \ mol^{-1}$$
Based on the above thermochemical equations find out which one of the following algebraic relationships is correct ?
(1) $z = x + y$ (2) $x = y - z$
(3) $x = y + z$ (4) $y = 2z - x$
Ans. (3)
Sol.
 $C_{(graphite)} + O_{2}(g) \rightarrow CO_{2}(g)\Delta_{r}H^{\circ} = xkJ/mol...(1)$
 $C_{(graphite)} + \frac{1}{2}O_{2}(g) \rightarrow CO_{2}(g)\Delta_{r}H^{\circ} = ykJ/mol...(2)$
 $CO(g) + \frac{1}{2}O_{2}(g) \rightarrow CO_{2}(g)\Delta_{r}H^{\circ} = zkJ/mol...(3)$
(1) $= (2) + (3)$
 $x = y + z$

12. An open vessel at 27°C is heated until two fifth of the air (assumed as an ideal gas) in it has escaped from the vessel. Assuming that the volume of the vessel remains constant, the temperature at which the vessel has been heated is :

(1) 750°C	(2) 500°C
(3) 750 K	(4) 500 K

Ans. (4)

 $\frac{2}{5}$ air escaped from vessel, $\therefore \frac{3}{5}$ air remain Sol. is vessel. P. V constant

$$n_1 T_1 = n_2 T_2$$

$$n_1(300) = \left(\frac{3}{5}n_1\right) T_2 \Longrightarrow T_2 = 500 \text{ K}$$

 $\wedge^{\circ}_{\rm m}$ for NaCl, HCl and NaA are 126.4, 425.9 13. and 100.5 S cm²mol⁻¹, respectively. If the conductivity of 0.001 M HA is 5×10^{-5} S cm⁻¹, degree of dissociation of HA is : (2) 0.125 (3) 0.25(1) 0.75(4) 0.50

Ans. (2)

Sol.

Ε

$$\Lambda_{\rm m}^{0}({\rm HA}) = \Lambda_{\rm m}^{0}({\rm HCl}) + \Lambda_{\rm m}^{0}({\rm NaA}) - \Lambda_{\rm m}^{0}({\rm NaCl})$$

= 425.9 + 100.5 - 126.4

 $= 400 \text{ S cm}^2 \text{ mol}^{-1}$ $\Lambda_{\rm m} = \frac{1000 \text{K}}{\text{M}} = \frac{1000 \times 5 \times 10^{-5}}{10^{-3}} = 50 \text{ S } \text{cm}^2 \text{ mol}^{-1}$ $\alpha = \frac{\Lambda_m}{\Lambda_m^0} = \frac{50}{400} = 0.125$

The major product in the following conversion 14. is :



(4)
$$CH_3O$$
 — CH — CH — CH_2 — CH_2 — CH_3O — I_3O —

Ans. (2) Sol.

(

H₃C-O-
$$\leftarrow$$
CH=CH-CH₃- $\stackrel{\text{HBr}}{\longrightarrow}$ HO- \leftarrow CH-CH₂-CH₃
Hydrolysis
of ether addition
acc. to markonikoff's
Rule

- 15. If K_{sp} of Ag_2CO_3 is 8×10^{-12} , the molar solubility of Ag_2CO_3 in 0.1M AgNO₃ is :
 - (1) 8×10⁻¹² M (2) 8×10⁻¹⁰ M

(3)
$$8 \times 10^{-11} \text{ M}$$
 (4) $8 \times 10^{-13} \text{ M}$

Ans. (2)

Sol.
$$Ag_2CO_3 (s) \rightleftharpoons 2Ag^+(aq.) + CO_3^{-2}(aq)$$

 $(0.1+2S) M \qquad S M$
 $Ksp = [Ag^+]^2[CO_3^{-2}]$
 $8 \times 10^{-12} = (0.1 + 2S)^2 (S)$
 $S = 8 \times 10^{-10} M$



- 16. Among the following, the false statement is :
 - (1) Latex is a colloidal solution of rubber particles which are positively charged
 - (2) Tyndall effect can be used to distingush between a colloidal solution and a true solution.
 - (3) It is possible to cause artificial rain by throwing electrified sand carrying charge opposite to the one on clouds from an aeroplane.
 - (4) Lyophilic sol can be coagulated by adding an electrolyte.
- Ans. (1)
- Sol. Colloidal solution fo rubber are negatively charged.
- 17. The pair that does NOT require calcination is:
 - (1) ZnO and MgO
 - (2) Fe₂O₃ and CaCO₃.MgCO₃
 - (3) ZnO and Fe₂O₃.xH₂O
 - (4) ZnCO₃ and CaO
- Ans. (1)
- Sol. ZnO & MgO both are in oxide form therefore no change on calcination.
- 18. The correct order of atomic radii is :
 - (1) Ce > Eu > Ho > N (2) N > Ce > Eu > Ho
 - (3) Eu > Ce > Ho > N (4) Ho > N > Eu > Ce
- Ans. (3)



19. The element that does NOT show catenation is:

(1) Sn (2) Ge (3) Si (4) Pb

Ans. (4)

- Sol. Catenation is not shown by lead.
- 20. The combination of plots which does not represent isothermal expansion of an ideal gas is:







- (1) (A) and (C) (2) (A) and (D) (4) (B) and (C)
- (3) (B) and (D)

Ans. (3)

Isothermal expansion $PV_m = K(Graph-C)$ Sol.

$$\mathbf{P} = \frac{\mathbf{K}}{\mathbf{V}_{\mathrm{m}}} \quad (\text{Graph-A})$$



21. The volume strength of $1M H_2O_2$ is:

- (Molar mass of $H_2O_2 = 34$ g mol⁻¹)
- (1) 16.8(2) 11.35

Ans. (2)

 $1L - 1M H_2O_2$ solution will produce 11.35 Sol. $L O_2$ gas at STP.





22. For a reaction consider the plot of ln k versus 1/T given in the figure. If the rate constant of this reaction at 400 K is 10^{-5} s⁻¹, then the rate constant at 500 K is :



Ans. (2)

Sol.

$$2.303 \log \frac{K_2}{10^{-5}} = 4606 \left[\frac{1}{400} - \frac{1}{500} \right]$$
$$\Rightarrow K_2 = 10^{-4} \text{ s}^{-1}$$

 $ln\frac{K_2}{K_1} = \frac{E_a}{R} \left[\frac{1}{T_1} - \frac{1}{T_2} \right]$

23. The element that shows greater ability to form $p\pi$ - $p\pi$ multiple bonds, is : С

(1) Si (2) Ge (3) Sn (4)
$$($$

Ans. (4)

- Sol. carbon atom have 2p orbitals able to form strongest $p\pi - p\pi$ bonds
- 24. The major product of the following reaction is: H₂C

(i) $NaNO_2/H^+$ (ii) $CrCO_3/H^+$ (iii) H_2SO_4 (conc.), Δ



Ans. (2)

Ε



25. The aldehydes which will not form Grignard product with one equivalent Grignard reagents are :

(A)
$$(B)$$
 (B) (C) (B) (C) (B) (C) (C) (C) (C)

Ans. (2)

Sol. Acid-base reaction of G.R are fast.







26. The major product of the following reaction is:





Ans. (1)



- **27.** Chlorine on reaction with hot and concentrated sodium hydroxide gives :
 - (1) Cl^- and ClO_2^- (2) Cl^- and ClO_3^- (3) Cl^- and ClO^- (4) ClO_3^- and ClO_2^-

Ans. (2)

Sol. $3Cl_2 + 6 OH^- \rightarrow 5Cl^- + ClO_3^- + 3H_2O$

28. The major product of the following reaction is:

 $\begin{array}{c} CH_{3}CH_{2}CH-CH_{2} & \xrightarrow{(i) \text{ KOH alc.}} \\ Br & Br & Br & \xrightarrow{(ii) \text{ NaNH}_{2}} \\ in \text{ liq NH}_{3} \end{array}$ $(1) CH_{3}CH_{2}C=CH$ $(2) CH_{3}CH_{2}CH-CH_{2} \\ I & I \\ NH_{2} NH_{2} \end{array}$ $(3) CH_{3}CH=C=CH_{2}$ $(4) CH_{3}CH=CHCH_{2}NH_{2}$ Ans. (1)

CH₃ - CH₂ - CH - CH₂
Br Br

$$\downarrow$$
 Alc. KOH
CH₃ - CH₂ - C = CH₂
Sol.
NaNH₂ \downarrow in liq. NH₃
CH₃ - CH₂ - C = CH

29. If the de Broglie wavelength of the electron in n^{th} Bohr orbit in a hydrogenic atom is equal to 1.5 $\pi a_0(a_0$ is Bohr radius), then the value of n/z is :

Sol. According to de-broglie's hypothesis

$$2\pi r_n = n\lambda \implies 2\pi \cdot a_0 = \frac{n^2}{z} = n \times 1.5\pi a_0$$

 $\frac{n}{z} = 0.75$

- **30.** The two monomers for the synthesis of Nylone 6, 6 are :
 - (1) HOOC(CH_2)₆COOH, $H_2N(CH_2)_6NH_2$
 - (2) HOOC(CH_2)₄COOH, $H_2N(CH_2)_4NH_2$
 - (3) HOOC(CH_2)₆COOH, $H_2N(CH_2)_4NH_2$
 - (4) HOOC(CH₂)₄COOH, H₂N(CH₂)₆NH₂

Ans. (4)

Sol. Nylon-6,6 is polymer of Hexamethylene diamine & Adipic acid \downarrow \downarrow \downarrow H₂N-(CH₂)₆-NH₂ HOOC-(CH₂)₄-COOH

6



Answers & Solutions

for

JEE (MAIN)-2019 (Online) Phase-2

(Physics, Chemistry and Mathematics)

Time : 3 hrs.

M.M.: 360

Important Instructions :

- 1. The test is of **3 hours** duration.
- 2. The Test Booklet consists of **90** questions. The maximum marks are **360**.
- 3. There are *three* parts in the question paper A, B, C consisting of **Physics**, **Chemistry** and **Mathematics** having 30 questions in each part of equal weightage.
- 4. Each question is allotted 4 (four) marks for each correct response. ¼ (one-fourth) marks will be deducted for indicating incorrect response of each question. No deduction from the total score will be made if no response is indicated for an item in the answer sheet.
- 5. There is only one correct response for each question.


Two particles move at right angle to each other. 1. Their de Broglie wavelengths are λ_1 and λ_2 respectively. The particles suffer perfectly inelastic collision. The de Broglie wavelength λ , of the final particle, is given by:

(1)
$$\lambda = \sqrt{\lambda_1 \lambda_2}$$

(2) $\lambda = \frac{\lambda_1 + \lambda_2}{2}$
(3) $\frac{2}{\lambda} = \frac{1}{\lambda_1} + \frac{1}{\lambda_2}$
(4) $\frac{1}{\lambda^2} = \frac{1}{\lambda_1^2} + \frac{1}{\lambda_2^2}$

Answer (4)

Sol.

$$p_{1} = \frac{h}{\lambda_{1}}$$

$$p_{2} = \frac{h}{\lambda_{2}}$$

$$\therefore \quad p_{f} = \sqrt{p_{1}^{2} + p_{2}^{2}}$$

$$\Rightarrow \quad \frac{h}{\lambda} = \sqrt{\frac{h^{2}}{\lambda_{1}^{2}} + \frac{h^{2}}{\lambda_{2}^{2}}}$$

$$\Rightarrow \quad \frac{1}{\lambda^{2}} = \frac{1}{\lambda_{1}^{2}} + \frac{1}{\lambda_{2}^{2}}$$

2. In SI units, the dimensions of
$$\sqrt{\frac{\epsilon_0}{\mu_0}}$$
 is:

(1) A	T ² M ⁻¹ L ⁻¹	(2)	AT-3ML ^{3/2}
(3) A	⁻¹ TML ³	(4)	$A^{2}T^{3}M^{-1}L^{-2}$

Answer (4)

Sol.
$$\left[\sqrt{\frac{\epsilon_0}{\mu_0}}\right] = \left[\frac{\epsilon_0}{\sqrt{\mu_0 \epsilon_0}}\right] = [LT^{-1}] \times [\epsilon_0]$$
$$\therefore \quad \mathbf{F} = \frac{\mathbf{q}^2}{4\pi \epsilon_0 \mathbf{r}^2}$$
$$\Rightarrow \quad [\epsilon_0] = \frac{[AT]^2}{[MLT^{-2}] \times [L^2]}$$
$$\therefore \quad \left[\sqrt{\frac{\epsilon_0}{\mu_0}}\right] = [LT^{-1}] \times [A^2 \mathbf{M}^{-1} \mathbf{L}^{-3} \mathbf{T}^4]$$
$$= [\mathbf{M}^{-1} \mathbf{L}^{-2} \mathbf{T}^3 \mathbf{A}^2]$$

A particle moves in one dimension from rest 3. under the influence of a force that varies with the distance travelled by the particle as shown in the figure. The kinetic energy of the particle after it has travelled 3 m is



(3) 5 J

Answer (4)

Sol. = Area under F-x graph

$$\Delta K.E = W = \frac{1}{2} \times (3+2) \times (3-2) + 2 \times 2$$

= 2.5 + 4
= 6.5 J

- 4. Radiation coming from transitions n = 2 to n = 1 of hydrogen atoms fall on He^+ ions in n = 1 and n = 2 states. The possible transition of helium ions as they absorb energy from the radiation is
 - (1) $n = 2 \rightarrow n = 4$
 - (2) $n = 2 \rightarrow n = 5$
 - (3) $n = 2 \rightarrow n = 3$
 - (4) $n = 1 \rightarrow n = 4$

Answer (1)

Sol. Energy released by hydrogen atom

$$\Delta E_1 = 13.6 \times \left(\frac{1}{1} - \frac{1}{4}\right) = \frac{3}{4} \times 13.6 \text{ eV}$$

= 10.2 eV

Also, energy absorbed by H_e^+ ion in transition $n = 2 \rightarrow n = 4$

$$\Delta E_2 = 13.6 \times 4 \times \left(\frac{1}{4} - \frac{1}{16}\right) = 10.2 \text{ eV}$$

So, possible transition is $n = 2 \rightarrow n = 4$



is 5 cm, the Reynolds number for the flow is of the order of : (density of water = 1000 kg/m^3 , coefficient of viscosity of water = 1 mPa s)

(1) 10 ²	(2) 10 ⁴
---------------------	---------------------

(3) 10^3 (4) 10^6

Answer (2)

Sol. Flow rate of water (Q) = 100 lit/min

$$= \frac{100 \times 10^{-3}}{60} = \frac{5}{3} \times 10^{-3} \text{ m}^{3}$$

$$\therefore \quad \text{Velocity of flow (v)} = \frac{Q}{A} = \frac{5 \times 10^{-3}}{3 \times \pi \times (5 \times 10^{-2})^{2}}$$

$$= \frac{10}{15\pi} = \frac{2}{3\pi} \text{ m/s}$$

$$= 0.2 \text{ m/s}$$

$$\therefore \quad \text{Reynold number } (R_{e}) = \frac{Dv\rho}{\eta}$$

$$= \frac{(10 \times 10^{-2}) \times \frac{2}{3\pi} \times 1000}{2 \times 10^{4}} \approx 2 \times 10^{4}$$

$$=\frac{(10\times10^{-2})\times\frac{2}{3\pi}\times1000}{1} \simeq 2\times10^{4}$$

Order of R_e = 10⁴

6. In figure, the optical fiber is I = 2 m long and has a diameter of $d = 20 \mu \text{m}$. If a ray of light is incident on one end of the fiber at angle $\theta_1 = 40^\circ$, the number of reflections it makes before emerging from the other end is close to:

(refractive index of fiber is 1.31 and $\sin 40^\circ = 0.64$)



Answer (4)

Sol. $1 \times \sin 40^{\circ} = 1.31 \sin \theta$

$$\Rightarrow \sin\theta = \frac{0.64}{1.31} \Rightarrow \theta \approx 30^{\circ}$$
$$I = 20 \ \mu m \times \cot\theta$$
$$\therefore \quad N = \frac{2}{20 \times 10^{-6} \times \cot\theta}$$
$$= \frac{2 \times 10^{6}}{20 \times \sqrt{3}} = 57735$$

code, the new resistance will be:

- **(1) 400** Ω
- **(2) 500** Ω
- **(3) 300** Ω
- **(4) 100** Ω

Answer (2)

Sol. 200 Ω = Red + Black + Brown

 $\textbf{Green} \equiv \textbf{5}$

So, Green + Black + Brown = 500 Ω

8. A 20 Henry inductor coil is connected to a 10 ohm resistance in series as shown in figure. The time at which rate of dissipation of energy (Joule's heat) across resistance is equal to the rate at which magnetic energy is stored in the inductor, is:





the corners of a square. They have accelerations of equal magnitude with directions as shown. The acceleration of the centre of mass of the particles is :



Answer (4)

Sol.
$$a_{CM} = \frac{(2m)a\hat{j} + 3m \times a\hat{i} + ma(-\hat{i}) + 4m \times a(-\hat{j})}{2m + 3m + 4m + m}$$
$$= \frac{2a\hat{i} - 2a\hat{j}}{10} = \frac{a}{5}(\hat{i} - \hat{j})$$

- (1) 2×10^{-8} T along y-direction
- (2) 6×10^{-8} T along z-direction
- (3) 2×10^{-8} T along z-direction
- (4) 6×10^{-8} T along x-direction

Answer (3)

Sol.
$$B_0 = \frac{E_0}{c} = \frac{6}{3 \times 10^8} = 2 \times 10^{-8} T$$

Propagation direction = $\hat{E} \times \hat{B}$

$$\hat{\mathbf{i}} = \hat{\mathbf{j}} \times \hat{\mathbf{B}}$$

 $\Rightarrow \hat{\mathbf{B}} = \hat{\mathbf{k}}$

11. Voltage rating of a parallel plate capacitor is 500 V. Its dielectric can withstand a maximum electric field of 10⁶ V/m. The plate area is 10⁻⁴ m². What is the dielectric constant if the capacitance is 15 pF?

(given $\varepsilon_0 = 8.86 \times 10^{-12} \text{ C}^2/\text{Nm}^2$)		
(1) 3.8	(2) 4.5	
(3) 8.5	(4) 6.2	

Sol.
$$C = \frac{k \in 0}{d}$$

 $E = \frac{V}{d}$
 $15 \times 10^{-12} = \frac{k \times 8.86 \times 10^{-12} \times 10^{-4} \times 10^{6}}{500}$

k = 8.5

- 12. The wavelength of the carrier waves in a modern optical fiber communication network is close to :
 - (1) 600 nm (2) 900 nm
 - (3) 1500 nm (4) 2400 nm

Answer (3)

Sol. Fact Based

Wavelength of carrier waves in modern optical fiber communication is most widely used near about 1500 nm.

- 13. A steel wire having a radius of 2.0 mm, carrying a load of 4 kg, is hanging from a ceiling. Given that $g = 3.1 \pi \text{ ms}^{-2}$, what will be the tensile stress that would be developed in the wire?
 - (1) $4.8 \times 10^{6} \text{ Nm}^{-2}$
 - (2) $3.1 \times 10^6 \text{ Nm}^{-2}$
 - (3) $5.2 \times 10^{6} \text{ Nm}^{-2}$
 - (4) $6.2 \times 10^6 \text{ Nm}^{-2}$

Answer (2)

В

Sol. Stress =
$$\frac{F}{A} = \frac{4 \times 3.1\pi}{\pi \times (2 \times 10^{-3})^2} = 3.1 \times 10^6 \text{ N/m}^2$$

14. A thin circular plate of mass M and radius R has its density varying as $\rho(\mathbf{r}) = \rho_0 \mathbf{r}$ with ρ_0 as constant and r is the distance from its center. The moment of inertia of the circular plate about an axis perpendicular to the plate and passing through its edge is $I = a MR^2$. The value of the coefficient a is

(1)
$$\frac{3}{2}$$
 (2) $\frac{1}{2}$

(3)
$$\frac{3}{5}$$
 (4) $\frac{8}{5}$



$$J^{1} = \int_{0}^{R} \rho_{0} \mathbf{r} \times 2\pi \mathbf{r} d\mathbf{r} \times \mathbf{r}^{2} = \frac{2\pi\rho_{0}R^{5}}{5}$$

$$\therefore I = I_{C} + MR^{2} = 2\pi\rho_{0}R^{5} \left(\frac{1}{3} + \frac{1}{5}\right) = \frac{16\pi\rho_{0}R^{5}}{15}$$
$$= \frac{8}{5} \left[\frac{2}{3}\pi\rho_{0}R^{3}\right]R^{2} = \frac{8}{5}MR^{2}$$

15. An alternating voltage v(t) = 220 sin 100π t volt is applied to a purely resistive load of 50 Ω . The time taken for the current to rise from half of the peak value to the peak value is

(1)	2.2 ms	(2)	7.2 ms
(3)	5 ms	(4)	3.3 ms

Answer (4)

Sol. I = I_m sin(100 π t)

$$\Rightarrow \frac{l_m}{2} = l_m \sin(100\pi t_1)$$
$$\Rightarrow \frac{\pi}{6} = 100\pi t_1$$

$$\Rightarrow$$
 $t_1 = \frac{1}{600} s$

$$T = \frac{2\pi}{100\pi} = \frac{1}{50} s$$

$$\therefore t_{req} = \frac{T}{4} - t_1 = \frac{1}{200} - \frac{1}{600} = \frac{2}{600} = \frac{1}{300} s = 3.3 \, \text{ms}$$

16. Four identical particles of mass M are located at the corners of a square of side 'a'. What should be their speed if each of them revolves under the influence of others' gravitational field in a circular orbit circumscribing the square?





17. The bob of a simple pendulum has mass 2 g and a charge of 5.0 μ C. It is at rest in a uniform horizontal electric field of intensity 2000 V/m. At equilibrium, the angle that the pendulum makes with the vertical is

$$(take g = 10 m/s^2)$$

(1)
$$\tan^{-1}(0.2)$$
 (2) $\tan^{-1}(5.0)$

(3)
$$\tan^{-1}(2.0)$$
 (4) $\tan^{-1}(0.5)$

Answer (4)

Sol.
$$T\cos\theta = mg$$

 $T\sin\theta = qE$
 $\tan\theta = \frac{qE}{mg}$
 $\tan\theta = \frac{5 \times 10^{-6} \times 2000}{2 \times 10^{-3} \times 10} = \frac{1}{2}$
 $\Rightarrow \tan^{-1}\left(\frac{1}{2}\right)$
18. $\frac{A}{1}$

A wire of length 2L, is made by joining two
wires A and B of same length but different radii
r and 2r and made of the same material. It is
vibrating at a frequency such that the joint of
the two wires forms a node. If the number of
antinodes in wire A is p and that in B is q then
the ratio
$$p : q$$
 is

(1) 4:9	(2) 1:2

(3) 3:5 (4) 1:4

Answer (2)



$$\begin{array}{cccc} & & & & \\ & & & \\ & \Rightarrow & \lambda_{A} = 2\lambda_{B} \\ & \Rightarrow & & \frac{p}{q} = \frac{1}{2} \end{array}$$

19. The reverse breakdown voltage of a Zener diode is 5.6 V in the given circuit.



The current Iz through the Zener is

(1) 15 mA	(2) 7 mA
(3) 10 mA	(4) 17 mA

Answer (3)

$$I_{800 \Omega} = \frac{5.6}{800} A = 7 \text{ mA}$$

 $I_{200 \Omega} = \frac{9-5.6}{200} = 17 \text{ mA}$
 $\therefore I_z = 17 - 7 = 10 \text{ mA}$

20. A thin strip 10 cm long is on a U shaped wire of negligible resistance and it is connected to a spring of spring constant 0.5 Nm⁻¹(see figure). The assembly is kept in a uniform magnetic field of 0.1 T. If the strip is pulled from its equilibrium position and released, the number of oscillations it performs before its amplitude decreases by a factor of e is N. If the mass of the strip is 50 grams, its resistance 10 Ω and air drag negligible, N will be close to

X X X X X X X X B X X X X X X X X X X X X	10 cm	
(1) 1000	(2)	5000
(3) 50000	(4)	10000
Answer (2)		

$$\mathbf{F} = -\mathbf{k}\mathbf{x} - \frac{\mathbf{B}^2\mathbf{I}^2}{\mathbf{R}} \times \mathbf{v}$$

So, it is case of damped oscillation

$$\Rightarrow A = A_0 e^{-\frac{bt}{2m}}$$

$$\Rightarrow \frac{A_0}{e} = A_0 e^{-\frac{bt}{2m}}$$

$$\Rightarrow t = \frac{2m}{\left(\frac{B^2l^2}{R}\right)} = \frac{2 \times 50 \times 10^{-3} \times 10}{0.01 \times 0.01} = 10000 \text{ s}$$
Time period, $T = 2\pi \sqrt{\frac{m}{k}} \approx 2 \text{ s}$

$$\therefore \text{ Number, } N = \frac{10000}{2} = 5000$$

21. Ship A is sailing towards north-east with velocity $\vec{v} = 30\hat{i} + 50\hat{j}$ km/hr where \hat{i} points east and \hat{j} , north. Ship B is at a distance of 80 km east and 150 km north of Ship A and is sailing towards west at 10 km/hr. A will be at minimum distance from B in:

(4) 2.6 hrs.

- (1) 2.2 hrs. (2) 4.2 hrs.
- (3) 3.2 hrs.

Sol.

$$\vec{j}(North)$$

$$\vec{v}_{BA} = \vec{v}_{BA}$$

$$\vec{v}_{A} = 30i + 50j \text{ km / hr}$$

$$\vec{v}_{BA} = (80\hat{i} + 150\hat{j}) \text{ km}$$

$$\vec{v}_{B} = (-10i) \text{ km / hr}$$

$$\vec{v}_{BA} = \vec{v}_{B} - \vec{v}_{A} = -10\hat{i} - 30\hat{i} - 50\hat{i} = -40\hat{i} - 50\hat{j}$$
Projection of

$$(\vec{r}_{BA}) \text{ on } \vec{v}_{BA} = \frac{(\vec{r}_{BA}) \cdot (\vec{v}_{BA})}{(\vec{v}_{BA})}$$

$$= \frac{(80\hat{i} + 150\hat{j})(-40\hat{i} - 50\hat{j})}{10\sqrt{41}} = \frac{\sqrt{mk}}{qB}$$

$$\therefore t = \frac{10 \times 107}{\sqrt{41} \times 10\sqrt{41}} = \frac{107}{41} = 2.6 \text{ Hrs}$$



potential difference between the points 'a' and 'b' is approximately (in V)



Answer (4)

Sol.
$$v = \frac{E_1 r_2 r_3 + E_2 r_3 r_1 + E_3 r_1 r_2}{r_1 r_2 + r_2 r_3 + r_3 r_1}$$

= $\frac{2(2 \times 2) + 4(2 \times 2) + 4(2 \times 2)}{4 + 4 + 4}$
= $\frac{40}{12} = \frac{10}{3} = 3.3$ Volt

23. Two identical beakers A and B contain equal volumes of two different liquids at 60°C each and left to cool down. Liquid in A has density of 8×10^2 kg/m³ and specific heat of 2000 Jkg⁻¹K⁻¹ while liquid in B has density of 10³ kgm⁻³ and specific heat of 4000 Jkg⁻¹K⁻¹. Which of the following best describes their temperature versus time graph schematically? (assume the emissivity of both the beakers to be the same)



Answer (2)

Sol.
$$mS\left(-\frac{dT}{dt}\right) = e\sigma AT^{4}$$
$$-\frac{dT}{dt} = \frac{e\sigma \times A \times T^{4}}{\rho \times Vol. \times S}$$
$$\frac{\left(-\frac{dT}{dt}\right)_{A}}{\left(-\frac{dT}{dt}\right)_{B}} = \frac{\rho_{B}}{\rho_{A}} \times (2) > 1$$

hollow spherical shell. Let the potential difference between the surface of the solid sphere and that of the outer surface of the hollow shell be V. If the shell is now given a charge of -4 Q, the new potential difference between the same two surfaces is:

(1) –2 V	(2) V	
(3) 2 V	(4) 4 V	

Answer (2)

Sol. In case-1

Electric field between spherical surface





Electric field between surfaces remain unchanged.



... Potential difference between them remain unchanged too.

25. A circular coil having N turns and radius r carries a current I. It is held in the XZ plane in a magnetic field $B\hat{i}$. The torque on the coil due to the magnetic field is:

(1)
$$\frac{B\pi r^2 I}{N}$$
(2)
$$\frac{Br^2 I}{\pi N}$$
(3)
$$B\pi r^2 I N$$
(4) Zero
Answer (3)
Sol.
$$|\vec{\tau}| = |\vec{\mu} \times \vec{B}|$$

$$= NIA \times B$$

$$\Rightarrow \tau = NI\pi r^2 B$$



So, A cools down at faster rate.

length 20 cm. A convergent mirror of focal length 10 cm is placed at a distance of 60 cm on the other side of the lens. The position and size of the final image will be:

- (1) 20 cm from the convergent mirror, twice the size of the object
- (2) 20 cm from the convergent mirror, same size as the object
- (3) 40 cm from the convergent lens, twice the size of the object
- (4) 40 cm from the convergent mirror, same size as the object

Answer (Bonus)

Sol.
$$v_1 = \frac{40 \times 20}{(40 - 20)} = 40 \text{ cm}$$

 $u_2 = 60 - 40 = 20 \text{ cm}$
 $\therefore \quad v_2 = \frac{20 \times 10}{(20 - 10)} = 20 \text{ cm}$

- ∴ Image traces back to object itself as image formed by lens is a centre of curvature of mirror.
- 27. If 10^{22} gas molecules each of mass 10^{-26} kg collide with a surface (perpendicular to it) elastically per second over an area 1 m² with a speed 10^4 m/s, the pressure exerted by the gas molecules will be of the order of:
 - (1) 10^8 N/m^2 (2) 10^3 N/m^2
 - (3) 10^{16} N/m^2 (4) 10^4 N/m^2

Answer (Bonus)

Sol. $P = \frac{N \times (2mv)}{\Delta t \times A}$

=

$$=\frac{10^{22} \times 2 \times 10^{-26} \times 10^4}{1 \times 1} = 2 \text{ N/m}^2$$

28. A thermally insulated vessel contains 150 g of water at 0°C. Then the air from the vessel is pumped out adiabatically. A fraction of water turns into ice and the rest evaporates at 0°C itself. The mass of evaporated water will be closest to:

(Latent heat of vaporization of water = 2.10×10^{6} J kg⁻¹ and Latent heat of Fusion of water = 3.36×10^{5} J kg⁻¹)

$$(0) = 20 g$$
 $(4) = 00 g$

Answer (3)

Sol. Let amount of water evaporated be m gram.

$$\therefore m \times L_v = (150 - m) \times L_s$$
$$m \times 540 = (150 - m) \times 80$$
$$\Rightarrow m \simeq 20 g$$

29. In an interference experiment the ratio of

amplitudes of coherent waves is $\frac{a_1}{a_2} = \frac{1}{3}$. The

ratio of maximum and minimum intensities of fringes will be:

Answer (1)

Sol.
$$\frac{a_1}{a_2} = \frac{3}{1}$$

 $\therefore \quad \frac{I_{max}}{I_{min}} = \left(\frac{a_2 + a_1}{a_2 - a_1}\right)^2 = \left(\frac{3 + 1}{3 - 1}\right)^2$

30. A boy's catapult is made of rubber cord which is 42 cm long, with 6 mm diameter of crosssection and of negligible mass. The boy keeps a stone weighing 0.02 kg on it and stretches the cord by 20 cm by applying a constant force. When released, the stone flies off with a velocity of 20 ms⁻¹. Neglect the change in the area of cross-section of the cord while stretched. The Young's modulus of rubber is closest to:

(1)
$$10^4 \text{ Nm}^{-2}$$
 (2) 10^3 Nm^{-2}

(3)
$$10^8 \text{ Nm}^{-2}$$
 (4) 10^6 Nm^{-2}

Sol.
$$\frac{1}{2} \cdot \left(\frac{YA}{L}\right) (\Delta I)^2 = \frac{1}{2} mv^2$$
$$\Rightarrow \quad Y = \frac{mv^2 L}{A(\Delta I)^2}$$
$$= \frac{0.02 \times 400 \times 0.42 \times 4}{\pi \times 36 \times 10^{-6} \times 0.04}$$
$$= 2.3 \times 10^6 \text{ N/m}^2$$
So, order is 10⁶.





(1) Zone refining

- (2) Vapour phase refining
- (3) Thermal reduction
- (4) Electrolysis

Answer (3)

- Sol. Ellingham diagram is used to select reducing agent so it help to predict feasibility of its thermal reduction.
- 2. Which one of the following equations does not correctly represent the first law of thermodynamics for the given processes involving an ideal gas? (Assume non-expansion work is zero)
 - (1) Isothermal process : q = -w
 - (2) Cyclic process : q = -w
 - (3) Isochoric process : $\Delta U = q$
 - (4) Adiabatic process : $\Delta U = -w$

Answer (4)

Sol. ∆U = q + W

Adiabatic process q = 0

 $\Delta U = W$

For isothermal, $\Delta U = 0$

For cyclic, $\Delta U = 0$

For isochoric, W = 0

3. The quantum number of four electrons are given below:

(I)
$$n = 4$$
, $l = 2$, $m_l = -2$, $m_s = -\frac{1}{2}$
(II) $n = 3$, $l = 2$, $m_l = 1$, $m_s = +\frac{1}{2}$
(III) $n = 4$, $l = 1$, $m_l = 0$, $m_s = +\frac{1}{2}$
(IV) $n = 3$, $l = 1$, $m_l = 1$, $m_s = -\frac{1}{2}$

The correct order of their increasing energies will be:

n + l

(1) IV < II < III < I
(2) I < III < I < IV
(3) IV < III < I < I
(4) I < I < III < IV

Answer (1)

Sol.

(I) n = 4 l = 2 4d 6 (II) n = 3 l = 2 3d 5 (III) n = 4 l = 1 4p 5 (IV) n = 3 l = 1 3p 4 more is n + l value, more is energy

3p < 3d < 4p < 4d

4. The major product of the following reaction is :















- 5. The correct order of the spin-only magnetic moment of metal ions in the following low-spin complexes, $[V(CN)_6]^{4-}$, $[Fe(CN)_6]^{4-}$, $[Ru(NH_3)_6]^{3+}$ and $[Cr(NH_3)_6]^{2+}$, is:
 - (1) $V^{2+} > Cr^{2+} > Ru^{3+} > Fe^{2+}$
 - (2) Cr²⁺ > V²⁺ > Ru³⁺ > Fe²⁺
 - (3) $V^{2+} > Ru^{3+} > Cr^{2+} > Fe^{2+}$
 - (4) $Cr^{2+} > Ru^{3+} > Fe^{2+} > V^{2+}$

Answer (1)

Sol.

No. of unpaired electrons

[V(CN) ₆] ^{4–}	V ⁺²	3
[Ru(NH ₃) ₆] ³⁺	Ru ⁺³	1
[Fe(CN) ₆] ^{4_}	Fe ⁺²	0
[Cr(NH ₃) ₆] ²⁺	Cr ⁺²	2

 \therefore Order of spin magnetic moment

 $V^{2+} > Cr^{2+} > Ru^{3+} > Fe^{2+}$

- 6. Which is wrong with respect to our responsibility as a human being to protect our environment?
 - (1) Using plastic bags
 - (2) Restricting the use of vehicles
 - (3) Avoiding the use of floodlighted facilities
 - (4) Setting up compost tin in gardens

Answer (1)

Sol. Use of plastic bags is hazardous to our environment

adsorbed on mass m of the adsorbent. The plot of log $\frac{x}{m}$ versus log p is shown in the given graph. $\frac{x}{m}$ is proportional to



(1)
$$p^{3/2}$$
 (2) p^3
(3) $p^{2/3}$ (4) p^2

Answer (3)

Sol.
$$\frac{x}{m} \propto p^{\frac{1}{n}}$$
 $\frac{x}{m} = kp^{\frac{1}{n}}$
Slope = $\frac{2}{3}$
 $\log \frac{x}{m} = \log k + \frac{1}{n} \log p$
Slope = $\frac{1}{n} = \frac{2}{3}$
 $\frac{x}{m} \propto p^{\frac{2}{3}}$

8. Given that $E_{O_2/H_2O}^{\Theta} = +1.23 \text{ V};$ $E_{S_2O_8^{D^-}/SO_4^{D^-}}^{\Theta} = 2.05 \text{ V}$ $E_{Br_2/Br^-}^{\Theta} = +1.09 \text{ V};$ $E_{Au^{3^+}/Au}^{\Theta} = +1.4 \text{ V}$

The strongest oxidizing agent is

- (1) Br₂
- (2) Au³⁺
- (3) $S_2O_8^{2-}$

(4) O₂

Answer (3)

Sol. More positive is the reduction potential stronger is the oxidising agent.

Reduction potential is maximum for S₂O₈²⁻







- 10. Maltose on treatment with dilute HCI gives
 - (1) D-Galactose
 - (2) D-Glucose and D-Fructose
 - (3) D-Glucose
 - (4) D-Fructose

- is composed of two α -D glucose units.
- 11. The vapour pressures of pure liquids A and B are 400 and 600 mmHg, respectively at 298 K. On mixing the two liquids, the sum of their initial volumes is equal to the volume of the final mixture. The mole fraction of liquid B is 0.5 in the mixture. The vapour pressure of the final solution, the mole fractions of components A and B in vapour phase, respectively are :
 - (1) 500 mmHg, 0.4, 0.6
 - (2) 500 mmHg, 0.5, 0.5
 - (3) 450 mmHg, 0.4, 0.6
 - (4) 450 mmHg, 0.5, 0.5

Answer (1)

Sol. P =
$$x_B p_B^\circ + x_A p_A^\circ$$

$$= 0.5 \times 600 + 0.5 \times 400 = 300 + 200 = 500$$

$$p_B = y_B P_{Total}$$

$$y_{\rm B} = \frac{p_{\rm B}}{P_{\rm Total}} = \frac{300}{500} = \frac{3}{5} = 0.6$$

$$y_A = \frac{p_A}{P_{Total}} = \frac{200}{500} = \frac{2}{5} = 0.4$$

12. For the reaction $2A + B \rightarrow C$, the values of initial rate at different reactant concentrations are given in the table below. The rate law for the reaction is

$[\Lambda](mol l^{-1})$	$[B](mol l^{-1})$	Initial Rate
		(mol $L^{-1}s^{-1}$)
0.05	0.05	0.045
0.10	0.05	0.090
0.20	0.10	0.72

(1) Rate =
$$k[A]^{2}[B]^{2}$$
 (2) Rate = $k[A][B]$

(4) Rate = $k[A][B]^2$

Sol. 2A + B
$$\longrightarrow$$
 P
Rate = k[A]^x [B]^y
Exp-1, 0.045 = k[0.05]^x [0.05]^y ...(i)



Exp-3, $0.72 = k[0.2]^{x} [0.1]^{y} ...(iii)$

Divide equation (i) by equation (ii)

$$\frac{0.045}{0.090} = \left(\frac{1}{2}\right)^{x} \Rightarrow x = 1$$

Divide equation (i) by equation (iii)

$$\frac{0.045}{0.72} = \left(\frac{0.05}{0.1}\right)^{y} \left(\frac{0.05}{0.2}\right)^{1}$$
$$\left(\frac{1}{2}\right)^{2} = \left(\frac{1}{2}\right)^{y} \Rightarrow y = 2$$

Rate law = $k[A]^1 [B]^2$.

13. For silver, $C_p(JK^{-1} \text{ mol}^{-1}) = 23 + 0.01T$. If the temperature (T) of 3 moles of silver is raised from 300 K to 1000 K at 1 atm pressure, the value of ΔH will be close to

(1) 21 kJ	(2) 13 kJ
(3) 62 kJ	(4) 16 kJ

Answer (3)

Sol. n = 3

$$T_{1} = 300$$

$$T_{2} = 1000$$

$$C_{p} = 23 + 0.01T$$

$$\Delta H = \int_{T_{1}}^{T_{2}} nC_{p} dT$$

$$= n \int_{300}^{1000} (23 + 0.01T) dT$$

$$= 3 \left[23T + \frac{0.01T^{2}}{2} \right]_{300}^{1000}$$

$$= 3[16100 + 4550]$$

$$= 3 \times 20650 = 61950 \text{ J}$$

$$= 61.95 \text{ kJ}$$

$$\approx 62$$

(1) Gd ^{or}	(2) Lu st
(3) La ³⁺	(4) Sm ³⁺

Answer (4)

Sol. Sm^{+3} = Partially filled f orbital = $4f^5$

 $Sm = 4f^{6}6s^{2}$

Sm⁺³ = Yellow.

 $Lu^{+3} = 4f^{14}$ colourless.

15. The correct order of hydration enthalpies of alkali metal ions is

(2) Li⁺ > Na⁺ > K⁺ > Cs⁺ > Rb⁺

Answer (4)

Sol. Smaller is size more is hydration energy.

Li⁺ < Na⁺ < K⁺ < Rb⁺ < Cs⁺ Size

$$Li^+ > Na^+ > K^+ > Rb^+ > Cs^+$$
 Hydration energy

16. The major product of the following reaction is





oxygen atoms occupy all the tetrahedral voids. The structure of bimetallic oxide is

- (1) A_2B_2O
- (2) AB₂O₄
- (3) A₄B₂O
- (4) A₂BO₄

Answer (2)

Sol. Lattice formed by B(ccp) = 4

O = tetrahedral voids = 8

Formula = AB_2O_4

18. If solubility product of $Zr_3(PO_4)_4$ is denoted by K_{sp} and its molar solubility is denoted by S, then which of the following relation between S and K_{sp} is correct?

(1)
$$\mathbf{S} = \left(\frac{\mathbf{K}_{sp}}{929}\right)^{\frac{1}{9}}$$
 (2) $\mathbf{S} = \left(\frac{\mathbf{K}_{sp}}{216}\right)^{\frac{1}{7}}$
(3) $\mathbf{S} = \left(\frac{\mathbf{K}_{sp}}{144}\right)^{\frac{1}{6}}$ (4) $\mathbf{S} = \left(\frac{\mathbf{K}_{sp}}{6912}\right)^{\frac{1}{7}}$

Answer (4)

Sol.
$$Zr_{3}(PO_{4})_{4} = 3Zr^{+4} + 4PO_{4}^{3-}$$

 $K_{sp} = \left[Zr^{+4}\right]^{3} \left[PO_{4}^{3-}\right]^{4} = (3S)^{3}(4S)^{4}$
 $K_{sp} = 6912 S^{7}$
 $S = \left(\frac{K_{sp}}{6912}\right)^{1/7}$

- 19. In order to oxidise a mixture of one mole of each of FeC_2O_4 , $Fe_2(C_2O_4)_3$, $FeSO_4$ and $Fe_2(SO_4)_3$ in acidic medium, the number of moles of KMnO₄ required is
 - (1) 1.5
 - (2) 2
 - (3) 3
 - (4) 1

Sol. $5e + MnO_4^- \longrightarrow Mn^{+2}$

(i) $FeC_2O_4 \longrightarrow Fe^{3+} + 2CO_2 + 3e^{3+}$

1 mole of FeC_2O_4 react with $\frac{3}{5}$ moles of acidified KMnO₄

- (ii) $Fe_2(C_2O_4)_3 \longrightarrow Fe^{3+} + CO_2 + 6e$ 1 mole of $Fe_2(C_2O_4)_3$ react with $\frac{6}{5}$ moles of KMnO₄
- (iii) $FeSO_4 \longrightarrow Fe^{3+} + e$
 - 1 mole of $FeSO_4$ react with $\frac{1}{5}$ moles of $KMnO_4$
- $\therefore \text{ Total moles required} = \frac{3}{5} + \frac{6}{5} + \frac{1}{5} = 2$
- 20. 100 mL of a water sample contains 0.81 g of calcium bicarbonate and 0.73 g of magnesium bicarbonate. The hardness of this water sample expressed in terms of equivalents of CaCO₃ is

(molar mass of calcium bicarbonate is 162 g mol^{-1} and magnesium bicarbonate is 146 g mol^{-1})

- (1) 5,000 ppm
- (2) 100 ppm
- (3) 10,000 ppm
- (4) 1,000 ppm

Answer (3)

Sol. Moles of $Ca(HCO_3)_2 = 0.005$

Moles of $Mg(HCO_3)_2 = 0.005$

Hardness in terms of $CaCO_3$ pm

 $=\frac{(0.005+0.005)\times100}{100}\times10^{6}$

= 10⁴ ppm





Answer (1)

Sol. It has four lone pairs but maximum it will be able to donate three lone pairs.

Maximum denticity is 3.

- 22. Diborane (B_2H_6) reacts independently with O_2 and H_2O to produce, respectively:
 - (1) H_3BO_3 and B_2O_3 (2) HBO_2 and H_3BO_3

(3)
$$B_2O_3$$
 and H_3BO_3 (4) B_2O_3 and $[BH_4]^-$

Answer (3)

- Sol. $B_2H_6 + 3O_2 \longrightarrow B_2O_3 + 3H_2O$ $B_2H_6 + 6H_2O \longrightarrow 2H_3BO_3 + 6H_2$
- The size of the iso-electronic species Cl⁻, Ar and Ca²⁺ is affected by
 - (1) Nuclear charge
 - (2) Principal quantum number of valence shell
 - (3) Azimuthal quantum number of valence shell
 - (4) Electron-electron interaction in the outer orbitals

Answer (1)

- **Sol.** Iso-electronic species differ in size due to different effective nuclear charge.
- 24. The IUPAC name of the following compound is:



- (1) 3-Hydroxy-4-methylpentanoic acid
- (2) 4-Methyl-3-hydroxypentanoic acid
- (3) 2-Methyl-3-hydroxypentan-5-oic acid
- (4) 4,4-Dimethyl-3-hydroxybutanoic acid

Answer (1)



- 25. Which of the following amines can be prepared by Gabriel phthalimide reaction?
 - (1) Neo-pentylamine (2) n-butylamine
 - (3) t-butylamine (4) Triethylamine

Answer (2)

Sol. Primary amines are prepared by Gabriel pthallimide synthesis



26. An organic compound 'X' showing the following solubility profile is



Benzamide is amphoteric

OH

Oleic acid will dissolve in NaOH as well as NaHCO $_3$ due to acidic nature.

 \rightarrow Insoluble

NaHCO₃ (10%)





Answer (1)



28. Assertion : Ozone is destroyed by CFCs in the upper stratosphere.

Reason : Ozone holes increase the amount of UV radiation reaching the earth.

- (1) Assertion and reason are both correct, and the reason is the correct explanation for the assertion.
- (2) Assertion is false, but the reason is correct.
- (3) Assertion and reason are correct, but the reason is not the explanation for the assertion.
- (4) Assertion and reason are incorrect.

Answer (3)

Sol. CFC's are responsible for depletion of ozone layer

$$CF_2CI_2 \xrightarrow{uv} CI_{(g)} + F_2CCI_{(g)}$$

$$\dot{\mathbf{CI}} + \mathbf{O}_{3} \longrightarrow \dot{\mathbf{CIO}} + \mathbf{O}_{2}$$

$$\dot{cio} + o \longrightarrow \dot{ci} + o_2$$

Both statements are correct.

Fehling solution. It however, reacts with Grignard reagent and gives positive iodoform test. The compound is





- $\rightarrow\,$ Reacts with Grignard's reagent due to acidic hydrogen.
- $\rightarrow\,$ Fehling solution test is negative as there is no CHO group.
- $\rightarrow~{\rm Neutral~FeCl}_{\rm 3}$ test is negative as there is no phenolic group
- 30. In the following compounds, the decreasing order of basic strength will be
 - (1) $NH_3 > C_2H_5NH_2 > (C_2H_5)_2NH$
 - (2) $C_2H_5NH_2 > NH_3 > (C_2H_5)_2NH_3$
 - (3) $(C_2H_5)_2NH > NH_3 > C_2H_5NH_2$
 - (4) $(C_2H_5)_2NH > C_2H_5NH_2 > NH_3$

Answer (4)

Sol. Correct order of K_b value

 $(C_2H_5)_2 N > (C_2H_5)_3N > NH_3$

In aqueous medium sec. amines are most basic.

3° amines are more basic than NH_3 as +I factor dominate over steric factor.



1. The shortest distance between the line y = xand the curve $y^2 = x - 2$ is :

(1)
$$\frac{11}{4\sqrt{2}}$$
 (2) $\frac{7}{8}$
(3) 2 (4) $\frac{7}{4\sqrt{2}}$

Answer (4)

Sol. The shortest distance between line y = x and parabola = the distance between line y = x and tangent of parabola having slope 1.

Let equation of tangent of parabola having slope 1 is,

$$y = m (x - 2) + \frac{a}{m}$$

where m = 1 and $a = \frac{1}{4}$

Equation of tangent $y = x - \frac{7}{4}$

Distance between the line y = x and the tangent

$$= \left| \frac{\frac{7}{4} - 0}{\sqrt{1^2 + 1^2}} \right| = \frac{7}{4\sqrt{2}}$$

2. The sum of the co-efficients of all even degree terms in x in the expansion of $(x + \sqrt{x^3 - 1})^6$ +

$(x - \sqrt{x^3 - 1})^6$, (x > 1)	is equal to :
(1) 24	(2) 32

(3) 26 (4) 29

Answer (1)

Sol.
$$(x + \sqrt{x^3 - 1})^6 + (x - \sqrt{x^3 - 1})^6$$

= $2 \Big[{}^6C_0 x^6 + {}^6C_2 x^4 (x^3 - 1) + {}^6C_4 x^2 (x^3 - 1)^2 + {}^6C_6 (x^3 - 1)^3 \Big]$
= $2 [x^6 + 15x^7 - 15x^4 + 15x^8 - 30x^5 + 15x^2 + x^9 - 3x^6 + 3x^3 - 1]$

Sum of coefficients of even powers of x = 2[1 - 15 + 15 + 15 - 3 - 1] = 24

3. If S₁ and S₂ are respectively the sets of local minimum and local maximum points of the function, $f(x) = 9x^4 + 12x^3 - 36x^2 + 25$, $x \in R$, then :

(1)
$$\mathbf{S}_1 = \{-2\}; \mathbf{S}_2 = \{0, 1\}$$

(2)
$$S_1 = \{-2, 1\}; S_2 = \{0\}$$

(3)
$$S_1 = \{-1\}; S_2 = \{0, 2\}$$

(4)
$$S_1 = \{-2, 0\}; S_2 = \{1\}$$

Answer (2)

Sol.
$$f(x) = 9x^4 + 12x^3 - 36x^2 + 72$$

$$f'(x) = 36[x^3 + x^2 - 2x] = 36x(x - 1)(x + 2)$$

Whenever derivative changes sign from negative to positive, we get local minima, and whenever derivative changes sign from positive to negative, we get local maxima (while moving left to right on x-axis)

$$S_1 = \{-2, 1\}$$

 $S_2 = \{0\}$

4. If
$$2y = \left(\cot^{-1}\left(\frac{\sqrt{3}\cos x + \sin x}{\cos x - \sqrt{3}\sin x}\right)\right)^2$$
, $x \in \left(0, \frac{\pi}{2}\right)$ then $\frac{dy}{dx}$ is equal to :

(1)
$$2x - \frac{\pi}{3}$$
 (2) $x - \frac{\pi}{6}$

(3)
$$\frac{\pi}{3} - x$$
 (4) $\frac{\pi}{6} - x$

Answer (2)

Sol.
$$2y = \left[\cot^{-1}\left(\frac{\sqrt{3}}{2}\cos x + \frac{1}{2}\sin x}{\frac{1}{2}\cos x - \frac{\sqrt{3}}{2}\sin x}\right)\right]^2$$

$$\Rightarrow 2y = \left[\cot^{-1}\left(\frac{\cos\left(\frac{\pi}{6} - x\right)}{\sin\left(\frac{\pi}{6} - x\right)}\right)\right]^2$$



$$\Rightarrow 2y = \begin{cases} \left(\frac{7\pi}{6} - x\right)^2 & \text{if } \frac{\pi}{6} - x \in \left(-\frac{\pi}{3}, 0\right) \\ \left(\frac{\pi}{6} - x\right)^2 & \text{if } \frac{\pi}{6} - x \in \left(0, \frac{\pi}{6}\right) \end{cases}$$
$$\Rightarrow \frac{dy}{dx} = \begin{cases} x - \frac{7\pi}{6} & \text{if } x \in \left(\frac{\pi}{6}, \frac{\pi}{2}\right) \\ x - \frac{\pi}{6} & \text{if } x \in \left(0, \frac{\pi}{6}\right) \end{cases}$$

Note: Only one given option is correct.

- 5. A point on the straight line, 3x + 5y = 15 which is equidistant from the coordinate axes will lie only in :
 - (1) 4th quadrant
 - (2) 1st quadrant
 - (3) 1st, 2nd and 4th quadrants
 - (4) 1st and 2nd quadrants

Answer (4)

Sol. A point which is equidistant from both the axes lies on either y = x and y = -x.

As it is given that the point lies on the line 3x + 5y = 15

So the required point is :

$$\begin{aligned} &3x+5y=15\\ &\underline{x+y=0}\\ &\overline{x=-\frac{15}{2},} \quad y=\frac{15}{2} \Rightarrow \left(\frac{15}{2},\frac{15}{2}\right) \Big\{2^{nd} \text{ quadrant}\Big\} \end{aligned}$$

$$\frac{\mathbf{x} = \mathbf{y}}{\mathbf{x} = \frac{15}{8}, \quad \mathbf{y} = \frac{15}{8} \Rightarrow \left(\frac{15}{8}, \frac{15}{8}\right) \left\{1^{st} \text{ quadrant}\right\}$$

6.
$$\int \frac{\sin \frac{5x}{2}}{\sin \frac{x}{2}} dx$$
 is equal to :

(where c is a constant of integration)

(1)
$$2x + \sin x + 2 \sin 2x + c$$

(2)
$$x + 2 \sin x + 2 \sin 2x + c$$

(3)
$$x + 2 \sin x + \sin 2x + c$$

(4)
$$2x + \sin x + \sin 2x + c$$

Sol.
$$\int \frac{2}{\sin\left(\frac{x}{2}\right)} dx$$

$$=\int \frac{2\cos\frac{x}{2}.\sin\frac{5x}{2}}{2\cos\frac{x}{2}.\sin\frac{x}{2}}dx$$

$$=\int \frac{\sin 3x + \sin 2x}{\sin x} dx$$

Now use sin2x = 2sinxcosx and $sin3x = 3sinx - 4sin^3x$

 $= \int (1 + 2\cos x + 2\cos 2x) dx$

 $= x + 2\sin x + \sin 2x + c$

7. If the tangents on the ellipse $4x^2 + y^2 = 8$ at the points (1, 2) and (a, b) are perpendicular to each other, then a^2 is equal to

(1)
$$\frac{64}{17}$$
 (2) $\frac{2}{17}$

(3)
$$\frac{4}{17}$$
 (4) $\frac{128}{17}$

Answer (2)

Sol. Equation of tangent at A(1, 2);

 $4x + 2y = 8 \implies 2x + y = 4$

So tangent at B(a, b) can be assumed as

$$x - 2y = c \implies y = \frac{1}{2}x - \frac{c}{2}$$

Condition for tangency;

 \Rightarrow c = $\pm \sqrt{34}$

$$-\frac{\mathbf{c}}{\mathbf{2}} = \pm \sqrt{\mathbf{2}\left(\frac{1}{\mathbf{2}}\right)^2 + \mathbf{8}} = \pm \sqrt{\frac{17}{2}}$$

Equation of tangent; $x - 2y = \pm \sqrt{34}$...(i) Equation of tangent at P(a, b); 4ax + by = 8...(ii)Comparing both the equations ;

$$\frac{4a}{1} = \frac{b}{-2} = \frac{8}{\pm\sqrt{34}}$$
$$\Rightarrow a = \pm \frac{2}{\sqrt{34}} \Rightarrow a^2 = \frac{2}{17}$$



by the lines, x + y = n, $n \in N$, where N is the set of all natural numbers, is

- (1) 105 (2) 160
- (3) 320 (4) 210

Answer (4)



Let the chord x + y = n cuts the circle $x^2 +$ y^2 = 16 at A and B length of perpendicular from

O on $AB = \left| \frac{0 + 0 - n}{\sqrt{1^2 + 1^2}} \right| = \frac{n}{\sqrt{2}}$

Length of chord $AB = 2\sqrt{4^2 - \left(\frac{n}{\sqrt{2}}\right)^2}$

$$=2\sqrt{16-\frac{n^2}{2}}$$

Here possible values of n are 1, 2, 3, 4, 5.

Sum of square of length of chords

$$=\sum_{n=1}^{5}4\left(16-\frac{n^{2}}{2}\right)$$

$$=64 \times 5 - 2.\frac{5 \times 6 \times 11}{6} = 210$$

If $f(x) = \log_{e}\left(\frac{1-x}{1+x}\right), |x| < 1$, then $f\left(\frac{2x}{1+x^{2}}\right)$ is 9. equal to :

(1) 2f(x)(2) $2f(x^2)$ (3) - 2f(x)(4) $(f(x))^2$

Answer (1)

Sol.
$$\therefore$$
 $f(x) = ln\left(\frac{1-x}{1+x}\right)$
 $f\left(\frac{2x}{1+x^2}\right) = ln\left(\frac{1-\frac{2x}{1+x^2}}{1+\frac{2x}{1+2x^2}}\right)$

$$(1+x^{2}+2x)$$
$$= \ln\left(\frac{1-x}{1+x}\right)^{2}$$
$$= 2\ln\left(\frac{1-x}{1+x}\right)$$
$$= 2f(x)$$

10. Let y = y(x) be the solution of the differential equation, $(x^{2} + 1)^{2} \frac{dy}{dx} + 2x (x^{2} + 1)y = 1$ such that y(0) = 0. If $\sqrt{a} y(1) = \frac{\pi}{32}$, then the value of 'a' is (1) $\frac{1}{2}$ (2) $\frac{1}{4}$ (4) $\frac{1}{16}$ (3) 1

Answer (4)

=

Sol.
$$(1+x^2)\frac{2dy}{dx} + 2x(1+x^2)y = 1$$

$$\Rightarrow \frac{dy}{dx} + \left(\frac{2x}{1+x^2}\right)y = \frac{1}{(1+x^2)^2}$$

It is a linear differential equation

$$\mathbf{l} \cdot \mathbf{F} \cdot = \mathbf{e}^{\int \frac{2x}{1+x^2} dx} = \mathbf{e}^{\ln(1+x^2)} = \mathbf{1} + \mathbf{x}^2$$

$$\Rightarrow \quad \mathbf{y} \cdot (\mathbf{1} + \mathbf{x}^2) = \int \frac{d\mathbf{x}}{\mathbf{1} + \mathbf{x}^2} + \mathbf{c}$$

$$\Rightarrow \quad \mathbf{y} \cdot (\mathbf{1} + \mathbf{x}^2) = \tan^{-1} \mathbf{x} + \mathbf{c}$$
If $\mathbf{x} = 0$ then $\mathbf{y} = 0$
So, $0 = 0 + \mathbf{c}$

$$\Rightarrow \quad \mathbf{c} = 0$$

$$\Rightarrow \quad \mathbf{y} \cdot (\mathbf{1} + \mathbf{x}^2) = \tan^{-1} \mathbf{x}$$
put $\mathbf{x} = \mathbf{1}$

$$2\mathbf{y} = \frac{\pi}{4}$$

$$\Rightarrow \quad \mathbf{2} \left(\frac{\pi}{32\sqrt{a}}\right) = \frac{\pi}{4}$$

$$\Rightarrow \quad \sqrt{a} = \frac{1}{4}$$

$$\Rightarrow \quad \mathbf{a} = \frac{1}{16}$$



$$\beta < \frac{\pi}{2}, \text{ then } \alpha - \beta \text{ is equal to}$$
(1) $\tan^{-1}\left(\frac{9}{14}\right)$
(2) $\cos^{-1}\left(\frac{9}{5\sqrt{10}}\right)$
(3) $\sin^{-1}\left(\frac{9}{5\sqrt{10}}\right)$
(4) $\tan^{-1}\left(\frac{9}{5\sqrt{10}}\right)$
Answer (3)
Sol. $\because \cos \alpha = \frac{3}{5}$
 $\Rightarrow \tan \alpha = \frac{4}{3}$

$$\tan (\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \cdot \tan \beta}$$
$$= \frac{\frac{4}{3} - \frac{1}{3}}{1 + \frac{4}{9}} = \frac{1}{\frac{13}{9}}$$
$$= \frac{9}{13}$$
$$\alpha - \beta = \tan^{-1}\left(\frac{9}{13}\right) = \sin^{-1}\left(\frac{9}{5\sqrt{10}}\right) = \cos^{-1}\left(\frac{13}{5\sqrt{10}}\right)$$

0

12. Let $f : [0, 2] \rightarrow R$ be a twice differentiable function such that f''(x) > 0, for all $x \in (0, 2)$. If $\phi(\mathbf{x}) = f(\mathbf{x}) + f(2 - \mathbf{x})$, then ϕ is

(1) Decreasing on (0, 2)

- (2) Increasing on (0, 2)
- (3) Decreasing on (0, 1) and increasing on (1, 2)
- (4) Increasing on (0, 1) and decreasing on (1, 2)

Answer (3)

Sol. $\phi(x) = f(x) + f(2 - x)$

 \Rightarrow and $\tan\beta = \frac{1}{3}$

differentiating w.r.t. x $\phi'(x) = f'(x) - f'(2 - x)$ For $\phi(x)$ to be increasing $\phi'(x) > 0$ \Rightarrow f'(x) > f'(2 - x) (: f''(x) > 0 then f'(x) is an increasing function) > 2 – x

$$\Rightarrow x > 2 -$$

$$\Rightarrow x > 1$$

So $\phi(x)$ is increasing in (1, 2) and decreasing in (0, 1)

(2) 2²⁵ (1) 2²³ (3) 224 (4) 226

Answer (2)

Sol. 2.²⁰C₀ + 5.²⁰C₁ + 8.²⁰C₂ + 62.²⁰C₂₀
=
$$\sum_{r=0}^{20} (3r+2)^{20}C_r = 3\sum_{r=0}^{20} r.^{20}C_r + 2\sum_{r=0}^{20} C_r$$

$$= 60 \sum_{r=1}^{\infty} {}^{19}C_{r-1} + 2 \sum_{r=0}^{\infty} {}^{20}C_r$$
$$= 60 \times 2^{19} + 2 \times 2^{20}$$
$$= 2^{21} [15 + 1] = 2^{25}$$

- 14. All possible numbers are formed using the digits 1, 1, 2, 2, 2, 2, 3, 4, 4 taken all at a time. The number of such numbers in which the odd
 - digits occupy even places is : (1) 180 (2) 175
 - (3) 162 (4) 160

Answer (1)

Sol. There are total 9 digits; out of which only 3 digits are odd.



Number of ways to arrange odd digits first

$$= {}^{4}C_{3} \cdot \frac{|3|}{|2|}$$

Total number of 9 digit numbers = $\begin{pmatrix} {}^{4}C_{3} \cdot \frac{|3|}{|2|} \end{pmatrix} \cdot \frac{|6|}{|2|4|}$ = 180

(2) 0

(4) $\frac{1}{2}$

15. The greatest value of $c \in R$ for which the system of linear equations

$$x - cy - cz = 0$$

 $cx - y + cz = 0$
 $cx + cy - z = 0$
has a non-trivial solution, is :
(1) -1 (2) 0
(3) 2 (4) $\frac{1}{2}$



$$\begin{vmatrix} 1 & -\mathbf{c} & -\mathbf{c} \\ \mathbf{c} & -\mathbf{1} & \mathbf{c} \\ \mathbf{c} & \mathbf{c} & -\mathbf{1} \end{vmatrix} = \mathbf{0}$$

$$\Rightarrow (\mathbf{1} - \mathbf{c}^2) + \mathbf{c} (-\mathbf{c} - \mathbf{c}^2) - \mathbf{c} (\mathbf{c}^2 + \mathbf{c}) = \mathbf{0}$$

$$\Rightarrow (\mathbf{1} + \mathbf{c}) (\mathbf{1} - \mathbf{c}) - 2\mathbf{c}^2 (\mathbf{1} + \mathbf{c}) = \mathbf{0}$$

$$\Rightarrow (\mathbf{1} + \mathbf{c}) (\mathbf{1} - \mathbf{c} - 2\mathbf{c}^2) = \mathbf{0}$$

$$\Rightarrow (\mathbf{1} + \mathbf{c})^2 (\mathbf{1} - 2\mathbf{c}) = \mathbf{0}$$

$$\Rightarrow \mathbf{c} = -\mathbf{1} \text{ or } \frac{1}{2}$$

- 16. Let O(0, 0) and A(0, 1) be two fixed points. Then the locus of a point P such that the perimeter of \triangle AOP is 4, is :
 - (1) $8x^2 9y^2 + 9y = 18$ (2) $9x^2 + 8y^2 8y = 16$ (3) $9x^2 - 8y^2 + 8y = 16$ (4) $8x^2 + 9y^2 - 9y = 18$

Answer (2)

Sol. Let point P(h, k)

..
$$OA = 1$$

So, $OP + AP = 3$
 $\sqrt{h^2 + k^2} + \sqrt{h^2 + (k - 1)^2} = 3$
Y
A(0, 1)
A(0, 1)
P(h, k)
X'
 $O(0, 0)$
Y'
 $\Rightarrow h^2 + (k - 1)^2 = 9 + h^2 + k^2 - 6\sqrt{h^2 + k^2}$
 $\Rightarrow 6\sqrt{h^2 + k^2} = 2k + 8$
 $\Rightarrow 9(h^2 + k^2) = k^2 + 16 + 8k$
 $\Rightarrow 9h^2 + 8k^2 - 8k - 16 = 0$
Locus of point P will be,
 $9x^2 + 8y^2 - 8y - 16 = 0$

is always correct?

(1) P(A | B) = P(B) - P(A)(2) $P(A | B) \le P(A)$ (3) $P(A | B) \ge P(A)$ (4) P(A | B) = 1Answer (3) Sol. $\cdots A \subset B$; so $A \cap B = A$ Now, $P\left(\frac{A}{B}\right) = \frac{P(A \cap B)}{P(B)}$ (A) P(A)

$$\Rightarrow P\left(\frac{A}{B}\right) = \frac{P(A)}{P(B)}$$

$$\therefore P(B) \le 1$$

So, $P\left(\frac{A}{B}\right) \ge P(A)$

- 18. If α and β be the roots of the equation $x^2 2x + 2 = 0$, then the least value of n for which
 - $\left(\frac{\alpha}{\beta}\right)^{n} = 1$ is : (1) 4 (2) 5 (3) 3 (4) 2

Answer (1)

Sol. $x^2 - 2x + 2 = 0$

roots of this equation are
$$\frac{2\pm\sqrt{-4}}{2} = 1\pm i$$

Then $\frac{\alpha}{\beta} = \frac{1+i}{1-i} = \frac{(1+i)^2}{1-i^2} = i$
or $\frac{\alpha}{\beta} = \frac{1-i}{1+i} = \frac{(1-i)^2}{1-i^2} = -i$
So, $\frac{\alpha}{\beta} = \pm i$
Now, $\left(\frac{\alpha}{\beta}\right)^n = 1 \Rightarrow (\pm i)^n = 1$

 \Rightarrow n must be a multiple of 4. minimum value of n = 4



(1)
$$\frac{26}{3}$$
 (2) $\frac{59}{6}$
(3) 8 (4) $\frac{53}{6}$

Answer (2)

Sol. $y \le x^2 + 3x$ represents region below the parabola.



Area of the required region

$$= \int_{0}^{1} (x^{2} + 3x) dx + \int_{1}^{3} 4 \cdot dx$$

= $\frac{1}{3} + \frac{3}{2} + 8$
= $\frac{59}{6}$
If $f(x) = \frac{2 - x \cos x}{2 + x \cos x}$ and $g(x) = \log_{e} x$, (x > 0) then

the value of the integral $\int_{1}^{\frac{\pi}{4}} g(f(x))dx$ is : (1) log_1 $(2) \log_{2} 3$ (3) log_2 (4) log_e

Answer (1)

20.

Sol.
$$g(f(x)) = ln\left(\frac{2 - x\cos x}{2 + x\cos x}\right)$$

Let $I = \int_{-\pi/4}^{\pi/4} ln\left(\frac{2 - x\cos x}{2 + x\cos x}\right) dx$...(i)
 $\left(Using property \int_{a}^{b} f(x) dx = \int_{a}^{b} f(a + b - x) dx\right)$
 $I = \int_{-\pi/4}^{\pi/4} ln\left(\frac{2 + x\cos x}{2 - x\cos x}\right) dx$...(ii)
Adding (i) and (ii)
 $2I = \int_{-\pi/2}^{\pi/2} ln(1) dx = 0$
 $\Rightarrow I = 0 = ln 1$

y + 2z - 4 = 0 and passing through the point

Answer (3)

(1, 1, 0) is : (1) 2x - z = 2(2) x - 3y - 2z = -2(3) x - y - z = 0(4) x + 3y + z = 4Sol. Let the equation of required plane be; $(2x - y - 4) + \lambda(y + 2z - 4) = 0$ \therefore This plane passes through (1, 1, 0) then $(2 - 1 - 4) + \lambda(1 + 0 - 4) = 0 \Rightarrow \lambda = -1$ Equation of required plane will be (2x - y - 4) - (y + 2z - 4) = 0 \Rightarrow 2x - 2y - 2z = 0 \Rightarrow x - y - z = 0 $\cos(\alpha+\beta)=\frac{3}{5}$, $\sin(\alpha-\beta)=\frac{5}{13}$ 22. If and $\mathbf{0} < \alpha, \beta < \frac{\pi}{4}$, then tan($\mathbf{2}\alpha$) is equal to : (1) $\frac{21}{16}$ (2) $\frac{63}{52}$ (3) $\frac{33}{52}$ (4) $\frac{63}{16}$ Answer (4) **Sol.** $\therefore \alpha + \beta$ and $\alpha - \beta$ both are acute angles. $\cos(\alpha+\beta)=\frac{3}{5}$ \Rightarrow tan $(\alpha + \beta) = \frac{4}{3}$ And $\sin(\alpha-\beta)=\frac{5}{12}$ \Rightarrow tan $(\alpha - \beta) = \frac{5}{12}$ Now, $\tan 2\alpha = \tan ((\alpha + \beta) + (\alpha - \beta))$ $=\frac{\tan(\alpha+\beta)+\tan(\alpha-\beta)}{1-\tan(\alpha+\beta)\cdot\tan(\alpha-\beta)}$ $=\frac{\frac{4}{3}+\frac{5}{12}}{1-\frac{4}{3}\cdot\frac{5}{12}}=\frac{63}{16}$ 23. Let $A = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}$, $(\alpha \in \mathbf{R})$ such that $A^{32} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Then a value of α is : (2) $\frac{\pi}{64}$ (1) $\frac{\pi}{32}$ (4) $\frac{\pi}{16}$ (3) 0 Answer (2)



$$\begin{bmatrix} \sin \alpha & \cos \alpha \end{bmatrix}$$

$$A^{2} = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

$$= \begin{bmatrix} \cos 2\alpha & -\sin 2\alpha \\ \sin 2\alpha & \cos 2\alpha \end{bmatrix}$$
Then $A^{4} = A^{2}.A^{2} = \begin{bmatrix} \cos 4\alpha & -\sin 4\alpha \\ \sin 4\alpha & \cos 4\alpha \end{bmatrix}$
similarly $A^{8} = A^{4}.A^{4} \begin{bmatrix} \cos 8\alpha & -\sin 8\alpha \\ \sin 8\alpha & \cos 8\alpha \end{bmatrix}$
and so on $A^{32} = \begin{bmatrix} \cos 32\alpha & -\sin 32\alpha \\ \sin 32\alpha & \cos 32\alpha \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$
So $\sin 32\alpha = 1$ and $\cos 32\alpha = 0$

$$\Rightarrow 32\alpha = 2n\pi + \frac{\pi}{2} \Rightarrow \alpha = \frac{n\pi}{16} + \frac{\pi}{64} \text{ where } n \in \mathbb{Z}$$
put $n = 0, \ \alpha = \frac{\pi}{64}$

- 24. The contrapositive of the statement "If you are born in India, then you are a citizen of India", is:
 - (1) If you are born in India, then you are not a citizen of India.
 - (2) If you are not born in India, then you are not a citizen of India.
 - (3) If you are a citizen of India, then you are born in India.
 - (4) If you are not a citizen of India, then you are not born in India.

Answer (4)

Sol. S: "If you are born in India, then you are a citizen of India."

Contrapositive of $\textbf{p} \rightarrow \textbf{q}$ is $\textbf{\neg q} \rightarrow \textbf{\neg p}$

So contrapositive of statement S will be :

"If you are not a citizen of India, then you are not born in India."

25. The sum of the solutions of the equation $|\sqrt{x}-2|+\sqrt{x}(\sqrt{x}-4)+2=0, (x>0)$ is equal to:

(1) 4	(2)	10
(·/ ·	(=)	

(3) 9 (4) 12

Answer (2)

 $\Rightarrow |t-2|+t^2-4t+4-2=0$ $\Rightarrow |t-2|+(t-2)^2-2=0$ Let |t-2| = z (Clearly $z \ge 0$) $\Rightarrow z+z^2-2=0$ $\Rightarrow z = 1 \text{ or } -2 \text{ (rejected)}$ $\Rightarrow |t-2| = 1 \Rightarrow t = 1, 3$ If $\sqrt{x} = 1 \Rightarrow x = 1$ If $\sqrt{x} = 3 \Rightarrow x = 9$

Sum of solutions = 10

26. The magnitude of the projection of the vector $2\hat{i}+3\hat{j}+\hat{k}$ on the vector perpendicular to the plane containing the vectors $\hat{i}+\hat{j}+\hat{k}$ and $\hat{i}+2\hat{j}+3\hat{k}$, is:

(1)
$$\sqrt{\frac{3}{2}}$$
 (2) $3\sqrt{6}$
(3) $\frac{\sqrt{3}}{2}$ (4) $\sqrt{6}$

Answer (1)

Sol. Let $\overline{a} = \hat{i} + \hat{j} + \hat{k}$ and $\overline{b} = \hat{i} + 2\hat{j} + 3\hat{k}$ vector

perpendicular to \overline{a} and \overline{b} is $\overline{a} \times \overline{b}$

$$\overline{\mathbf{a}} \times \overline{\mathbf{b}} = \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ 1 & 1 & 1 \\ 1 & 2 & 3 \end{vmatrix} = \hat{\mathbf{i}} - 2\hat{\mathbf{j}} + \hat{\mathbf{k}}$$

Projection of vector $\overline{c} = 2\hat{i} + 3\hat{j} + \hat{k}$ on $\overline{a} \times \overline{b}$ is

$$= \left| \frac{\overline{\mathbf{c}} \cdot \left(\overline{\mathbf{a}} \times \overline{\mathbf{b}} \right)}{\left| \overline{\mathbf{a}} \times \overline{\mathbf{b}} \right|} \right| = \left| \frac{2 - 6 + 1}{\sqrt{6}} \right|$$
$$= \frac{3}{\sqrt{6}} = \sqrt{\frac{3}{2}}$$

- 27. The mean and variance of seven observations are 8 and 16, respectively. If 5 of the observations are 2, 4, 10, 12, 14, then the product of the remaining two observations is:
 - (1) 45 (2) 40

(3) 48 (4) 49

Answer (3)



Mean
$$(\bar{\mathbf{x}}) = \frac{\sum x_i}{N} = \frac{2+4+10+12+14+x+y}{7} = 8$$

 $\Rightarrow x + y = 14$ (i)
Variance $(\sigma^2) = \frac{\sum x_i^2}{N} - (\bar{\mathbf{x}})^2 = 16$
 $\Rightarrow \frac{2^2 + 4^2 + 10^2 + 12^2 + 14^2 + x^2 + y^2}{7} - (8)^2 = 16$
 $\Rightarrow x^2 + y^2 = 100$ (ii)
From (i) and (ii) (x, y) = (6, 8) or (8, 6)
xy = 48
28. $\lim_{x \to 0} \frac{\sin^2 x}{\sqrt{2} - \sqrt{1 + \cos x}}$ equals:
(1) $\sqrt{2}$ (2) $2\sqrt{2}$

(3) 4 (4)
$$4\sqrt{2}$$

Answer (4)

Sol.
$$\lim_{x \to 0} \frac{\sin^2 x}{\sqrt{2} - \sqrt{1 + \cos x}} = \lim_{x \to 0} \frac{\sin^2 x}{\sqrt{2} - \sqrt{2\cos^2 \frac{x}{2}}} \quad \begin{bmatrix} 0\\0 \end{bmatrix}$$

$$= \lim_{x \to 0} \frac{\sin^2 x}{\sqrt{2} \left[1 - \cos \frac{x}{2} \right]}$$
$$= \lim_{x \to 0} \frac{\sin^2 x}{2\sqrt{2} \sin^2 \frac{x}{4}}$$
$$= \lim_{x \to 0} \frac{\left(\frac{\sin x}{x}\right)^2 .16}{2\sqrt{2} \left(\frac{\sin \frac{x}{4}}{\frac{x}{4}}\right)^2}$$

$$=\frac{16}{2\sqrt{2}}=4\sqrt{2}$$

29. The sum of all natural numbers 'n' such that 100 < n < 200 and H.C.F. (91, n) > 1 is :

(1) 3303	(2) 3121
(3) 3203	(4) 3221

Answer (2)

7 or 13

Sum of such numbers = Sum of no. divisible by 7 + sum of the no. divisible by 13 – Sum of the numbers divisible by 91

= (105 + 112 + + 196) + (104 + 117 + + 195) – 182

= 2107 + 1196 – 182

= 3121

30. The length of the perpendicular from the point (2, -1, 4) on the straight line,

$$\frac{x+3}{10} = \frac{y-2}{-7} = \frac{z}{1}$$
 is:

- (1) Greater than 3 but less than 4
- (2) Greater than 2 but less than 3
- (3) Greater than 4
- (4) Less than 2

Answer (1)



Let P be the foot of perpendicular from point A(2, -1, 4) on the given line. So P can be assumed as P($10\lambda - 3$, $-7\lambda + 2$, λ)

DR's of AP \propto to 10 λ – 5, –7 λ + 3, λ – 4

· AP and given line are perpendicular, so

$$10(10\lambda - 5) - 7(-7\lambda + 3) + 1(\lambda - 4) = 0$$

$$\Rightarrow \lambda = \frac{1}{2}$$

$$AP = \sqrt{(10\lambda - 5)^2 + (-7\lambda + 3)^2 + (\lambda - 4)^2}$$

$$= \sqrt{0 + \frac{1}{4} + \frac{49}{4}}$$

$$=\sqrt{12.5}$$
 ; $\sqrt{12.5}\in$ (3, 4)



Answers & Solutions

for

JEE (MAIN)-2019 (Online) Phase-2

(Physics, Chemistry and Mathematics)

Time : 3 hrs.

M.M.: 360

Important Instructions :

- 1. The test is of **3 hours** duration.
- 2. The Test Booklet consists of **90** questions. The maximum marks are **360**.
- 3. There are *three* parts in the question paper A, B, C consisting of **Physics**, **Chemistry** and **Mathematics** having 30 questions in each part of equal weightage.
- 4. Each question is allotted 4 (four) marks for each correct response. ¼ (one-fourth) marks will be deducted for indicating incorrect response of each question. No deduction from the total score will be made if no response is indicated for an item in the answer sheet.
- 5. There is only one correct response for each question.



1. A cell of internal resistance r drives current through an external resistance R. The power delivered by the cell to the external resistance will be maximum when :

(1) R = 1000 r	(2) R = r
(3) R = 2r	(4) R = 0.001

Answer (2)

Sol. For maximum power in external resistance, Internal resistance = External resistance ⇒ R = r

r

- 2. A body of mass m_1 moving with an unknown velocity of $v_1\hat{i}$, undergoes a collinear collision with a body of mass m_2 moving with a velocity $v_2\hat{i}$. After collision, m_1 and m_2 move with
 - velocities of $v_3 \hat{i}$ and $v_4 \hat{i}$, respectively.
 - If $m_2 = 0.5 m_1$ and $v_3 = 0.5 v_1$, then v_1 is :

(1)
$$v_4 - \frac{v_2}{4}$$
 (2) $v_4 - v_2$
(3) $v_4 + v_2$ (4) $v_4 - \frac{v_2}{2}$

Answer (2)

Sol.
$$m_1v_1 + m_2v_2 = m_1v_3 + m_2v_4$$

 $m_1v_1 + 0.5m_1v_2 = 0.5m_1v_1 + 0.5m_1v_4$
 $v_1 = v_4 - v_2$

3. In a line of sight radio communication, a distance of about 50 km is kept between the transmitting and receiving antennas. If the height of the receiving antenna is 70 m, then the minimum height of the transmitting antenna should be :

(Radius of the Earth = 6.4×10^6 m).

- (1) 20 m (2) 51 m
- (3) 32 m (4) 40 m

Answer (3)

Sol.
$$\sqrt{2 \times 70 \times R_E} + \sqrt{2 \times h_R \times R_E} = 50 \times 10^3$$

Putting R_E = 6.4 × 10⁶ m and solving we get

h_R = 32 m

4. The electric field in a region is given by $\vec{E} = (Ax+B)\hat{i}$, where E is in NC⁻¹ and x is in metres. The values of constants are A = 20 SI unit and B = 10 SI unit. If the potential at x = 1 is V₁ and that at x = -5 is V₂, then V₁ - V₂ is :

(1) 180 V (2) -520 V

(3) 320 V (4) -48 V

Answer (1)

Sol.
$$dV = -\vec{E} \cdot \vec{dr} = -(Ax + B)dx$$

$$\int_{V_{2}}^{V_{1}} dV = \int_{-5}^{1} -(Ax+B) dx$$
$$V_{1} - V_{2} = \left(-A\frac{x^{2}}{2} - Bx\right)_{-5}^{1}$$
$$= \left(-\frac{A}{2} - B\right) + \left(\frac{A}{2}25 + B(-5)\right)$$
$$= 12A - 6B = 240 - 60 = 180 \text{ V}$$

5. A parallel plate capacitor has 1 μ F capacitance. One of its two plates is given +2 μ C charge and the other plate, +4 μ C charge. The potential difference developed across the capacitor is :

Answer (2)

6. The given diagram shows four processes i.e., isochoric, isobaric, isothermal and adiabatic. The correct assignment of the processes, in the same order is given by :





7. Two magnetic dipoles X and Y are placed at a separation d, with their axes perpendicular to each other. The dipole moment of Y is twice that of X. A particle of charge q is passing through their mid-point P, at angle θ = 45° with the horizontal line, as shown in figure. What would be the magnitude of force on the particle at that instant? (d is much larger than the dimensions of the dipole)



(1) 0

(2)
$$\sqrt{2} \left(\frac{\mu_0}{4\pi}\right) \frac{M}{\left(\frac{d}{2}\right)^3} \times qv$$

$$(3) \ \left(\frac{\mu_0}{4\pi}\right) \frac{2M}{\left(\frac{d}{2}\right)^3} \times qv$$

$$\textbf{(4)} \quad \left(\frac{\mu_0}{4\pi}\right) \frac{\textbf{M}}{\left(\frac{\textbf{d}}{\textbf{2}}\right)^3} \times \textbf{qv}$$

Answer (1)

Sol. $\vec{F}_m = q(\vec{V} \times \vec{B})$

 $\vec{\mathbf{B}} = \vec{\mathbf{B}}_{x} + \vec{\mathbf{B}}_{y}$

$$\begin{array}{c} X \\ \hline S \\ \hline M \\ \hline d \\ \hline 2 \\ \hline d \\ \hline 2 \\ \hline S \\ \hline N \\ \hline B_x \\ \hline B_x \\ \hline B_y \\ \hline B_{net} \\ is parallel to \ \vec{V} \end{array}$$

 $\implies ~\vec{F}=0$

drops to half its value for every 10 oscillations.

The time it will take to drop to $\frac{1}{1000}$ of the original amplitude is close to:

- (1) 100 s
- (2) 10 s
- (3) 50 s
- (4) 20 s

Answer (4)

Sol. Time for 10 oscillations =
$$\frac{10}{5} = 2$$
 s

$$A = A_0 e^{-kt}$$

$$\frac{1}{2} = e^{-2k} \implies \ln 2 = 2k$$

$$10^{-3} = e^{-kt} \implies 3\ln 10 = kt$$

$$t = \frac{3\ln 10}{k} = \frac{3\ln 10}{\ln 2} \times 2$$

$$= 6 \times \frac{2.3}{0.69} \approx 20 s$$

- 9. If Surface tension (S), Moment of Inertia (I) and Planck's constant (h), were to be taken as the fundamental units, the dimensional formula for linear momentum would be:
 - (1) S^{1/2}I^{3/2}h⁻¹
 - (2) S^{3/2}l^{1/2}h⁰
 - (3) S^{1/2}I^{1/2}h⁻¹
 - (4) S^{1/2}I^{1/2}h⁰

Sol. [p] = MLT⁻¹ = [I^x h^y S^z]
= M^xL^{2x} (ML²T⁻¹)^y (MT⁻²)^z
= M^{x+y+z} L^{2x+2y} T^{-y-2z}
x + y + z = 1
2(x + y) = 1
$$\Rightarrow$$
 x + y = $\frac{1}{2}$ \Rightarrow z = $\frac{1}{2}$
y + 2z = 1 \Rightarrow y = 0 \Rightarrow x = $\frac{1}{2}$
[p] = \sqrt{IS}



difference of $\frac{\pi}{4}$ between the emf e and current i. Which of the following circuits will exhibit this?

- (1) RL circuit with R = 1 k Ω and L = 10 mH
- (2) RL circuit with R = 1 k Ω and L = 1 mH
- (3) RC circuit with R = 1 k Ω and C = 10 μF
- (4) RC circuit with R = 1 k Ω and C = 1 μF

Answer (3)

Sol. As $\phi = \frac{\pi}{4}$, $x_c = R$

11. The magnetic field of an electromagnetic wave is given by:

$$\vec{B} = 1.6 \times 10^{-6} \cos(2 \times 10^7 z + 6 \times 10^{15} t) (2\hat{i} + \hat{j}) \frac{Wb}{m^2}$$

The associated electric field will be:

(1)
$$\vec{E} = 4.8 \times 10^2 \cos(2 \times 10^7 z + 6 \times 10^{15} t) (-\hat{i} + 2\hat{j}) \frac{V}{m}$$

(2)
$$\vec{E} = 4.8 \times 10^2 \cos(2 \times 10^7 z - 6 \times 10^{15} t) (2\hat{i} + \hat{j}) \frac{V}{m}$$

(3)
$$\vec{E} = 4.8 \times 10^2 \cos(2 \times 10^7 z - 6 \times 10^{15} t) (-2\hat{j} + \hat{i}) \frac{V}{m}$$

(4) $\vec{E} = 4.8 \times 10^2 \cos(2 \times 10^7 z + 6 \times 10^{15} t) (\hat{i} - 2\hat{j}) \frac{V}{m}$

Answer (1)

Sol. Amplitude of electric field, $E = B_0C$

- $= 1.6 \times 10^{-6} \times \sqrt{5} \times 3 \times 10^{8}$
- $= 4.8 \times 10^2 \sqrt{5} \text{ V/m}$

Also $\vec{E} \times \vec{B}$ is along $-\hat{k}$ (the direction of propagation)

$$\Rightarrow \vec{\mathsf{E}} = 4.8 \times 10^2 \cos(2 \times 10^7 z + 6 \times 10^{15} t) (-\hat{i} + 2\hat{j}) \frac{V}{m}$$

12. In the circuit shown, a four-wire potentiometer is made of a 400 cm long wire, which extends between A and B. The resistance per unit length of the potentiometer wire is $r = 0.01 \Omega/$ cm. If an ideal voltmeter is connected as shown with jockey J at 50 cm from end A, the expected reading of the voltmeter will be:



Answer (4)

Sol. Resistance of potentiometer wire, $R_p = 400 \times 0.01$ = 4 Ω

$$\Rightarrow I = \frac{3}{6} = 0.5 \text{ A}$$

$$\Rightarrow \text{ Reading of voltmeter} = I R_{AJ}$$

$$= 0.5 \times 50 \times 0.01$$

$$= 0.25 \text{ V}$$

13. A nucleus A, with a finite de-Broglie wavelength $\lambda_{\rm A}$, undergoes spontaneous fission into two nuclei B and C of equal mass. B flies in the same direction as that of A, while C flies in the opposite direction with a velocity equal to half of that of B. The de-Broglie wavelength $\lambda_{\rm B}$ and $\lambda_{\rm C}$ of B and C are respectively:

(1)
$$2 \lambda_A, \lambda_A$$
 (2) $\lambda_A, \frac{\lambda_A}{2}$
(3) $\frac{\lambda_A}{2}, \lambda_A$ (4) $\lambda_A, 2 \lambda_A$

Answer (3)

Sol.
$$\lambda_A = \frac{h}{mV_A}$$

Conservation of linear momentum

. .

. .

$$\Rightarrow mV_{A} = \frac{m}{2}V - \frac{m}{2} \cdot \frac{V}{2} = \frac{mV}{4}$$
$$\Rightarrow \lambda_{A} = \frac{4h}{mV}$$
$$\therefore V_{A} = \frac{V}{4}$$
$$\lambda_{B} = \frac{h}{\frac{m}{2}V} = \frac{2h}{mV} = \frac{\lambda_{A}}{2}$$
$$\lambda_{c} = \frac{h}{\frac{m}{2} \cdot \frac{V}{2}} = \frac{4h}{mV} = \lambda_{A}$$



energy delivered by the rocket launcher, what should be the minimum energy that the launcher should have if the same rocket is to be launched from the surface of the moon? Assume that the density of the earth and the moon are equal and that the earth's volume is 64 times the volume of the moon.

(1)
$$\frac{E}{64}$$
 (2) $\frac{E}{4}$
(3) $\frac{E}{16}$ (4) $\frac{E}{32}$

Answer (3)

Sol.
$$E = \frac{GM_Em}{R_E}$$

$$E' = \frac{GM_mm}{R_M}$$

$$\rho R_{E}^{3} = 64 \rho R_{M}^{3}$$

$$\Rightarrow R_{E} = 4 R_{M}$$

$$\frac{E'}{E} = \frac{M_{M}}{M_{E}} \cdot \frac{R_{E}}{R_{M}} = \frac{1}{64} \cdot 4 =$$

$$\Rightarrow E' = \frac{E}{16}$$

15. In a simple pendulum experiment for determination of acceleration due to gravity (g), time taken for 20 oscillations is measured by using a watch of 1 second least count. The mean value of time taken comes out to be 30 s. The length of pendulum is measured by using a meter scale of least count 1 mm and the value obtained is 55.0 cm. The percentage error in the determination of g is close to:

1 16

- (1) 6.8%
- (2) 0.2%
- (3) 3.5%
- (4) 0.7%

Sol.
$$T = 2\pi \sqrt{\frac{I}{g}} \Rightarrow g = 4\pi^2 \frac{I}{T^2}$$

$$\frac{\Delta g}{g} = \frac{\Delta I}{I} + \frac{2\Delta T}{T} = \left(\frac{0.1}{55} + \frac{2\times 1}{30}\right) \times 100$$
$$\approx 6.8\%$$

16. An electric dipole is formed by two equal and opposite charges q with separation d. The charges have same mass m. It is kept in a uniform electric field E. If it is slightly rotated from its equilibrium orientation, then its angular frequency ω is:

(1)
$$\sqrt{\frac{2 qE}{md}}$$

(2) $2\sqrt{\frac{qE}{md}}$
(3) $\sqrt{\frac{qE}{2md}}$

(4)
$$\sqrt{\frac{qE}{md}}$$







of a platform of height 5 m. When released, it slips off the table in a very short time τ = 0.01 s, remaining essentially horizontal. The angle by which it would rotate when it hits the ground will be (in radians) close to:



- (1) 0.3
- (2) 0.02
- (3) 0.28
- (4) 0.5

Answer (4)

Sol. Initial angular acceleration before it slips off

 $mg\frac{l}{2} = l\alpha$

- $\alpha = \frac{3g}{2I}$
- :. Angular speed acquire by the box in time τ = 0.01 s

 $\omega = \alpha t = \frac{3g}{2l} \times 0.01 = \frac{3 \times 10 \times 0.01}{2 \times 0.3} = \frac{1}{2} \text{ rad/sec}$

... The angle by which it would rotate when hits the ground

 $\theta = \omega t'$

[Assuming ω = constant and t' = time of fall =

$$\sqrt{\frac{2H}{g}} = 1 \sec \theta$$

 $\therefore \quad \theta = \frac{1}{2} \text{ radians}$

in the figure. If the shaded portion HBGO is cut-off, the coordinates of the centre of mass of the remaining portion will be:



Answer (2)

Sol. X- coordinate of CM of remaining sheet

$$X_{cm} = \frac{MX - mx}{M - m}$$
$$= \frac{(4m) \times \left(\frac{a}{2}\right) - M\left(\frac{3a}{4}\right)}{4m - m} = \frac{5a}{12}$$
Similarly $y_{cm} = \frac{5b}{12}$

$$\therefore \quad \mathsf{CM}\left(\frac{5a}{12},\frac{5b}{12}\right)$$

19. In the figure shown, what is the current (in Ampere) drawn from the battery? You are given :





COI. Equivalent resistance of the given chedit

$$R_{eq} = 45 + \frac{10 \times 50}{10 + 50} = \left(45 + \frac{50}{6}\right)\Omega = \frac{160}{3}\Omega$$

$$\therefore \quad I = \frac{15}{\frac{160}{3}} = \frac{9}{32}A$$

20. Let $|\overline{A_1}| = 3$, $|\overline{A_2}| = 5$ and $|\overline{A_1} + \overline{A_2}| = 5$. The value of $(2\overline{A_1} + 3\overline{A_2}) \cdot (3\overline{A_1} - 2\overline{A_2})$ is : (1) - 106.5 (2) - 118.5 (3) - 99.5 (4) - 112.5

Answer (2)

Sol.
$$(2\overrightarrow{A_1} + 3\overrightarrow{A_2}) \cdot (3\overrightarrow{A_1} - 2\overrightarrow{A_2}) = 6 |\overrightarrow{A_1}|^2 + 5\overrightarrow{A_1} \cdot \overrightarrow{A_2} - 6 |\overrightarrow{A_2}|^2$$

= $(6 \times 9) + 5\overrightarrow{A_1} \cdot \overrightarrow{A_2} - (6 \times 25) \dots (i)$

As
$$25 = 9 + 25 + 2A_1 \cdot A_2$$

 $\Rightarrow 2\overrightarrow{A_1} \cdot \overrightarrow{A_2} = -9$...(ii)

From (i) and (ii),

$$\Rightarrow (2\overline{A_1} + 3\overline{A_2}) \cdot (3\overline{A_1} - 2\overline{A_2}) = 54 - 22.5 - 150$$
$$= -118.5$$

21. Calculate the limit of resolution of a telescope objective having a diameter of 200 cm, if it has to detect light of wavelength 500 nm coming from a star.

(1) 457.5 × 10⁻⁹ radian

- (2) 305 × 10⁻⁹ radian
- (3) 152.5 × 10⁻⁹ radian
- (4) 610 × 10⁻⁹ radian

Answer (2)

Sol.
$$\theta = \frac{1.22\lambda}{D}$$

 $\theta = \frac{1.22 \times 500 \times 10^{-9}}{200 \times 10^{-2}} = \frac{1.22 \times 500 \times 10^{-9}}{2}$
 $\theta = 305 \times 10^{-9}$ radian

xy-plane as shown in the figure.



These wires carry currents of equal magnitude I, whose directions are shown in the figure. The net magnetic field at point P will be

(1) Zero (2)
$$\frac{+\mu_0 I}{\pi d}(\hat{z})$$

(3)
$$-\frac{\mu_0 I}{2\pi d}(\hat{x} + \hat{y})$$
 (4) $\frac{\mu_0 I}{2\pi d}(\hat{x} + \hat{y})$

Answer (1)

Sol.
$$\vec{\mathbf{B}} = \vec{\mathbf{B}_1} + \vec{\mathbf{B}_2} = \frac{\mu_0 \mathbf{1}}{2\pi d} \left[\hat{\mathbf{k}} - \hat{\mathbf{k}} \right] = \mathbf{0}$$

23. A positive point charge is released from rest at a distance r_0 from a positive line charge with uniform density. The speed (v) of the point charge, as a function of instantaneous distance r from line charge, is proportional to :





Zne₀i

$$\begin{split} & \bigvee_{V_{G}}^{V_{P}} dV = \int -\frac{\lambda}{2\pi\epsilon_{0}r} dr \\ & \Rightarrow \quad V_{P} - V_{G} = \frac{\lambda}{2\pi\epsilon_{0}} ln \frac{r}{r_{0}} \\ & \frac{1}{2}mv^{2} = q(V_{P} - V_{G}) \\ & \Rightarrow \quad v \propto \left[ln \left(\frac{r}{r_{0}}\right) \right]^{\frac{1}{2}} \end{split}$$

24. A convex lens (of focal length 20 cm) and a concave mirror, having their principal axes along the same lines, are kept 80 cm apart from each other. The concave mirror is to the right of the convex lens. When an object is kept at a distance of 30 cm to the left of the convex lens, its image remains at the same position even if the concave mirror is removed. The maximum distance of the object for which this concave mirror, by itself would produce a virtual image would be :

(1) 30 cm (2)	25 c	m
---------------	------	---

(3) 20 cm (4) 10 cm

Answer (4)



So clearly radius of curvature of mirror is 20 cm. Now if the object is placed within focal plane i.e. 10 cm then image formed by mirror is virtual.

(1) 0.1	(2) 1
(3) 2	(4) 5

Answer (2)

- Sol. Densities of nucleus happens to be constant, irrespective of mass number.
- 26. A common emitter amplifier circuit, built using an npn transistor, is shown in the figure. Its dc current gain is 250, $R_c = 1 \text{ k}\Omega$ and $V_{cc} = 10 \text{ V}$. What is the minimum base current for V_{CE} to reach saturation?



Answer (4)

Sol. For saturation, $V_{CC} - i_c \times R_c = 0$

$$\Rightarrow i_{c} = \frac{V_{CC}}{R_{c}} = \frac{10}{10^{3}} = 10^{-2} \text{ A}$$
$$\therefore \quad \beta = \frac{I_{C}}{I_{B}} = 250$$

:.
$$I_{\rm B} = \frac{I_{\rm C}}{250} = \frac{10^{-2}}{250} = 40 \,\mu {\rm A}$$

27. A solid sphere and solid cylinder of identical radii approach an incline with the same linear velocity (see figure). Both roll without slipping all throughout. The two climb maximum heights

 h_{sph} and h_{cyl} on the incline. The ratio $\frac{h_{sph}}{h_{cyl}}$ is given by





Sol.
$$mgh_{sph} = \frac{1}{2}mv^2 + \frac{1}{2} \cdot \frac{2}{5}mR^2 \cdot \left(\frac{v}{R}\right)^2$$

 $= \frac{7}{10}mv^2$...(i)
 $mgh_{cylinder} = \frac{1}{2}mv^2 + \frac{1}{2}\frac{mR^2}{2}\left(\frac{v}{R}\right)^2 = \frac{3}{4}mv^2$
 $\Rightarrow \frac{h_{sph}}{h_{cylinder}} = \frac{7 \times 4}{10 \times 3} = \frac{14}{15}$

28. A particle starts from origin O from rest and moves with a uniform acceleration along the positive x-axis. Identify all figures that correctly represent the motion qualitatively. (a = acceleration, v = velocity,

x = displacement, t = time)



Sol. a = Constant

$$x = \frac{1}{2}at^2$$
 [Particle starts from the origin]

A, B and D are correct graphs.

Wire B is 1.5 m long and has radius 2 mm. If the two wires stretch by the same length for a given load, then the value of R is close to

(1)	1.3 mm	(2)	1.9 mm
(3)	1.5 mm	(4)	1.7 mm

Answer (4)

Sol.
$$\Delta L = \frac{FL}{YA}$$

 $\Rightarrow \frac{L_A}{Y_A r_A^2} = \frac{L_B}{Y_B r_B^2}$
 $\Rightarrow r_A^2 = \sqrt{\frac{L_A}{L_B} \cdot \frac{Y_B}{Y_A}} \cdot r_B$
 $= \sqrt{\frac{2 \times 2 \times 4}{3 \times 7}} \times 2 \text{ mm}$
 $= \frac{4}{4.58} \times 2 = 1.7 \text{ mm}$

30. The temperature, at which the root mean square velocity of hydrogen molecules equals their escape velocity from the earth, is closest to

[Boltzmann constant k_B = 1.38 × 10⁻²³ J/K

Avogadro Number $N_{\Delta} = 6.02 \times 10^{26}$ /kg

Radius of Earth : 6.4×10^6 m

Gravitational acceleration on Earth = 10 ms⁻²]

- (1) 10⁴ K
- (2) 650 K

Answer (1)

Sol.
$$V_{rms} = \sqrt{\frac{3RT}{M}} = 11.2 \times 10^3 \text{ m/s}$$

$$\Rightarrow T = \frac{M}{3R} \times (11.2 \times 10^3)^2$$

$$= \frac{2 \times 10^{-3}}{3 \times 8.3} \times 125.44 \times 10^6 \simeq 10^4 \text{ K}$$





1. 0.27 g of a long chain fatty acid was dissolved in 100 cm³ of hexane. 10 mL of this solution was added dropwise to the surface of water in a round watch glass. Hexane evaporates and a monolayer is formed. The distance from edge to centre of the watch glass is 10 cm. What is the height of the monolayer?

[Density of fatty acid = 0.9 g cm⁻³; π = 3]

- (1) 10⁻⁸ m (2) 10⁻⁴ m
- (3) 10^{-2} m (4) 10^{-6} m

Answer (4)

Sol. 0.27 gm in 100 ml of hexane

:. in 10 ml of aqueous solution only 0.027 gm acid is present

volume of 0.027 g acid $=\frac{0.027}{0.9}$ ml

:.
$$\pi r^2 h = \frac{0.027}{0.9}$$
 (given r = 10 cm, π = 3)

- :. $h = 10^{-4} cm$
 - = 10⁻⁶ m
- 2. The Mond process is used for the :
 - (1) purification of Zr and Ti
 - (2) extraction of Mo
 - (3) purification of Ni
 - (4) extraction of Zn

Answer (3)

Sol. Nickel is purified by Mond's process

$$\frac{\text{Ni}_{(\text{impure})} + 4\text{CO} \xrightarrow{330-350 \text{ k}} \text{Ni(CO)}_{4}}{\text{Ni(CO)}_{4} \xrightarrow{450-470 \text{ k}} \frac{\text{Ni}}{(\text{pure})} + 4\text{CO}}$$

- 3. The correct statement about ICI_5 and ICI_4^- is :
 - (1) ICI_5 is square pyramidal and ICI_4^- is tetrahedral.
 - (2) both are isostructural.
 - (3) ICI_5 is square pyramidal and ICI_4^- is square planar.
 - (4) ICI_5 is trigonal bipyramidal and ICI_4^- is tetrahedral.

Answer (3)

Sol. ICl_5 is sp^3d^2 hybridised (5 bond pairs, 1 lone pair)



 ICl_4^- is $sp^3 d^2$ hybridised (4 bond pairs, 2 lone pairs)



4. For the following reactions, equilibrium constants are given :

$$S(s) + O_2(g) \Longrightarrow SO_2(g); K_1 = 10^{52}$$

 $2S(s) + 3O_2(g) \Longrightarrow 2SO_3(g); K_2 = 10^{129}$

The equilibrium constant for the reaction,

$$2SO_2(g) + O_2(g) \Longrightarrow 2SO_3(g)$$
 is :

- (1) 10¹⁵⁴
- (2) 10²⁵
- (3) 1077
- (4) **10**¹⁸¹

Answer (2)

Sol. eqⁿ 1 S + O₂ \rightleftharpoons SO₂ K₁ = 10⁵² eqⁿ 2 2S + 3O₂ \rightleftharpoons 2SO₃ K₂ = 10¹²⁹ eqⁿ 3 2SO₂ + O₂ \rightleftharpoons 2SO₃ eqⁿ 3 = eqⁿ 2 - 2 (eqⁿ 1) = $\frac{10^{129}}{(10^{52})^2} = 10^{25}$



The radius of atom B is twice that of atom A. The unit cell edge length is 50% more in solid 2 than in 1. What is the approximate packing efficiency in solid 2?



(1) 45% (2) 65%

(3) 75% (4) 90%

Answer (4)

Sol. Volume occupied by atoms in solid 2

$$=\frac{4}{3}\pi r^{3}+\frac{4}{3}\pi (2r)^{3}=12 \pi r^{3}$$

relationship between edge length (a) and radius of atom (r)

$$=$$
 6r $=$ $\sqrt{3}$ a \Rightarrow a $=$ $\frac{6r}{\sqrt{3}}$

packing efficiency = $\frac{12\pi r^3}{\left(\frac{6r}{\sqrt{3}}\right)^3} \times 100 = 90\%$

- 6. The compound that inhibits the growth of tumors is
 - (1) cis-[Pt(Cl)₂(NH₃)₂]
 - (2) trans-[Pt(Cl)₂(NH₃)₂]
 - (3) cis-[Pd(Cl)₂(NH₃)₂]
 - (4) trans-[Pd(Cl)₂(NH₃)₂]

Answer (1)

Sol. Cis-platin is used as an anti-cancer drug.

$$\begin{array}{c} \mathbf{O} \quad \mathbf{H} \\ (1) \quad \left[(\mathbf{CH}_{2})_{4} - \mathbf{C} - \mathbf{N} \right]_{n} \\ (2) \quad \left[\mathbf{C} - (\mathbf{CH}_{2})_{6} - \mathbf{N} \right]_{n} \\ (3) \quad \left[(\mathbf{CH}_{2})_{6} - \mathbf{C} - \mathbf{N} \right]_{n} \\ (4) \quad \left[\mathbf{C} - (\mathbf{CH}_{2})_{5} - \mathbf{N} \right]_{n} \end{array}$$

Answer (4)

Sol. Nylon-6 is produced from caprolactam

$$\begin{array}{c} O \\ \downarrow \\ \mathsf{NH} \\ \downarrow \\ \mathsf{NH} \end{array} \rightarrow \begin{array}{c} \left[\begin{array}{c} O \\ \parallel \\ \mathsf{C} - (\mathsf{CH}_2)_5 - \mathsf{NH} \right]_n \\ \mathsf{Nylon-6} \end{array} \right]$$

- 8. Which one of the following alkenes when treated with HCl yields majorly an anti Markovnikov product ?
 - (1) $F_3C-CH=CH_2$
 - (2) $CH_3O-CH=CH_2$
 - (3) $H_2N-CH=CH_2$

(4)
$$CI-CH=CH_2$$

Answer (1)

Sol.
$$CF_3 - CH = CH_2$$

 \downarrow H^+
 $CF_3 - CH - CH_3 + CF_3 - CH_2 - CH_2$
 \downarrow H^+
 $CF_3 - CH - CH_3 + CF_3 - CH_2 - CH_2$
 \downarrow $(More stable)$
 $(Less stable due to -l effect of -CF_3)$
 $CF_3 - CH_2 - CH_2 - CH_2 - CH_2$

- 9. The strength of 11.2 volume solution of H_2O_2 is [Given that molar mass of H = 1 g mol⁻¹ and O = 16 g mol⁻¹]
 - (1) 13.6%
 - (2) 1.7%
 - (3) 3.4%
 - (4) 34%



301. 11.2 V 01 $\Pi_2 U_2$

$$H_2O_2 \longrightarrow H_2O + \frac{1}{2}O_2$$

11.2 L of O_2 at STP = 0.5 mol

It means 1 L of given H_2O_2 solution consist 1 mole of H_2O_2 (i.e., 34 g)

strength = $\frac{34}{1000}$ × 100 = 3.4%

- 10. The covalent alkaline earth metal halide (X = Cl, Br, I) is
 - (1) BeX₂
 - (2) SrX₂
 - (3) CaX₂
 - (4) MgX₂

Answer (1)

Sol. According to Fajan's rule, graeter the polarising power of cation greater would be the covalent character.

Since Be^{2+} has maximum polarising power among given cation, therefore, BeX_2 would be most covalent among given alkaline with metal halides.

- 11. Polysubstitution is a major drawback in :
 - (1) Reimer Tiemann reaction
 - (2) Acetylation of aniline
 - (3) Friedel Craft's acylation
 - (4) Friedel Craft's alkylation

Answer (4)

- Sol. Polysubstitution is a major drawback in Friedel Craft's alkylation
- 12. The IUPAC symbol for the element with atomic number 119 would be :

(1)	une		(2)	uun

(3) uue (4) unh

Answer (3)

Sol. Symbol for 1 is u

and for 9 is e

 \therefore IUPAC symbol for 119 is uue

temperature becomes 200 K. If $C_V = 28 \text{ J K}^{-1} \text{ mol}^{-1}$, calculate ΔU and ΔpV for this process. (R = 8.0 J K⁻¹ mol⁻¹)

Answer (4)

Sol. $\Delta U = n c_{vm} \Delta T = 5 \times 28 \times 100$ = 14 kJ $\Delta (PV) = nR(T_2 - T_1)$ = 5 × 8 × 100 = 4 kJ

- 14. The calculated spin-only magnetic moments (BM) of the anionic and cationic species of $[Fe(H_2O)_6]_2$ and $[Fe(CN)_6]$, respectively, are:
 - (1) 2.84 and 5.92 (2) 4.9 and 0
 - (3) 0 and 5.92 (4) 0 and 4.9

- Sol. $[Fe(H_2O)_6]_2$ $[Fe(CN)_6]$ $[Fe(H_2O)_6]^{2+}$ $[Fe(CN)_6]^{4-}$ 4 unpaired no unpaired electron electrons $\mu = 0$ $\mu = 4.9$
- 15. The major product of the following reaction is:







16. The major product in the following reaction is:







Answer (4)



removes the more acidic H-atom and the conjugate base of the given compound attacks at CH_3 group to give the final product shown above.

17. The major product of the following reaction is :



Answer (2)





18. The ion that has sp^3a^2 hybridization for the central atom, is :

(1) [ICI ₂]⁻	(2) [IF ₆]⁻
(3) [BrF₂] [−]	(4) [ICI₄] ⁻

Sol. Species	Hybridisation
	sp³ d
	sp³ d²
BrF_2^-	sp³ d
IF_6^-	sp³ a ^β



place :

$$\begin{split} & \operatorname{Fe}^{2+}(\operatorname{aq}) + \operatorname{Ag}^{+}(\operatorname{aq}) \to \operatorname{Fe}^{3+}(\operatorname{aq}) + \operatorname{Ag}(\operatorname{s}) \\ & \operatorname{Given that} \\ & \operatorname{E}^{o}_{\operatorname{Ag}^{+}/\operatorname{Ag}} = \operatorname{xV} \\ & \operatorname{E}^{o}_{\operatorname{Fe}^{2+}/\operatorname{Fe}} = \operatorname{yV} \\ & \operatorname{E}^{o}_{\operatorname{Fe}^{3+}/\operatorname{Fe}} = \operatorname{zV} \\ & (1) \times - \operatorname{y} \\ & (2) \times + \operatorname{y} - \operatorname{z} \\ & (3) \times + 2\operatorname{y} - 3\operatorname{z} \\ & (4) \times - \operatorname{z} \end{split}$$

Answer (3)

Sol.

$$\begin{array}{c} Ag^{+} + Fe^{+2} \longrightarrow Fe^{3+} + Ag \\ (aq) & (aq) \end{array}$$
$$F^{o}_{cell} = E^{o}_{Ag^{+}/Ag} - E^{o}_{Fe^{3+}/Fe^{2+}} \end{array}$$

To calculate

E^o_{Fe³⁺/Fe²⁺}

$$F_{e^{3^{+}}} \xrightarrow{F_{e^{2^{+}}}} F_{e^{2^{+}}} \xrightarrow{F_{e^{2^{+}}}} F_{e^{2^{+}}}$$

$$E_{F_{e^{3^{+}}/F_{e^{2^{+}}}}^{o}} = 3z - 2y$$

 $E^{o}_{Ag^{+}/Ag} = x$

 E_{cell}^{o} = x – 3z + 2y

20. Which of the following compounds will show the maximum 'enol' content?

(2) CH₃COCH₃

Maximum enol content

21. The major product obtained in the following reaction is:




are 0.5, 2, 35 and 40 kbar, respectively. The correct plot for the given data is :



Partial
pressure
$$y$$

 $(0, 0)$ x_{H_2O}

gas K_H
 w 0.5
 x 2
 y 35
 z 50

23. If p is the momentum of the fastest electron ejected from a metal surface after the irradiation of light having wavelength λ , then for 1.5 p momentum of the photoelectron, the wavelength of the light should be :

(Assume kinetic energy of ejected photoelectron to be very high in comparison to work function):



Answer (2)

Sol. In photoelectric effect,

$$\frac{hc}{\lambda}$$
 = w + KE of electron

It is given that KE of ejected electron is very high in comparison to w.

$$\frac{hc}{\lambda} = KE \implies \frac{hc}{\lambda} = \frac{P^2}{2m}$$

New wavelength

$$\frac{hc}{\lambda_1} = \frac{(1.5P)^2}{2m} \implies \lambda' = \frac{4}{9}\lambda$$

- 24. Fructose and glucose can be distinguished by:
 - (1) Fehling's test (2) Seliwanoff's test
 - (3) Barfoed's test (4) Benedict's test



- and ketose
- 25. Among the following molecules/ions,

Which one is diamagnetic and has the shortest bond length?

(1)	0 ₂	(2) O ₂ ²⁻
(3)	N ₂ ^{2–}	(4) C ₂ ²⁻

Answer (4)

Sol. Bond length
$$\sim \frac{1}{bondorder}$$

and diamagnetic species has no unpaired electron in their molecular orbitals.

	Bond order	Magnetic character
C ₂ ^{2–}	3	diamagnetic
N_{2}^{2-}	2	paramagnetic
0 ₂ ^{2–}	1	diamagnetic
O ₂	2	paramagnetic

 \therefore C₂²⁻ has least bond length and is diamagnetic

26. For a reaction scheme $A \xrightarrow{k_1} B \xrightarrow{k_2} C$, if the rate of formation of B is set to be zero then the concentration of B is given by :

(1)
$$k_1 k_2 [A]$$
 (2) $(k_1 - k_2) [A]$
(3) $\left(\frac{k_1}{k_2}\right) [A]$ (4) $(k_1 + k_2) [A]$

Answer (3)

Sol. $A \xrightarrow{k_1} B \xrightarrow{k_2} C$

 k_2

$$\frac{d[B]}{dt} = k_1[A] - k_2[B] = 0$$

[B] = $\frac{k_1[A]}{k_1}$

- 27. The maximum prescribed concentration of copper in drinking water is :
 - (1) 3 ppm (2) 0.05 ppm
 - (3) 0.5 ppm (4) 5 ppm

Answer (1)

Sol. Maximum prescribed concentration of Cu in drinking water is 3 ppm.



Answer (3)



29. The percentage composition of carbon by mole in methane is

(1) 80%	(2) 75%
(3) 20%	(4) 25%

Answer (3)

Sol. In CH₄

one atom of carbon among 5 atoms (1C + 4H atoms)

... Mole % of C =
$$\frac{1}{5} \times 100$$

= 20%

- 30. The statement that is INCORRECT about the interstitial compounds is
 - (1) They are chemically reactive.
 - (2) They are very hard.
 - (3) They have high melting points.
 - (4) They have metallic conductivity.

Answer (1)

Sol. Interstitial compounds are inert.



- The vector equation of plane through the line of intersection of the planes x + y + z = 1 and 2x + 3y + 4z = 5 which is perpendicular to the plane x - y + z = 0 is :
 - (1) $\vec{r} \cdot (\hat{i} \hat{k}) + 2 = 0$ (2) $\vec{r} \cdot (\hat{i} \hat{k}) 2 = 0$ (3) $\vec{r} \times (\hat{i} - \hat{k}) + 2 = 0$ (4) $\vec{r} \times (\hat{i} + \hat{k}) + 2 = 0$

Answer (1)

Sol. Equation of the plane passing through the line of intersection of x + y + z = 1 and 2x + 3y + 4z= 5 is

$$(2x + 3y + 4z - 5) + \lambda (x + y + z - 1) = 0$$

$$(2 + \lambda) x + (3 + \lambda) y + (4 + \lambda) z + (-5 - \lambda) = 0 \dots(i)$$
(i) is perpendicular to $x - y + z = 0$

$$\Rightarrow (2 + \lambda) (1) + (3 + \lambda) (-1) + (4 + \lambda) (1) = 0$$

$$2 + \lambda - 3 - \lambda + 4 + \lambda = 0$$

$$\lambda = -3$$

$$\Rightarrow \text{ Equation of required plane is}$$

$$-x + z - 2 = 0$$

$$\rightarrow$$
 X - Z + Z = 0

$$\Rightarrow \vec{\mathbf{r}} \cdot (\hat{\mathbf{i}} - \hat{\mathbf{k}}) + \mathbf{2} = \mathbf{0}$$

- 2. Which one of the following statements is not a tautology?
 - (1) $(p \land q) \rightarrow (\sim p) \lor q$
 - (2) $(p \land q) \rightarrow p$

(3)
$$(p \lor q) \rightarrow (p \lor (~q))$$

(4)
$$p \rightarrow (p \lor q)$$

Answer (3)

Sol. By help of truth table :

·										
р	q	~q	pv~q	~p	p∧~q	pvq	p→pvq	p∧q	(p∧q)→p	~pvq
Т	т	F	Т	F	F	Т	т	Т	т	т
Т	F	т	т	F	т	Т	т	F	т	F
F	т	F	F	т	F	Т	т	F	т	т
F	F	т	Т	Т	F	F	т	F	Т	т
	$(\mathbf{p} \wedge \mathbf{q}) \rightarrow (\mathbf{\sim} \mathbf{p}) \mathbf{vq} (\mathbf{pvq}) \rightarrow (\mathbf{pv}(\mathbf{\sim} \mathbf{q})) $									
	Т						т			
	т						Т			
	Т				F					
	Т					т				

3. The number of four-digit numbers strictly greater than 4321 that can be formed using the digits 0, 1, 2, 3, 4, 5 (repetition of digits is allowed) is :

(1) 360	(2) 306
(3) 288	(4) 310

Answer (4)

Sol. 0, 1, 2, 3, 4, 5

$$5 + + + + = 6 \times 6 \times 6 = 216$$

$$4 + 5 + + + = 6 \times 6 = 36$$

$$4 + 4 + + + = 6 \times 6 = 36$$

$$4 + 4 + + + = 6 \times 6 = 36$$

$$4 + 4 + + + = 3 \times 6 = 18$$

$$(5/4/3) + + + = 4$$

$$4 + 3 + + + = 4$$

- ⇒ Required numbers = 216 + 36 + 36 + 18 + 4 = 310
- 4. In an ellipse, with centre at the origin, if the difference of the lengths of major axis and minor axis is 10 and one of the foci is at

 $(0, 5\sqrt{3})$, then the length of its latus rectum is :

(1) 5	(2)	6
-------	-----	---

(3) 8 (4) 10

Answer (1)

Sol. : Focus is $(0, 5\sqrt{3}) \Rightarrow |\mathbf{b}| > |\mathbf{a}|$ Let $\mathbf{b} > \mathbf{a} > 0$ foci $(0, \pm \mathbf{b}\mathbf{e})$ $\mathbf{a}^2 = \mathbf{b}^2 - \mathbf{b}^2\mathbf{e}^2 \Rightarrow \mathbf{b}^2\mathbf{e}^2 = \mathbf{b}^2 - \mathbf{a}^2$

 $be = \sqrt{b^2 - a^2}$



$$\Rightarrow b - a = 5 \qquad \dots(ii)$$

From (i) and (ii)
$$b + a = 15 \qquad \dots(iii)$$
$$\Rightarrow b = 10, a = 5$$

Length of L·R· = $\frac{2a^2}{b} = \frac{50}{10} = 5$

5. A student scores the following marks in five tests : 45, 54, 41, 57, 43. His score is not known for the sixth test. If the mean score is 48 in the six tests, then the standard deviation of the marks in six tests is :

(1)
$$\frac{100}{\sqrt{3}}$$
 (2) $\frac{10}{\sqrt{3}}$
(3) $\frac{100}{3}$ (4) $\frac{10}{3}$

Answer (2)

Sol.
$$\bar{\mathbf{x}} = \frac{41+45+54+57+43+\mathbf{x}}{6} = 48$$

 $\mathbf{x} + 240 = 288$
 $\mathbf{x} = 48$
 $\sigma^2 = \frac{1}{6} \begin{bmatrix} (48-41)^2 + (48-45)^2 + (48-54)^2 \\ + (48-57)^2 + (48-43)^2 + (48-48)^2 \end{bmatrix}$
 $= \frac{1}{6} (49+9+36+81+25)$
 $= \frac{200}{6} = \frac{100}{3}$
 $\sigma = \frac{10}{\sqrt{3}}$

6. The tangent and the normal lines at the point $(\sqrt{3}, 1)$ to the circle $x^2 + y^2 = 4$ and the x-axis form a triangle. The area of this triangle (in square units) is :

(1)
$$\frac{2}{\sqrt{3}}$$
 (2) $\frac{4}{\sqrt{3}}$
(3) $\frac{1}{3}$ (4) $\frac{1}{\sqrt{3}}$

Answer (1)

y
P(
$$\sqrt{3}, 1$$
)
 $x^2 + y^2 = 4$
 \therefore Coordinate of $A = \left(\frac{4}{\sqrt{3}}, 0\right)$
Area $= \frac{1}{2} \times OA \times PM$
 $= \frac{1}{2} \times \frac{4}{\sqrt{3}} \times 1 = \frac{2}{\sqrt{3}}$ Square units
If the fourth term in the binomial expansion
of $\left(\sqrt{\frac{1}{x^{1+\log_{10}x}}} + x^{\frac{1}{12}}\right)^6$ is equal to 200, and
 $x > 1$, then the value of x is :
(1) 10 (2) 10³
(3) 100 (4) 10⁴

Answer (1)

7.

 $\sqrt{3}x + y = 4$

Sol.
$$T_4 = {}^6C_3 \left(\sqrt{x^{\left(\frac{1}{1+\log_{10} x}\right)}} \right)^3 \left(\frac{1}{x^{12}} \right)^3 = 200$$

 $\Rightarrow \left[20x^{\frac{3}{2(1+\log_{10} x)}} \cdot x^{\frac{1}{4}} = 200 \right]$
 $x^{\frac{1}{4} + \frac{3}{2(1+\log_{10} x)}} = 10$

Taking \log_{10} on both sides and put $\log_{10} x = t$

$$\left(\frac{1}{4} + \frac{3}{2(1+t)}\right)t = 1$$

$$\left(\frac{(1+t)+6}{4(1+t)}\right) \times t = 1 \implies t^2 + 7t = 4 + 4t$$

$$t^2 + 3t - 4 = 0 \implies t^2 + 4t - t - 4 = 0$$

$$\Rightarrow t(t+4) - 1 (t+4) = 0$$

$$\Rightarrow t = 1 \text{ or } t = -4$$

$$\log_{10} x = 1$$

$$\Rightarrow x = 10 \text{ or if } \log_{10} x = -4$$

$$\Rightarrow x = 10^{-4}$$

Note: There seems a printing error in this question in the original question paper.



first quadrant, passes through the point :

(1)
$$\left(\frac{3}{4}, \frac{7}{4}\right)$$
 (2) $\left(-\frac{1}{3}, \frac{4}{3}\right)$
(3) $\left(\frac{1}{4}, \frac{3}{4}\right)$ (4) $\left(-\frac{1}{4}, \frac{1}{2}\right)$

Answer (1)

Sol. Intersection point of

$$x^{2} + y^{2} = 5, \quad y^{2} = 4x$$

$$\Rightarrow x^{2} + 4x - 5 = 0$$

$$\Rightarrow x^{2} + 5x - x - 5 = 0$$

$$\Rightarrow x(x + 5) - 1(x + 5) = 0$$

$$\therefore x = 1, -5$$

Intersection point in 1^{st} quadrant be (1, 2) equation of tangent to $y^2 = 4x$ at (1, 2) is

$$y \times 2 = 2(x + 1)$$

$$\Rightarrow y = x + 1$$

$$\Rightarrow x - y + 1 = 0 \qquad \dots(i)$$

$$\left(\frac{3}{4}, \frac{7}{4}\right) \text{ lies on (i)}$$

9. If three distinct numbers a, b, c are in G.P. and the equations $ax^2 + 2bx + c = 0$ and $dx^2 + 2ex + f = 0$ have a common root, then which one of the following statements is correct?

(1) d, e, f are in A.P. (2)
$$\frac{d}{a}$$
, $\frac{e}{b}$, $\frac{f}{c}$ are in G.P.
(3) $\frac{d}{a}$, $\frac{e}{b}$, $\frac{f}{c}$ are in A.P. (4) d, e, f are in G.P.

Answer (3)

Sol. Since a, b, c are in G.P.

⇒ b² = ac
Given, ax² + 2bx + c = 0
⇒ ax² + 2
$$\sqrt{ac}$$
 x + c = 0
⇒ $(\sqrt{a}$ x + $\sqrt{c})^2$ = 0
⇒ x = $-\sqrt{\frac{c}{a}}$

 \therefore ax² + 2bx + c = 0 and dx² + 2ex + f = 0 have common root

$$\forall a$$

$$\Rightarrow d \cdot \frac{c}{a} + 2e\left(-\sqrt{\frac{c}{a}}\right) + f = 0$$

$$\frac{d}{a} - \frac{2e}{\sqrt{ac}} + \frac{f}{c} = 0$$

$$\Rightarrow \frac{d}{a} - \frac{2e}{b} + \frac{f}{c} = 0$$

$$\Rightarrow \frac{2e}{b} = \frac{d}{a} + \frac{f}{c}$$

$$\Rightarrow \frac{d}{a}, \frac{e}{b}, \frac{f}{c} \text{ are in A.P.}$$
10. If $z = \frac{\sqrt{3}}{2} + \frac{i}{2}(i = \sqrt{-1})$, then
$$(1 + iz + z^5 + iz^8)^9 \text{ is equal to :}$$

$$(1) 0 \qquad (2) (-1 + 2i)^9$$

$$(3) - 1 \qquad (4) 1$$

Answer (3)

Sol.
$$z = \frac{\sqrt{3}}{2} + \frac{i}{2} = -i\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) = -i\omega$$

where ω is not real cube root of unity

$$\Rightarrow (1 + iz + z^5 + iz^8)^9 = (1 + \omega - i\omega^2 + i\omega^2)^9$$
$$= (1 + \omega)^9$$
$$= (-\omega^2)^9$$
$$= -\omega^{18}$$
$$= -1$$

11. The number of integral values of m for which the equation $(1 + m^2)x^2 - 2(1 + 3m)x + (1 + 8m) = 0$ has no real root is :

(1) Infinitely many
(2) 3
(3) 2
(4) 1

Answer (1)

Sol. $(1 + m^2)x^2 - 2(1 + 3m)x + (1 + 8m) = 0$

equation has no real solution

 $\Rightarrow D < 0$ $4(1 + 3m)^{2} < 4(1 + m^{2}) (1 + 8m)$ $1 + 9m^{2} + 6m < 1 + 8m + m^{2} + 8m^{3}$



$$2m(4m^2 - 4m + 1) > 0$$

 $2m(2m - 1)^2 > 0$
 $m > 0, m \neq \frac{1}{2}$

- \Rightarrow number of integral values of m are infinitely many.
- 12. Let the numbers 2, b, c be in an A.P. and

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & b & c \\ 4 & b^2 & c^2 \end{bmatrix}.$$
 If det(A) \in [2, 16], then c lies

in the interval :

(1) [2, 3) (2) $(2 + 2^{3/4}, 4)$ (3) [3, 2 + 2^{3/4}] (4) [4, 6]

Answer (4)

Sol.
$$|A| = \begin{vmatrix} 1 & 1 & 1 \\ 2 & b & c \\ 4 & b^2 & c^2 \end{vmatrix}$$

 $c_2 \rightarrow c_2 - c_1, c_3 \rightarrow c_3 - c_1$
 $|A| = \begin{vmatrix} 1 & 0 & 0 \\ 2 & b - 2 & c - 2 \\ 4 & (b - 2)(b + 2) & (c - 2)(c + 2) \end{vmatrix}$
 $= (b - 2)(c - 2)(c - b)$
2, b, c are in A.P. $\Rightarrow (b - 2) = (c - b) = d, c - 2 = 2d$
 $\Rightarrow |A| = d.2d.d = 2d^3$
 $\because |A| \in [2, 16] \Rightarrow 1 \le d^3 \le 8 \Rightarrow 1 \le d \le 2$

$$4 \leq 2d + 2 \leq 6 \implies 4 \leq c \leq 6$$

13. The minimum number of times one has to toss a fair coin so that the probability of observing at least one head is at least 90% is :

(1) 5	(2) 2
(3) 4	(4) 3

$$P(x \ge 1) \ge \frac{9}{10}$$

$$1 - P(x = 0) \ge \frac{9}{10}$$

$$1 - {}^{n}C_{0}\left(\frac{1}{2}\right)^{n} \ge \frac{9}{10}$$

$$\frac{1}{2^{n}} \le 1 - \frac{9}{10} \Longrightarrow \frac{1}{2^{n}} \le \frac{1}{10}$$

$$2^{n} \ge 10 \implies n \ge 4$$

$$\Rightarrow n_{min} = 4$$
Let $\vec{a} = 3\hat{i} + 2\hat{j} + x\hat{k}$ and $\vec{b} = \hat{i} - \hat{j} + \hat{k}$, for some real x. Then $|\vec{a} \times \vec{b}| = r$ is possible if :
$$(1) \ 3\sqrt{\frac{3}{2}} < r < 5\sqrt{\frac{3}{2}}$$

(2)
$$\sqrt{\frac{3}{2}} < r \le 3\sqrt{\frac{3}{2}}$$

(3) $0 < r \le \sqrt{\frac{3}{2}}$
(4) $r \ge 5\sqrt{\frac{3}{2}}$

Answer (4)

14.

Sol.
$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 2 & x \\ 1 & -1 & 1 \end{vmatrix}$$

$$= (2+x)\hat{i} + (x-3)\hat{j} - 5\hat{k}$$
$$|\vec{a} \times \vec{b}| = r = \sqrt{(2+x)^2 + (x-3)^2 + (-5)^2}$$
$$\Rightarrow r = \sqrt{4 + x^2 + 4x + x^2 + 9 - 6x + 25}$$
$$= \sqrt{2x^2 - 2x + 38} = \sqrt{2(x^2 - x + \frac{1}{4}) + 38 - \frac{1}{2}}$$
$$= \sqrt{2(x - \frac{1}{2})^2 + \frac{75}{2}}$$
$$\Rightarrow r \ge \sqrt{\frac{75}{2}} \Rightarrow r \ge 5\sqrt{\frac{3}{2}}$$



(1)
$$2 - \frac{3}{2^{17}}$$

(2) $1 - \frac{11}{2^{20}}$
(3) $2 - \frac{21}{2^{20}}$
(4) $2 - \frac{11}{2^{19}}$

Answer (4)

Sol. $S = \sum_{k=1}^{20} k \cdot \frac{1}{2^k}$ $S = \frac{1}{2} + 2 \cdot \frac{1}{2^2} + 3 \cdot \frac{1}{2^3} + \dots + 20 \cdot \frac{1}{2^{20}}$ $\frac{1}{2}S = \frac{1}{2^2} + 2 \cdot \frac{1}{2^3} + \dots + 19 \cdot \frac{1}{2^{20}} + 20 \cdot \frac{1}{2^{21}}$ On subtracting $\frac{S}{2} = \left(\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots + \frac{1}{2^{20}}\right) - 20 \cdot \frac{1}{2^{21}}$ $= \frac{\frac{1}{2}\left(1 - \frac{1}{2^{20}}\right)}{1 - \frac{1}{2}} - 20 \cdot \frac{1}{2^{21}} = 1 - \frac{1}{2^{20}} - 10 \cdot \frac{1}{2^{20}}$ $\frac{S}{2} = 1 - 11 \cdot \frac{1}{2^{20}} \Rightarrow S = 2 - 11 \cdot \frac{1}{2^{19}} = 2 - \frac{11}{2^{19}}$

16. Two vertical poles of heights, 20 m and 80 m stand apart on a horizontal plane. The height (in meters) of the point of intersection of the lines joining the top of each pole to the foot of the other, from this horizontal plane is :

(1) 16	(2) 18
--------	--------

(3) 15 (4) 12

Answer (1)



$$y = \frac{80}{x_1}x$$
 ...(i)
 $\frac{x}{x_1} + \frac{y}{20} = 1$...(ii)

For intersection point, from equations (i) and (ii)

$$\frac{y}{80} + \frac{y}{20} = 1$$

$$\Rightarrow y + 4y = 80$$

$$\Rightarrow y = 16 m$$

$$\Rightarrow \text{ Height of intersection point is 16 m}$$

17. Let $f : [-1, 3] \rightarrow R$ be defined as

$$f(x) = \begin{cases} |x| + [x], & -1 \le x < 1 \\ x + |x|, & 1 \le x < 2 \end{cases}$$

 $|\mathbf{x}+[\mathbf{x}], \mathbf{2} \le \mathbf{x} \le \mathbf{3},$

where [t] denotes the greatest integer less than or equal to t. Then, f is discontinuous at :

- (1) Only one point
- (2) Only two points
- (3) Only three points
- (4) Four or more points

Answer (3)

Sol.
$$f(x) = \begin{cases} |x| + [x], & -1 \le x < 1 \\ x + |x|, & 1 \le x < 2 \\ x + [x], & 2 \le x \le 3 \end{cases}$$

$$=\begin{cases} -x-1, & -1 \le x < 0 \\ x, & 0 \le x < 1 \\ 2x, & 1 \le x < 2 \\ x+2, & 2 \le x < 3 \\ 6, & x = 3 \end{cases}$$

$$\Rightarrow f(-1) = 0, f(-1^+) = 0$$

f(0⁻) = -1, f(0) = 0, f(0^+) = 0
f(1⁻) = 1, f(1) = 2, f(1^+) = 2
f(2⁻) = 4, f(2) = 4, f(2^+) = 4
f(3⁻) = 5, f(3) = 6
f(x) is discontinuous at x = {0, 1,



3}

y = y(x) at any point (x,y) is $\frac{1}{x^2}$. If the curve passes through the centre of the circle $x^2 + y^2 - 2x - 2y = 0$, then its equation is (1) $x \log_e |y| = -2(x - 1)$ (2) $x \log_e |y| = x - 1$ (3) $x \log_e |y| = 2(x - 1)$ (4) $x^2 \log_e |y| = -2(x - 1)$

Answer (3)

Sol.
$$\frac{dy}{dx} = \frac{2y}{x^2} \implies \int \frac{dy}{y} = 2\int \frac{dx}{x^2}$$
$$\implies \ln|y| = -\frac{2}{x} + C \dots(i)$$
(i) passes through (1, 1)
$$\implies C = 2$$
$$\implies \ln|y| = -\frac{2}{x} + 2$$
$$x \ln|y| = -2 + 2x$$
$$x \ln|y| = -2(1-x) = 2(x-1)$$

19. Suppose that the points (h, k), (1, 2) and (-3, 4) lie on the line L_1 . If a line L_2 passing through the points (h, k) and (4, 3) is perpendicular to

L_1 , then $\frac{k}{h}$ equals	
(1) 3	(2) - <mark>1</mark> 7
(3) 0	(4) $\frac{1}{3}$

Answer (4)

Sol. (h, k), (1, 2) and (-3, 4) and collinear

$$\begin{vmatrix} h & k & 1 \\ 1 & 2 & 1 \\ -3 & 4 & 1 \end{vmatrix} = 0 \Rightarrow -2h - 4k + 10 = 0$$

$$\Rightarrow h + 2k = 5 \qquad \dots(i)$$

$$m_{L_1} = \frac{4 - 2}{-3 - 1} = \frac{2}{-4} = -\frac{1}{2} \Rightarrow m_{L_2} = 2$$

$$\Rightarrow m_{L_2} = \frac{k - 3}{h - 4} = 2 \Rightarrow k - 3 = 2h - 8$$

$$2h - k = 5 \qquad \dots(ii)$$

from (i) and (ii)
$$h = 3, k = 1 \Rightarrow \frac{k}{h} = \frac{1}{3}$$

 $A(\lambda)$: A(4) = 2 : 5, then λ equals



Answer (4)

Sol.



21. If a point R(4, y, z) lies on the line segment joining the points P(2, -3, 4) and Q(8, 0, 10), then the distance of R from the origin is

(1)	√53	(2)	2√21

(3) 6 (4) $2\sqrt{14}$

Answer (4)

Sol. P, Q, R are collinear

$$\Rightarrow \overrightarrow{PR} = \lambda \overrightarrow{PQ}$$

$$2\hat{i} + (y+3)\hat{j} + (z-4)\hat{k} = \lambda[6\hat{i}+3\hat{j}+6\hat{k}]$$

$$\Rightarrow 6\lambda = 2, y+3 = 3\lambda, z-4 = 6\lambda$$

$$\Rightarrow \lambda = \frac{1}{3}, y = -2, z = 6$$

$$\Rightarrow \text{ point R } (4, -2, 6)$$

$$\Rightarrow OR = \sqrt{(4)^2 + (-2)^2 + (6)^2} = \sqrt{16 + 4 + 36}$$

$$= \sqrt{56} = 2\sqrt{14}$$



smallest, then a ratio of lengths of the sides of this triangle is

(1)
$$4:5:6$$

(2) $3:4:5$
(3) $5:9:13$
(4) $5:6:7$
Answer (1)
Sol. Let $a > b > c$
 $\therefore A = 2C$
 $\Rightarrow A + B + C = \pi$

$$\Rightarrow \mathbf{B} = \pi - \mathbf{3C} \qquad \dots (\mathbf{i})$$

Sol

- \Rightarrow sinA + sinC = 2sinB ...(ii)
- \Rightarrow sinA = sin(2C), sinB = sin3C
- \Rightarrow From (ii), sin2C + sinC = 2sin3C $(2\cos C + 1)\sin C = 2\sin C (3 - 4 \sin^2 C)$
- $\Rightarrow 2\cos C + 1 = 6 8(1 \cos^2 C)$
- \Rightarrow 8cos²C 2cosC 3 =0
- \Rightarrow cosC = $\frac{3}{4}$ or cosC = $-\frac{1}{2}$
- ∴ C is acute angle

$$\Rightarrow \cos C = \frac{3}{4}, \sin A = 2 \sin C \cos C = 2 \times \frac{\sqrt{7}}{4} \times \frac{3}{4}$$
$$\Rightarrow \sin C = \frac{\sqrt{7}}{4}, \sin B = \frac{3\sqrt{7}}{4} - \frac{4\sqrt{7}}{4} \times \frac{7}{16} = \frac{5\sqrt{7}}{16}$$

- \Rightarrow sinA : sinB : sinC :: a : b : c is 6 : 5 : 4
- 23. The height of a right circular cylinder of maximum volume inscribed in a sphere of radius 3 is

24.

- (1) $\frac{2}{3}\sqrt{3}$
- (2) $2\sqrt{3}$
- (3) $\sqrt{3}$
- (4) $\sqrt{6}$

Answer (2)

Sol. Let radius of base and height of cylinder be r and h respectively.

$$\therefore r^{2} + \frac{h^{2}}{4} = 9 \qquad \dots(i)$$

$$\therefore Volume of cylinder$$

$$V = \pi r^{2}h$$

$$V = \pi h \left(9 - \frac{h^{2}}{4}\right)$$

$$V = 9\pi h - \frac{\pi}{4}h^{3}$$

$$\therefore \frac{dV}{dh} = 9\pi - \frac{3}{4}\pi h^{2}$$
For maxima/minima

$$\frac{dV}{dh} = 0$$

$$\Rightarrow h = \sqrt{12}$$
and
$$\frac{d^{2}V}{dh^{2}} = -\frac{3}{2}\pi h$$

$$\therefore \left(\frac{d^{2}V}{dh^{2}}\right)_{n=\sqrt{12}} < 0$$

$$\Rightarrow Volume is maximum when h = 2\sqrt{3}$$
24. If the system of linear equations

$$x - 2y + kz = 1$$

$$2x + y + z = 2$$

$$3x - y - kz = 3$$
has a solution (x, y, z), z \neq 0, then (x, y) lies on
the straight line whose equation is
(1) $3x - 4y - 4 = 0$
(2) $3x - 4y - 1 = 0$
(3) $4x - 3y - 1 = 0$
(4) $4x - 3y - 4 = 0$



$$\Delta = \begin{vmatrix} 1 & -2 & k \\ 2 & 1 & 1 \\ 3 & -1 & -k \end{vmatrix} = 1(-k+1) + 2(-2k-3) + k(-2-3)$$

$$= -k+1 - 4k - 6 - 5k = -10k - 5 = -5(2k+1)$$

$$\Delta_{1} = \begin{vmatrix} 1 & -2 & k \\ 2 & 1 & 1 \\ 3 & -1 & -k \end{vmatrix} = -5(2k+1)$$

$$\Delta_{2} = \begin{vmatrix} 1 & 1 & k \\ 2 & 2 & 1 \\ 3 & 3 & -k \end{vmatrix} = 0, \Delta_{3} = \begin{vmatrix} 1 & -2 & 1 \\ 2 & 1 & 2 \\ 3 & -1 & 3 \end{vmatrix} = 0$$

$$\Rightarrow \Delta = 0$$

$$\Rightarrow \Delta = 0$$

$$\Rightarrow k = -\frac{1}{2}$$

$$\therefore \text{ System of equation has infinite many solutions.}$$
Let $z = \lambda \neq 0$ then $X = \frac{10 - 3\lambda}{10}$ and $Y = -\frac{2\lambda}{5}$

$$\therefore (x, y) \text{ must lie on line $4x - 3y - 4 = 0$
25. If $f(1) = 1$, $f'(1) = 3$, then the derivative of $f(f(f(x))) + (f(x))^{2}$ at $x = 1$ is:
(1) 33 (2) 12
(3) 9 (4) 15
Answer (1)
Sol. Let $g(x) = f(f(f(x))) + (f(x))^{2}$
On differentiating both sides w.r.t. x we get $g'(x) = f'(f(f(x))) f'(x) + 2f(x) f'(x)$
 $g'(1) = f'(f(f(1))) f'(f(1)) f'(1) + 2f(1) f'(1)$
 $= 3 \times 3 \times 3 + 2 \times 1 \times 3 = 27 + 6 = 33$
26. Let $f(x) = \int_{0}^{x} g(t) dt$ (2) $2 \int_{5}^{x+5} g(t) dt$
(3) $\int_{5}^{x} g(t) dt$ (2) $2 \int_{5}^{x+5} g(t) dt$
Answer (1)$$

$$g(-x) = g(x), \qquad \dots(ii)$$

$$f(x + 5) = g(x) \qquad \dots(iii)$$
From (i)

$$f'(x) = g(x)$$
Let $I = \int_{0}^{x} f(t) dt,$
Put $t = \lambda - 5$

$$\Rightarrow I = \int_{5}^{x+5} f(\lambda - 5) d\lambda$$

$$\therefore f(x + 5) = g(x)$$

$$\Rightarrow f(-x + 5) = g(-x) = g(x) \qquad \dots(iv)$$

$$I = \int_{5}^{x+5} f(\lambda - 5) d\lambda,$$

$$(\because f(0) = 0, g(x) \text{ is even } \Rightarrow f(x) \text{ is odd})$$

$$I = -\int_{5}^{x+5} g(\lambda) d\lambda = \int_{x+5}^{5} g(t) dt \qquad (from (iv))$$
Let $f(x) = a^{x} (a > 0)$ be written as $f(x) = f_{1}(x) + f_{2}(x),$
where $f_{1}(x)$ is an even function and $f_{2}(x)$ is an odd function. Then $f_{1}(x + y) + f_{1}(x - y)$ equals:
(1) $2f_{1}(x)f_{1}(y) \qquad (2) $2f_{1}(x + y)f_{1}(x - y)$$

(1)
$$2f_1(x)f_1(y)$$
 (2) $2f_1(x + y)f_1(x - y)$
(3) $2f_1(x + y)f_2(x - y)$ (4) $2f_1(x)f_2(y)$

Answer (1)

27.

Sol.
$$f(x) = a^{x} = \left(\frac{a^{x} + a^{-x}}{2}\right) + \left(\frac{a^{x} - a^{-x}}{2}\right)$$

where $f_{1}(x) = \frac{a^{x} + a^{-x}}{2}$ is even function
 $f_{2}(x) = \frac{a^{x} - a^{-x}}{2}$ is odd function
 $\Rightarrow f_{1}(x + y) + f_{1}(x - y)$
 $= \left(\frac{a^{x+y} + a^{-x-y}}{2}\right) + \left(\frac{a^{x-y} + a^{-x+y}}{2}\right)$
 $= \frac{1}{2} \left[a^{x}(a^{y} + a^{-y}) + a^{-x}(a^{y} + a^{-y})\right]$
 $= \frac{(a^{x} + a^{-x})(a^{y} + a^{-y})}{2}$
 $= 2f_{1}(x).f_{1}(y)$



$$\int x^{3}(1+x^{6})^{\frac{2}{3}}$$

constant of integration, then the function f(x) is equal to:

(1)
$$-\frac{1}{2x^3}$$
 (2) $\frac{3}{x^2}$
(3) $-\frac{1}{6x^3}$ (4) $-\frac{1}{2x^2}$

Answer (1)

Sol.
$$I = \int \frac{dx}{x^3 (1 + x^6)^{2/3}} = \int \frac{dx}{x^7 (1 + x^{-6})^{2/3}}$$

Put $1 + x^{-6} = t^3$
 $\Rightarrow -6x^{-7}dx = 3t^2 dt$
 $\Rightarrow \frac{dx}{x^7} = -\frac{1}{2}t^2 dt$
 $\Rightarrow I = \int -\frac{1}{2}\frac{t^2 dt}{t^2}$
 $= -\frac{1}{2}t + C$
 $= -\frac{1}{2}(1 + x^{-6})^{\frac{1}{3}} + C$
 $= -\frac{1}{2}\frac{(1 + x^6)^{\frac{1}{3}}}{x^2} + C$
 $= -\frac{1}{2x^3}x(1 + x^6)^{\frac{1}{3}} + C$

- 29. If the eccentricity of the standard hyperbola passing through the point (4, 6) is 2, then the equation of the tangent to the hyperbola at (4, 6) is:
 - (1) 2x 3y + 10 = 0 (2) x 2y + 8 = 0(3) 3x - 2y = 0 (4) 2x - y - 2 = 0

Answer (4)

Sol. Let equation of hyperbola be

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \qquad \dots(i)$$

$$\Rightarrow b^2 = a^2(e^2 - 1)$$

$$\therefore e = 2 \qquad \Rightarrow \qquad b^2 = 3a^2 \qquad \dots(ii)$$

$$\Rightarrow \frac{16}{a^2} - \frac{36}{b^2} = 1 \qquad \dots (iii)$$

From (ii) and (iii)

$$a^2 = 4$$
, $b^2 = 12$
 \Rightarrow Equation of hyperbola is $\frac{x^2}{4} - \frac{y^2}{12} = 1$

Equation of tangent to the hyperbola at (4, 6) is

$$\frac{4x}{4} - \frac{6y}{12} = 1$$
$$\Rightarrow x - \frac{y}{2} = 1$$
$$\Rightarrow 2x - y = 2$$

30. Let $f: R \rightarrow R$ be a differentiable function satisfying f'(3) + f'(2) = 0.

Then
$$\lim_{x \to 0} \left(\frac{1 + f(3 + x) - f(3)}{1 + f(2 - x) - f(2)} \right)^{\frac{1}{x}}$$
 is equal to:
(1) e (2) 1
(3) e^{2} (4) e^{-1}

Answer (2)

Sol. I =
$$\lim_{x \to 0} \left(\frac{1 + f(3 + x) - f(3)}{1 + f(2 - x) - f(2)} \right)^{\frac{1}{x}}$$

$$\Rightarrow$$
 I = e^{I₁}, where

$$I_{1} = \lim_{x \to 0} \left(\left(\frac{1 + f(3 + x) - f(3)}{1 + f(2 - x) - f(2)} - 1 \right) \frac{1}{x} \right)$$

$$= \lim_{x \to 0} \left(\left(\frac{f(3+x) - f(3) - f(2-x) + f(2)}{1 + f(2-x) - f(2)} \right) \frac{1}{x} \right)$$

form :
$$\frac{0}{0}$$

Using L.H. Rule

$$I_{1} = \lim_{x \to 0} \left(\frac{f'(3+x) + f'(2-x)}{1} \right) \cdot \lim_{x \to 0} \left(\frac{1}{1 + f(2-x) - f(2)} \right)$$
$$= f'(3) + f'(2) = 0$$
$$\Rightarrow I = e^{I_{1}} = 1$$



Answers & Solutions

for

JEE (MAIN)-2019 (Online) Phase-2

(Physics, Chemistry and Mathematics)

Time : 3 hrs.

M.M.: 360

Important Instructions :

- 1. The test is of **3 hours** duration.
- 2. The Test Booklet consists of **90** questions. The maximum marks are **360**.
- 3. There are *three* parts in the question paper A, B, C consisting of **Physics**, **Chemistry** and **Mathematics** having 30 questions in each part of equal weightage.
- 4. Each question is allotted 4 (four) marks for each correct response. ¼ (one-fourth) marks will be deducted for indicating incorrect response of each question. No deduction from the total score will be made if no response is indicated for an item in the answer sheet.
- 5. There is only one correct response for each question.



1. Determine the charge on the capacitor in the following circuit :



Answer (1)

Sol. At steady state current through capacitor is zero.

$$V_{c} = V_{1}$$

$$V_{1} = \frac{5}{6} \times V_{0} \times \frac{3}{9}$$



$$V_1 = \frac{5 \times 72 \times 3}{6 \times 9} = 20 \text{ V}$$

 $Q_1 = CV_1$ = 200 µC

2. The magnetic field of a plane electromagnetic wave is given by :

 $\vec{\mathbf{B}} = \vec{\mathbf{B}}_0 \hat{\mathbf{i}} [\cos(kz - \omega t) + \mathbf{B}_1 \hat{\mathbf{j}} \cos(kz + \omega t)]$

where $B_0 = 3 \times 10^{-5} \text{ T}$ and $B_1 = 2 \times 10^{-6} \text{ T}$.

The rms value of the force experienced by a stationary charge $Q = 10^{-4}$ C at z =0 is closest to:

(1)	0.6 N	(2)	0.9	Ν
(3)	3 × 10 ⁻² N	(4)	0.1	Ν

Answer (1)

Sol. $\left| \vec{E}_1 \right| = CB_1$

$$\left| \vec{\mathsf{E}}_2 \right| = \mathsf{CB}_2$$

Also $\vec{E}_1 \perp E_2$

$$F_{net} = \frac{\theta}{\sqrt{2}} \sqrt{E_1^2 + E_2^2}$$
$$= \frac{10^{-4}}{\sqrt{2}} \times 3 \times 10^8 \times 30 \times 10^{-6}$$
$$= \frac{90 \times 10^8 \times 10^{-10}}{\sqrt{2}}$$
$$\approx 0.6 \text{ N}$$

3. A stationary horizontal disc is free to rotate about its axis. When a torque is applied on it, its kinetic energy as a function of θ , where θ is the angle by which it has rotated, is given as $k\theta^2$. If its moment of inertia is I then the angular acceleration of the disc is:

(1)
$$\frac{2\mathbf{k}}{\mathbf{l}}\theta$$
 (2) $\frac{\mathbf{k}}{\mathbf{l}}\theta$
(3) $\frac{\mathbf{k}}{2\mathbf{l}}\theta$ (4) $\frac{\mathbf{k}}{4\mathbf{l}}\theta$

Answer (1)

Sol.
$$\tau = \frac{dE}{d\theta}$$

 $2k\theta = I\alpha$
 $\alpha = \frac{2k\theta}{I}$

- 4. The stream of a river is flowing with a speed of 2 km/h. A swimmer can swim at a speed of 4 km/h. What should be the direction of the swimmer with respect to the flow of the river to cross the river straight?
 - (1) 60° (2) 90°
 - (3) 150° (4) 120°

Answer (4)

Sol. Draw velocity diagram

$$v_{r}$$

$$v_{sr}$$

$$v_{sg}$$

$$\theta$$

$$sin \theta = \frac{v_{r}}{v_{sr}} = \frac{1}{2}$$

$$\theta = 30^{\circ}$$

$$\phi = 90 + \theta = 120^{\circ}$$



 $\left(\frac{1}{n}\right)^{th}$ part is hanging below the edge of the surface. To lift the hanging part of the cable upto the surface, the work done should be:

(1)
$$\frac{\text{MgL}}{n^2}$$
 (2) nMgL
(3) $\frac{\text{MgL}}{2n^2}$ (4) $\frac{2\text{MgL}}{n^2}$

Answer (3)

Sol.
$$m_1 = \frac{M}{n}$$

 $U_i = \frac{-M}{n} \times g \frac{L}{2n}$
 $W = \frac{MgL}{2n^2}$

6. A signal Acos ω t is transmitted using v₀sin ω_0 t as carrier wave. The correct amplitude modulated (AM) signal is:

(1)
$$\mathbf{v}_0 \sin \omega_0 \mathbf{t} + \frac{\mathbf{A}}{2} \sin(\omega_0 - \omega) \mathbf{t} + \frac{\mathbf{A}}{2} \sin(\omega_0 + \omega) \mathbf{t}$$

(2) $(v_0 + A) \cos \omega t \sin \omega_0 t$

- (3) $v_0 \sin \omega_0 t + A \cos \omega t$
- (4) $v_0 \sin[\omega_0(1 + 0.01 \text{ Asin}\omega t)t]$

Answer (1)

Sol. A = $(v_o + A\cos \omega t) \sin \omega_o t$

=
$$v_0 \sin(\omega_0 t) + \frac{A}{2} [\sin(\omega_0 - \omega)t + \sin(\omega_0 + \omega)t]$$

7. The following bodies are made to roll up (without slipping) the same inclined plane from a horizontal plane : (i) a ring of radius R, (ii) a

solid cylinder of radius $\frac{R}{2}$ and (iii) a solid

sphere of radius $\frac{R}{4}$. If, in each case, the speed

of the center of mass at the bottom of the incline is same, the ratio of the maximum heights they climb is :

(1) 14:15:20(2) 10:15:7(3) 4:3:2(4) 2:3:4

Answer (Bonus)

$$\begin{split} & \text{For ring} \to h_1 = \frac{1}{2} \frac{(2mr'^2)}{mg} \frac{v^2}{(r')^2} = \frac{v^2}{g} \\ & \text{For cylinder} \to h_2 = \frac{1}{2} \left(\frac{3}{2}mr'^2 \right) \frac{v^2}{(r')^2} = \frac{3}{4} \frac{v^2}{g} \\ & \text{For sphere} \to h_3 = \frac{1}{2} \times \frac{\frac{7}{5}m(r')^2}{g} \frac{v^2}{(r')^2} = \frac{7}{10} \frac{v^2}{g} \\ & h_1 : h_2 : h_3 \\ & 2 : \frac{3}{2} : \frac{14}{10} \\ & 20 : 15 : 14 \end{split}$$

8. A wire of resistance R is bent to form a square ABCD as shown in the figure. The effective resistance between E and C is

(E is mid-point of arm CD)

(1)
$$\frac{3}{4}R$$
 (2) R
(3) $\frac{1}{16}R$ (4) $\frac{7}{64}R$

Answer (4)

Sol.
$$R_1 = \frac{R}{8}, R_2 = \frac{7R}{8}$$

$$\frac{1}{R_{eq}} = \frac{8}{R} + \frac{8}{7R}$$

$$R_{eq} = \frac{7R}{64}$$

9. In the density measurement of a cube, the mass and edge length are measured as (10.00 \pm 0.10) kg and (0.10 \pm 0.01) m, respectively. The error in the measurement of density is

(1) 0.31 kg/m³ (2) 0.01 kg/m³

(3) 0.10 kg/m³ (4) 0.07 kg/m³

Answer (Bonus)



$$\frac{d\rho}{\rho} = \left[\frac{dM}{M} + \frac{dV}{V}\right]$$
$$\frac{d\rho}{10,000} = \frac{0.1}{10} + \frac{0.03}{0.1}$$
$$d\rho = 3100 \text{ kg/m}^3$$

/n 1)^o

10. A rectangular coil (Dimension 5 cm × 2.5 cm) with 100 turns, carrying a current of 3 A in the clock-wise direction, is kept centered at the origin and in the X-Z plane. A magnetic field of 1 T is applied along X-axis. If the coil is tilted through 45° about Z-axis, then the torque on the coil is

(1)	0.55 Nm	(2)	0.27	Nm
(3)	0.42 Nm	(4)	0.38	Nm

Answer (2)

Sol. $\tau = \vec{M} \times \vec{B}$

$$\tau = 100 \times 3 \times 5 \times 2.5 \times 10^{-4} \times 1 \times \frac{1}{\sqrt{2}}$$
$$= 0.27 \, \text{Nm}$$

11. A simple pendulum oscillating in air has period T. The bob of the pendulum is completely immersed in a non-viscous liquid. The density

of the liquid is $\frac{1}{16}$ th of the material of the bob.

If the bob is inside liquid all the time, its period of oscillation in this liquid is :

(1) $2T\sqrt{\frac{1}{14}}$	(2) $4T\sqrt{\frac{1}{15}}$
(3) $4T\sqrt{\frac{1}{14}}$	(4) $2T\sqrt{\frac{1}{10}}$

Answer (2)

Sol.
$$T = 2\pi \sqrt{\frac{I}{g_{eff}}}$$

 $\frac{T'}{T} = \sqrt{\frac{g_{eff}}{g'_{eff}}}$
 $g'_{eff} = g - \frac{g}{16} = \frac{15}{16}$
 $\frac{T'}{T} = \sqrt{\frac{16}{15}}$

L is varied by adjusting the separation between windings. The inductance of solenoid will be proportional to

(1)	1/L	(2)	L
(3)	1/L ²	(4)	L ²

Answer (1)

Sol. B = μ_0 ni

$$\phi = \mu_0 \operatorname{ni}(\mathbf{nL}) \mathbf{A}$$

(Inductance) L' =
$$\mu_0 n^2 L A$$
 $\left[\eta = \frac{N}{L} \right]$
L' = $\mu_0 NA\left(\frac{N}{L}\right)$
L' $\propto \frac{1}{L}$

- 13. For a given gas at 1 atm pressure, rms speed of the molecules is 200 m/s at 127°C. At 2 atm pressure and at 227°C, the rms speed of the molecules will be:
 - (1) $100\sqrt{5}$ m/s (2) 100 m/s
 - (3) $80\sqrt{5}$ m/s (4) 80 m/s

Answer (1)

Sol.
$$V \propto \sqrt{T}$$

 $\frac{V_1}{V_2} = \sqrt{\frac{T_1}{T_2}}$

$$\frac{200}{V_2} = \sqrt{\frac{400}{500}}$$

$$V_2 = 200\sqrt{\frac{5}{4}} = 100\sqrt{5}$$
 m/s

14. A capacitor with capacitance 5 μ F is charged to 5 μ C. If the plates are pulled apart to reduce the capacitance to 2 μ F, how much work is done?

(1) 2.55×10^{-6} J (2) 6.25×10^{-6} J (3) 3.75×10^{-6} J (4) 2.16×10^{-6} J

Answer (3)

2.10 ~ 10



2C₁

$$U_{f} = \frac{Q^{2}}{2C_{2}}$$
$$W = \frac{Q^{2}}{2} \left[\frac{1}{C_{2}} - \frac{1}{C_{1}} \right]$$
$$= \frac{(5)^{2}}{2} \left[\frac{1}{2} - \frac{1}{5} \right]$$
$$W = 3.75 \ \mu J$$

15. Following figure shows two processes A and B for a gas. If ΔQ_A and ΔQ_B are the amount of heat absorbed by the system in two cases, and ΔU_A and ΔU_B are changes in internal energies, respectively, then:



Answer (1)

Sol.
$$\Delta W_A > \Delta W_B$$

 $\Delta U_A > \Delta U_B$
From first law

$$\Delta \mathbf{Q} = \Delta \mathbf{U} + \Delta \mathbf{W}$$

$$\Delta \mathbf{Q}_{\mathsf{A}} > \Delta \mathbf{Q}_{\mathsf{B}}$$

16. A concave mirror for face viewing has focal length of 0.4 m. The distance at which you hold the mirror from your face in order to see your image upright with a magnification of 5 is:
(1) 0.32 m
(2) 0.24 m

of thermodynamics,

(3) 1.60 m (4) 0.16 m

Answer (1)

Sol.
$$\frac{1}{V} + \frac{1}{U} = -\frac{1}{40}$$

$$\frac{V}{U} = -5$$

$$50 \ 0 \ 40$$

U = - 32 cm

17. A solid sphere of mass 'M' and radius 'a' is surrounded by a uniform concentric spherical shell of thickness 2a and mass 2M. The gravitational field at distance '3a' from the centre will be:

(1)
$$\frac{GM}{9a^2}$$
 (2) $\frac{2GM}{9a^2}$
(3) $\frac{GM}{3a^2}$ (4) $\frac{2GM}{3a^2}$

Answer (3)

Sol.
$$E = \frac{GM}{(3a)^2} + \frac{2GM}{(3a)^2}$$
$$E = \frac{GM}{3a^2}$$

- 18. A moving coil galvanometer has resistance 50 Ω and it indicates full deflection at 4 mA current. A voltmeter is made using this galvanometer and a 5 k Ω resistance. The maximum voltage, that can be measured using this voltmeter, will be close to:
 - (1) 10 V (2) 20 V
 - (3) 15 V (4) 40 V

Answer (2)

Sol.
$$V = I_g (R_s + R_g)$$

= 4 × 10⁻³ [5050]
 $\approx 20 V$

19. The pressure wave, $P = 0.01 \sin[1000 t - 3x]Nm^{-2}$, corresponds to the sound produced by a vibrating blade on a day when atmospheric temperature is 0°C. On some other day when temperature is T, the speed of sound produced by the same blade and at the same frequency is found to be 336 ms⁻¹. Approximate value of T is :

(1) 4°C	(2) 12°C
---------	----------

(3) 15°C (4) 11°C

Answer (1)

Sol. $V_1 = \frac{1000}{3}$ m/s



$$\frac{dV}{V} = \frac{1}{2} \frac{dT}{T}$$
$$\frac{8 \times 3}{3 \times 1000} = \frac{1}{2} \times \frac{dT}{273}$$
$$dT = \frac{273 \times 2 \times 8}{1000} = 4.36^{\circ}C$$

20. Taking the wavelength of first Balmer line in hydrogen spectrum (n = 3 to n = 2) as 660 nm, the wavelength of the 2^{nd} Balmer line (n = 4 to n = 2) will be :

(1)	889.2 nm	(2) 488.9 nm
(3)	388.9 nm	(4) 642.7 nm

Answer (2)

Sol.
$$\frac{1}{\lambda_1} = R \left[\frac{1}{2^2} - \frac{1}{3^2} \right]$$
$$\frac{1}{\lambda_2} = R \left[\frac{1}{2^2} - \frac{1}{4^2} \right]$$
$$\frac{5\lambda_1}{36} = \frac{12\lambda_2}{4 \times 16}$$
$$\lambda_2 = \frac{5 \times 660 \times 64}{36 \times 12}$$

= 489 nm

21. A body of mass 2 kg makes an elastic collision with a second body at rest and continues to move in the original direction but with one fourth of its original speed. What is the mass of the second body?

(1)	1.5 kg	(2) 1.8 kg	J
(3)	1.0 kg	(4) 1.2 kg	J

Answer (4)

Sol.
$$2V = \frac{2V}{4} + m_2V_2$$

 $V = V_2 - \frac{V}{4}$
 $\frac{3V}{2} = m_2\left(V + \frac{V}{4}\right)$
 $m_2 = \frac{6}{5}$ kg

HCI molecules in its gaseous phase is \overline{v} , m is its mass and k_B is Boltzmann constant, then its temperature will be :

(1)
$$\frac{m\overline{v}^{2}}{5k_{B}}$$
(2)
$$\frac{m\overline{v}^{2}}{6k_{B}}$$
(3)
$$\frac{m\overline{v}^{2}}{7k_{B}}$$
(4)
$$\frac{m\overline{v}^{2}}{3k_{B}}$$
Answer (4)
Sol.
$$\overline{v} = \sqrt{\frac{3 \text{ RT}}{M}}$$

$$\overline{v} = \sqrt{\frac{3 \text{ RT}}{mN_{A}}}$$

$$\overline{v} = \sqrt{\frac{3 k_{B}T}{m}}$$

23. A system of three charges are placed as shown in the figure :



If D >> d, the potential energy of the system is best given by :

(1)
$$\frac{1}{4\pi\epsilon_{0}} \left[-\frac{q^{2}}{d} - \frac{qQd}{D^{2}} \right]$$

(2)
$$\frac{1}{4\pi\epsilon_{0}} \left[+\frac{q^{2}}{d} + \frac{qQd}{D^{2}} \right]$$

(3)
$$\frac{1}{4\pi\epsilon_{0}} \left[-\frac{q^{2}}{d} + \frac{2qQd}{D^{2}} \right]$$

(4)
$$\frac{1}{4\pi\epsilon_{0}} \left[-\frac{q^{2}}{d} - \frac{qQd}{2D^{2}} \right]$$



Sol.
$$U = -\frac{Kq^2}{d} + QV$$

$$\mathbf{V} = -\frac{\mathbf{KP}}{\mathbf{D}^2} = -\frac{\mathbf{Kqd}}{\mathbf{D}^2}$$

$$\mathbf{U} = -\frac{\mathbf{Kq}^2}{\mathbf{d}} - \frac{\mathbf{KQqd}}{\mathbf{D}^2}$$

24. A rigid square loop of side 'a' and carrying current I_2 is lying on a horizontal surface near a long current I_1 carrying wire in the same plane as shown in figure. The net force on the loop due to the wire will be :



- (1) Repulsive and equal to $\frac{\mu_0 I_1 I_2}{4\pi}$
- (2) Repulsive and equal to $\frac{\mu_0 I_1 I_2}{2\pi}$
- (3) Zero

(4) Attractive and equal to
$$\frac{\mu_0 I_1 I_2}{3\pi}$$

Answer (1)

Sol. $F = I_2 a (B_1 - B_2)$

$$\mathbf{B_1} = \frac{\mu_0 \mathbf{I_1}}{2\pi \mathbf{a}}$$

$$\mathbf{B_2} = \frac{\mu_0 \mathbf{I_1}}{4\pi \mathbf{a}}$$

$$\mathbf{F} = \frac{\mu_0 \mathbf{I_1} \mathbf{I_2}}{4\pi}$$

the stationary wave is Y = 0.3 sin (0.157x) cos ($200\pi t$). The length of the string is : (All quantities are in SI units)

- (1) 60 m
- (2) 20 m
- (3) 40 m
- (4) 80 m

Answer (4)

Sol. λ = 0.157

$$=\frac{3.14}{20}=\frac{\pi}{20}$$

In 4th harmonic

$$\therefore \frac{2\pi}{\lambda} = \pi/20$$

λ **= 40 m**

$$\therefore \quad \mathbf{L} = \frac{4\lambda}{2} = 80 \text{ m}$$

- 26. An NPN transistor is used in common emitter configuration as an amplifier with 1 k Ω load resistance. Signal voltage of 10 mV is applied across the base-emitter. This produces a 3 mA change in the collector current and 15 μ A change in the base current of the amplifier. The input resistance and voltage gain are :
 - (1) 0.33 kΩ, 1.5
 - (2) 0.33 kΩ, 300
 - (3) 0.67 kΩ, 200
 - (4) 0.67 kΩ, 300
- Answer (4)



---- I_B

$$R_{i} = \frac{10 \times 10^{-3}}{15 \times 10^{-6}} = 0.67 \, \text{K}\Omega$$

Voltage Gain
$$= \beta \left(\frac{R_o}{R_i} \right) = 300$$

27. The electric field of light wave is given as

$$\vec{E} = 10^{-3} \cos\left(\frac{2\pi x}{5 \times 10^{-7}} - 2\pi \times 6 \times 10^{14} t\right) \hat{x} \frac{N}{C}$$
. This

light falls on a metal plate of work function 2eV. The stopping potential of the photoelectrons is :

Given, E (in eV) =
$$\frac{12375}{\lambda (in \text{ Å})}$$

(1) 0.48 V
(2) 2.48 V
(3) 0.72 V
(4) 2.0 V
Answer (1)
Sol. $\lambda = 5 \times 10^{-7} \text{ m}$
 $v = \frac{3 \times 10^8}{5 \times 10^{-7}} = 6 \times 10^{14} \text{ Hz}$
E = $\frac{12375}{5 \times 10^{-7}} = 2.475 \text{ eV}$

$$E = \frac{125.2}{5000} = 2.47$$

 $K_{max} = E - \phi$
= 0.48 eV

A ball is thrown vertically up (taken as + z-axis) from the ground. The correct momentum-height (p-h) diagram is:



Sol. $V = \sqrt{V_0^2 - 2gh}$

Direction of velocity changes at top most point

- 29. If 'M' is the mass of water that rises in a capillary tube of radius 'r', then mass of water which will rise in a capillary tube of radius '2r' is
 - (1) 2 M
 - (2) M
 - (3) 4 M

$$(4) \frac{M}{2}$$

Answer (1)

Sol. $h \propto 1/r$

 $\mathbf{M} \propto \pi \mathbf{r^2} \mathbf{h}$

 $\textbf{M} \propto \textbf{r}$

30. The figure shows a Young's double slit experimental setup. It is observed that when a thin transparent sheet of thickness t and refractive index μ is put in front of one of the slits, the central maximum gets shifted by a distance equal to n fringe widths. If the wavelength of light used is λ , t will be



(3)
$$\frac{nD\lambda}{a(\mu-1)}$$

(4)
$$\frac{D\lambda}{a(\mu-1)}$$

Answer (Bonus)

(1)

(2)

Sol. (μ –1) t = n λ

$$t = \frac{n\lambda}{\mu - 1}$$



- 1. C₆₀, an allotrope of carbon contains
 - (1) 16 hexagons and 16 pentagons
 - (2) 18 hexagons and 14 pentagons
 - (3) 20 hexagons and 12 pentagons
 - (4) 12 hexagons and 20 pentagons

Answer (3)

- **Sol.** Fullerene (C_{60}) contains 20 six membered rings and 12 five membered rings.
- 2. The major product of the following reaction is

 $CH_{3}C \equiv CH \xrightarrow{(i) DCI (1 equiv.)}_{(ii) DI} \rightarrow$

- (1) $CH_3C(I)(CI)CHD_2$ (2) $CH_3CD(I)CHD(CI)$
- (3) CH₃CD(CI)CHD(I) (4) CH₃CD₂CH(CI)(I)

Answer (1)



Both addition follow Markownikov's rule.

3. The standard Gibbs energy for the given cell reaction in kJ mol⁻¹ at 298 K is

Zn(s) + $Cu^{2+}(aq) \longrightarrow Zn^{2+}(aq)$ + Cu(s),

E° = 2 V at 298 K

(Faraday's constant, F = 96000 C mol⁻¹)

- (1) 192
- (2) 384
- (3) -384
- (4) -192

Answer (3)

Sol. $\Delta G^\circ = - nFE^\circ_{cell}$

= -384 kJ/mole

- 4. The number of water molecule(s) not coordinated to copper ion directly in $CuSO_4 \cdot 5H_2O$, is
 - (1) 4
 (2) 1

 (3) 2
 (4) 3

Answer (2)

- Sol. In $CuSO_4 \cdot 5H_2O$, four H_2O molecules are directly coordinated to the central metal ion while one H_2O molecule is hydrogen bonded.
- 5. Magnesium powder burns in air to give
 - (1) $Mg(NO_3)_2$ and Mg_3N_2
 - (2) MgO and $Mg(NO_3)_2$
 - (3) MgO and Mg_3N_2
 - (4) MgO only

Answer (3)

Sol. Mg burn in air and produces a mixture of nitride and oxide.

So, Mg_3N_2 and MgO are formed.

6. The one that will show optical activity is (en = ethane-1,2-diamine)



collegebatch.com



or centre of symmetry, hence it is optically active.

- Liquid 'M' and liquid 'N' form an ideal solution. The vapour pressures of pure liquids 'M' and 'N' are 450 and 700 mmHg, respectively, at the same temperature. Then correct statement is
 - $(x_{M} = Mole fraction of 'M' in solution;$
 - x_N = Mole fraction of 'N' in solution;
 - y_{M} = Mole fraction of 'M' in vapour phase;

 y_N = Mole fraction of 'N' in vapour phase)

(1)
$$\frac{x_{M}}{x_{N}} = \frac{y_{M}}{y_{N}}$$
 (2) $\frac{x_{M}}{x_{N}} > \frac{y_{M}}{y_{N}}$
(3) $\frac{x_{M}}{x_{N}} < \frac{y_{M}}{y_{N}}$ (4) $(x_{M} - y_{M}) < (x_{N} - y_{N})$

Answer (2)

Sol. $P_M^o = 450 \text{ mmHg}, P_N^o = 700 \text{ mmHg}$

$$\begin{split} \mathbf{P}_{\mathbf{M}} &= \mathbf{P}_{\mathbf{M}}^{\mathbf{o}} \mathbf{X}_{\mathbf{M}} = \mathbf{Y}_{\mathbf{M}} \mathbf{P}_{\mathbf{T}} \\ & \Rightarrow \quad \mathbf{P}_{\mathbf{M}}^{\mathbf{o}} = \frac{\mathbf{Y}_{\mathbf{M}}}{\mathbf{X}_{\mathbf{M}}} (\mathbf{P}_{\mathbf{T}}) \\ & \text{Similarly, } \mathbf{P}_{\mathbf{N}}^{\mathbf{o}} = \frac{\mathbf{Y}_{\mathbf{N}}}{\mathbf{X}_{\mathbf{N}}} (\mathbf{P}_{\mathbf{T}}) \\ & \text{Given, } \mathbf{P}_{\mathbf{M}}^{\mathbf{o}} < \mathbf{P}_{\mathbf{N}}^{\mathbf{o}} \\ & \Rightarrow \quad \frac{\mathbf{Y}_{\mathbf{M}}}{\mathbf{X}_{\mathbf{M}}} < \frac{\mathbf{Y}_{\mathbf{N}}}{\mathbf{X}_{\mathbf{N}}} \end{split}$$

$$\Rightarrow \frac{\mathbf{I}_{\mathsf{M}}}{\mathsf{Y}_{\mathsf{N}}} < \frac{\mathbf{\Lambda}_{\mathsf{M}}}{\mathsf{X}_{\mathsf{N}}}$$

_

8. The major product of the following reaction is

$$CH_{3}CH = CHCO_{2}CH_{3} \xrightarrow{\text{LiAlH}_{4}}$$
(1)
$$CH_{3}CH_{2}CH_{2}CH_{2}OH$$
(2)
$$CH_{3}CH_{2}CH_{2}CHO$$
(3)
$$CH_{3}CH_{2}CH_{2}CO_{2}CH_{3}$$
(4)
$$CH_{3}CH = CHCH_{2}OH$$

Answer (4)

$$CH_3 - CH = CH - CO_2Me$$

LIAIH
 $CH_3 - CH = CH - CH_2$

9. For any given series of spectral lines of atomic hydrogen, let $\Delta \overline{v} = \overline{v}_{max} - \overline{v}_{min}$ be the difference in maximum and minimum frequencies in cm⁻¹. The ratio $\Delta \overline{v}_{Lyman} / \Delta \overline{v}_{Balmer}$ is

(1)	9	:	4	(2)	27	:	5
1)	9	:	4	(2)	27	:	5

Answer (1)

Sol.
$$\overline{\mathbf{v}} \propto \Delta \mathbf{E}$$

For H-atom

$$\overline{v} = \mathbf{R} \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$$

For Lyman series,

$$\overline{v}$$
 (max) $\propto 13.6 \left(1 - \frac{1}{\infty} \right)$
 \overline{v} (min) $\propto 13.6 \left(1 - \frac{1}{4} \right)$

$$\therefore \quad \overline{v}_{max} - \overline{v}_{min} \propto 13.6 \left(\frac{1}{4}\right)$$

For Balmer series,

$$\overline{v} (\max) \propto 13.6 \left(\frac{1}{4} - \frac{1}{\infty}\right)$$
$$\overline{v} (\min) \propto 13.6 \left(\frac{1}{4} - \frac{1}{9}\right)$$
$$\therefore \quad \overline{v}_{\min} - \overline{v}_{\min} \propto 13.6 \left(\frac{1}{9}\right)$$
$$\frac{v_{\text{Lyman}}}{v_{\text{Balmer}}} = \frac{9}{4}$$

10. Among the following, the set of parameters that represents path functions, is



- (C) w, path function
- (D) H TS = G, state function
- 11. The major product of the following reaction is



Answer (2)

Sol. Alkaline $KMnO_4$ converts \bigcirc -R with a

benzylic hydrogen into benzoic acid.



12. The increasing order of reactivity of the following compounds towards aromatic electrophilic substitution reaction is





13. The given plots represent the variation of the concentration of a reactant R with time for two different reactions (i) and (ii). The respective orders of the reactions are



Answer (3)

Sol. Graph-(i) : In[Reactant] vs time is linear

Hence, 1st order

Graph-(ii) : [Reactant] vs time is linear

Hence, zero order

14. The osmotic pressure of a dilute solution of an ionic compound XY in water is four times that of a solution of 0.01 M $BaCl_2$ in water. Assuming complete dissociation of the given ionic compounds in water, the concentration of XY (in mol L⁻¹) in solution is

(1)
$$16 \times 10^{-4}$$
 (2) 4×10^{-4}

(3) 6×10^{-2} (4) 4×10^{-2}

Answer (3)

Sol. $\pi_{XY} = 4\pi_{BaCl_2}$

$$= 6 \times 10^{-2} \frac{\text{mol}}{\text{L}}$$

15. Consider the van der Waals constants, a and b, for the following gases.

Gas	Ar	Ne	Kr	Xe
a/(atm dm ⁶ mol ⁻²)	1.3	0.2	5.1	4.1
b/(10 ⁻² dm ³ mol ⁻¹)	3.2	1.7	1.0	5.0
Which gas is expected to have the highest critical temperature?				
(1) Ne (2) Kr				
(3) Xe	(4)	Ar		

Answer (2)



So, species with greatest value of $\frac{a}{b}$ has

greatest value of critical temperature i.e. Kr.

16. The major product of the following reaction is



Answer (2)



More stable product due to conjugation.

17. Among the following, the molecule expected to be stabilized by anion formation is

 $\begin{array}{c} {\sf C}_2,\,{\sf O}_2,\,{\sf NO},\,{\sf F}_2\\ (1)\,\,{\sf F}_2 & (2)\,\,{\sf NO}\\ (3)\,\,{\sf C}_2 & (4)\,\,{\sf O}_2 \end{array}$

Answer (3)

Sol. C₂ has s-p mixing and the HOMO is $\pi 2p_x = \pi 2p_y$ and LUMO is $\sigma 2p_z$. So, the extra electron will occupy bonding molecular orbital and this will lead to an increase in bond order.

 C_2^- has more bond order than C_2 .

- 18. Excessive release of CO_2 into the atmosphere results in
 - (1) Depletion of ozone
 - (2) Polar vortex
 - (3) Formation of smog
 - (4) Global warming

Answer (4)

Sol. CO_2 causes global warming.

 $i_2(g) + 3i_2(g) \rightarrow 2i_3(g),$

Identify dihydrogen (H_2) as a limiting reagent in the following reaction mixtures.

- (1) 35 g of N_2 + 8 g of H_2
- (2) 28 g of N_2 + 6 g of H_2
- (3) 56 g of N_2 + 10 g of H_2
- (4) 14 g of N_2 + 4 g of H_2

Answer (3)

Sol. 28 g N_2 react with 6 g H_2

$$\begin{array}{c} \mathsf{N}_2 + \mathsf{3H}_2 \\ \overset{1 \text{ mole }}{\underset{6 \text{ g}}{3 \text{ moles }}} \xrightarrow{3 \text{ moles }} \mathsf{2NH} \end{array}$$

For 56 g of N_2 , 12 g of H_2 is required.

20. Match the catalysts (Column I) with products (Column II).

	Column I		Column II	
	Catalyst		Product	
(A)	V ₂ O ₅	(i)	Polyethylene	
(B)	TiCl ₄ /Al(Me) ₃	(ii)	Ethanal	
(C)	PdCl ₂	(iii)	H_2SO_4	
(D)	Iron Oxide	(iv)	NH ₃	
(1)	(A)-(iv); (B)-(iii); (C)	-(ii);	(D)-(i)	
(2)	(A)-(iii); (B)-(iv); (C)-(i); (D)-(ii)			
(3)	(A)-(iii); (B)-(i); (C)-((ii); ((D)-(iv)	
(4)	(A)-(ii); (B)-(iii); (C)-	(i); ((D)-(iv)	

Answer (3)

Sol.	(A) $V_2O_5 \rightarrow$	Preparation of $\rm H_2SO_4$ in contacts process
	(B) TiCl ₄ + Al(Me) ₃ \rightarrow	Polyethylene (Ziegler- Natta catalyst)
	(C) $PdCl_2 \rightarrow$	Ethanal (Wacker's process)
	(D) Iron oxide \rightarrow	NH ₃ in Haber's process
21.	The element havi between its first energies, is	ng greatest difference and second ionization
	(1) K	(2) Sc
	(3) Ca	(4) Ba

Answer (1)

Sol. Alkali metals have high difference in the first ionisation and the second ionisation energy as they achieve stable noble gas configuration after first ionisation.



(1) $NO_2 < NO < N_2O_3 < N_2O$ (2) $N_2O < NO < N_2O_3 < NO_2$ (3) $NO_2 < N_2O_3 < NO < N_2O$ (4) $N_2O < N_2O_3 < NO < NO_2$

Answer (2)

Sol. (oxide) (oxidation state) N₂O + 1 NO + 2

-
+ 3
+ 4

So, $N_2O < NO < N_2O_3 < NO_2$

- 23. Which of the following statements is not true about sucrose?
 - (1) The glycosidic linkage is present between C₁ of α -glucose and C₁ of β -fructose
 - (2) On hydrolysis, it produces glucose and fructose
 - (3) It is a non-reducing sugar
 - (4) It is also named as invert sugar

Answer (1)

Sol. Sucrose contains glycosidic link between C₁ of α -D glucose and C₂ of β -D-Fructose.

 $C_{12}H_{22}O_{11}+H_2O \longrightarrow Glu cose+Fructose$

- 24. The ore that contains the metal in the form of fluoride is
 - (1) malachite (2) sphalerite
 - (3) magnetite (4) cryolite

Answer (4)

- Sol. Magnetite Fe_3O_4 SphaleriteZnSCryolite Na_3AIF_6 Malachite $CuCO_3 \cdot Cu(OH)_2$
- 25. The correct IUPAC name of the following compound is



- (3) 2-methyl-5-nitro-1-chlorobenzene
- (4) 2-chloro-1-methyl-4-nitrobenzene

Answer (4)



All Groups attached are to be treated as substituents and lowest set of locant rule is followed.

2-Chloro-1-methyl-4-nitrobenzene

26. Aniline dissolved in dilute HCI is reacted with sodium nitrite at 0°C. This solution was added dropwise to a solution containing equimolar mixture of aniline and phenol in dil. HCI. The structure of the major product is



Answer (2)

Sol. In acidic medium aniline is more reactive than phenol that's why electrophilic aromatic

substitution of $Ph_{N_{a}}$ takes place with aniline



- 27. The aerosol is a kind of colloid in which
 - (1) solid is dispersed in gas
 - (2) gas is dispersed in solid
 - (3) liquid is dispersed in water
 - (4) gas is dispersed in liquid

Answer (1)



- 28. The degenerate orbitals of $[Cr(H_2O)_6]^{3+}$ are
 - (1) d_{xz} and d_{yz}
 - (2) $d_{x^2-y^2}$ and d_{xy}
 - (3) d_{z^2} and d_{xz}
 - (4) d_{yz} and d_{z^2}

Answer (1)

Sol. Cr³⁺ has d³ configuration and forms an octahedral inner orbitals complex.

The set of degenerate orintals are (d_{xy}, d_{yz} and d_{xz}) and (d_{x²-y²} and d_{z²}).

29. The major product of the following reaction is





30. The organic compound that gives following qualitative analysis is





- 1. Let p, q \in R. If $2-\sqrt{3}$ is a root of the quadratic equation, x^2 + px + q = 0, then :
 - (1) $q^2 4p 16 = 0$ (2) $p^2 4q + 12 = 0$ (3) $p^2 - 4q - 12 = 0$ (4) $q^2 + 4p + 14 = 0$

Answer (3)

Sol. p, q are rational numbers

 $\therefore 2+\sqrt{3} \text{ in the other root}$ Now, p = -4, q = 1 $\Rightarrow p^2 - 4q - 12 = 16 - 4 - 12$ = 0

Note:- (Erratum) p, q, should be given as rational numbers instead of real numbers

2. If the line, $\frac{x-1}{2} = \frac{y+1}{3} = \frac{z-2}{4}$ meets the plane, x + 2y + 3z = 15 at a point P, then the distance

of P from the origin is :



Answer (2)

Sol. Let point on line is p(2r + 1, 3r – 1, 4r + 2)

It lies on the plane x + 2y + 3z = 15

$$\therefore 2r + 1 + 6r - 2 + 12r + 6 = 15$$

$$\Rightarrow r = \frac{1}{2}$$

$$\therefore P = \left(2, \frac{1}{2}, 4\right)$$

$$\therefore OP = \sqrt{4 + \frac{1}{4} + 16} = \sqrt{\frac{81}{4}} = \frac{9}{2}$$

3. If a tangent to the circle $x^2 + y^2 = 1$ intersects the coordinate axes at distinct points P and Q, then the locus of the mid-point of PQ is :

(1)
$$x^{2} + y^{2} - 16x^{2}y^{2} = 0$$

(2) $x^{2} + y^{2} - 2x^{2}y^{2} = 0$
(3) $x^{2} + y^{2} - 4x^{2}y^{2} = 0$
(4) $x^{2} + y^{2} - 2xy = 0$

Answer (3)

Sol. Let any tangent to circle $x^2 + y^2 = 1$ is x cos $\theta + y \sin \theta = 1$

$$x \cos\theta + y \sin\theta = 1$$

$$P\left(\frac{1}{\cos\theta},0\right); \quad Q\left(0,\frac{1}{\sin\theta}\right)$$

 $\therefore \quad \text{Mid-point of PQ let } M\left(\frac{1}{2\cos\theta}, \frac{1}{2\sin\theta}\right) = (h, k)$

$$\Rightarrow \cos\theta = \frac{1}{2h}; \quad \sin\theta = \frac{1}{2k}$$

... On squaring and adding

$$\frac{1}{h^2} + \frac{1}{k^2} = 4 \qquad \Rightarrow x^2 + y^2 = 4x^2y^2$$

4. If f(x) is a non-zero polynomial of degree four, having local extreme points at x = -1, 0, 1; then the set

$$S = \{x \in R : f(x) = f(0)\}$$

contains exactly :

- (1) Four irrational numbers
- (2) Four rational numbers
- (3) Two irrational and one rational number
- (4) Two irrational and two rational numbers

Answer (3)

Sol.
$$f'(x) = A(x + 1) \times (x - 1)$$

$$= A(x^3 - x)$$

$$\Rightarrow f(\mathbf{x}) = \mathbf{A}\left(\frac{\mathbf{x}^4}{4} - \frac{\mathbf{x}^2}{2}\right) + \mathbf{C}$$

Now f(0) = C

$$\therefore f(\mathbf{x}) = f(\mathbf{0}) \qquad \Rightarrow \quad A\left(\frac{\mathbf{x}^4}{4} - \frac{\mathbf{x}^2}{2}\right) = \mathbf{0}$$

$$\Rightarrow \quad \frac{\mathbf{x}^2}{2}\left(\frac{\mathbf{x}^2}{2} - 1\right) = \mathbf{0}$$

$$\Rightarrow \quad \mathbf{x} = \mathbf{0}, \mathbf{0}, -\sqrt{2}, \sqrt{2}$$

$$\therefore \quad \mathbf{S} = \left\{\mathbf{0}, -\sqrt{2}, \sqrt{2}\right\}$$



 $1, K = \sqrt{3}$ where $K \neq 0$, then K is equal to .

(1)
$$\sqrt{6}$$
 (2) $2\sqrt{6}$
(3) $2\sqrt{\frac{10}{3}}$ (4) $4\sqrt{\frac{5}{3}}$

Answer (2)

Sol. Mean of given observation = $\frac{k}{4}$

$$\therefore \sigma^2 = 5 \text{(given)} \qquad \dots(i)$$
Also $\sigma^2 = \frac{\left(\frac{k}{4} + 1\right)^2 + \left(\frac{k}{4}\right)^2 + \left(\frac{k}{4} - 1\right)^2 + \left(\frac{3k}{4}\right)^2}{4} \qquad \dots(ii)$

... from (i) and (ii)

$$\frac{\frac{12k^2}{16} + 2}{4} = 5 \quad \Rightarrow \frac{12k^2}{16} = 18$$

$$\Rightarrow$$
 k² = 24 \Rightarrow k = 2 $\sqrt{6}$

 $\left(\frac{2}{x} + x^{\log_{8} x}\right)^{6} (x > 0) \text{ is } 20 \times 8^{7}, \text{ then a value of } x$ is : (1) 8^{3} (2) 8 (3) 8^{-2} (4) 8^{2}

Answer (4)

Sol.
$$T_4 = 20 \times 8^7 = {}^6C_3 \left(\frac{2}{x}\right)^3 \times \left(x^{\log_8 x}\right)^3$$

$$\Rightarrow 8 \times 20 \times \left(\frac{x^{\log_8 x}}{x}\right)^3 = 20 \times 8^7$$

$$\Rightarrow \left(\frac{x^{\log_8 x}}{x}\right)^2 = \left(8^2\right)^3$$

$$\Rightarrow \frac{x^{\log_8 x}}{x} = 64$$
 Take \log_8 both side

$$\Rightarrow (\log_8 x)^2 - (\log_8 x) = 2$$

$$\Rightarrow \log_8 x = -1 \quad \text{or} \quad \log_8 x = 2$$

$$\Rightarrow x = \frac{1}{8} \qquad \text{or} \qquad x = 8^2$$

(Here C is a constant of integration)

(1)
$$-3\cot^{-1/3}x + C$$

(2) $-3\tan^{-1/3}x + C$
(3) $-\frac{3}{4}\tan^{-4/3}x + C$
(4) $3\tan^{-1/3}x + C$

Answer (2)

Sol.
$$I = \int \sec^{\frac{2}{3}} x \cdot \csc^{\frac{4}{3}} dx$$

$$I = \int \frac{\sec^2 x \, dx}{\tan^3 x}$$
 Put tan x = t

$$\Rightarrow$$
 sec²x dx = dt

$$\Rightarrow I = \frac{t^{\frac{-1}{3}}}{\left(\frac{-1}{3}\right)} + C \quad \Rightarrow I = -3(tanx)^{\frac{-1}{3}} + C$$

- 8. If one end of a focal chord of the parabola, $y^2 = 16x$ is at (1, 4), then the length of this focal chord is :
 - (1) 24
 - (2) 20
 - (3) 22
 - (4) 25

Answer (4)

Sol.
$$\therefore$$
 y² = 16x \Rightarrow a = 4
One end of focal chord (1, 4) \therefore 2 at = 4

$$\Rightarrow t = \frac{1}{2}$$

Length of focal chord = $a\left(t + \frac{1}{t}\right)^2$
= $4 \times \left(2 + \frac{1}{2}\right)^2$
= 25

collegebatch.com

$$f(\mathbf{x}) = \begin{cases} \frac{\sqrt{2}\cos x - 1}{\cot x - 1}, & \mathbf{x} \neq \frac{\pi}{4} \\ \mathbf{k}, & \mathbf{x} = \frac{\pi}{4} \end{cases}$$

is continuous, then k is equal to :

(1) 1 (2)
$$\frac{1}{2}$$

(3) $\frac{1}{\sqrt{2}}$ (4) 2

Answer (2)

Sol.
$$\lim_{x \to \frac{\pi}{4}} \frac{\sqrt{2} \cos x - 1}{\cot x - 1} = k \qquad \therefore \text{ By L hospital rule.}$$

$$\Rightarrow \lim_{\mathbf{x} \to \frac{\pi}{4}} \frac{\sqrt{2} \sin \mathbf{x}}{\operatorname{cosec}^2 \mathbf{x}} = \mathbf{k} \qquad \Rightarrow \mathbf{k} = \frac{1}{2}$$

10. All the points in the set

$$\mathbf{S} = \left\{ \frac{\alpha + \mathbf{i}}{\alpha - \mathbf{i}} : \alpha \in \mathbf{R} \right\} (\mathbf{i} = \sqrt{-1})$$

lie on a :

- (1) Straight line whose slope is 1
- (2) Circle whose radius is $\sqrt{2}$
- (3) Circle whose radius is 1
- (4) Straight line whose slope is -1

Answer (3)

Sol.
$$\therefore$$
 $S = \frac{\alpha + i}{\alpha - i}$ Let $S = x + iy$
 $\Rightarrow x + iy = \frac{(\alpha + i)^2}{\alpha^2 + 1}$ (by rationalisation)
 $\Rightarrow x + iy = \frac{(\alpha^2 - 1)}{\alpha^2 + 1} + \frac{i(2\alpha)}{\alpha^2 + 1}$ (On comparing both sides)
 $\Rightarrow x = \frac{\alpha^2 - 1}{\alpha^2 + 1}$... (i) $y = \frac{2\alpha}{\alpha^2 + 1}$... (ii)
By squaring and adding

 \Rightarrow x² + y² = 1

where $\vec{\beta}_1$ is parallel to $\vec{\alpha}$ and $\vec{\beta}_2$ is perpendicular to $\vec{\alpha}$, then $\vec{\beta}_1 \times \vec{\beta}_2$ is equal to :

(1)
$$\frac{1}{2}(3\hat{i}-9\hat{j}+5\hat{k})$$
 (2) $\frac{1}{2}(-3\hat{i}+9\hat{j}+5\hat{k})$
(3) $-3\hat{i}+9\hat{j}+5\hat{k}$ (4) $3\hat{i}-9\hat{j}-5\hat{k}$

Answer (2)

Sol.
$$\vec{\beta} = \vec{\beta_1} - \vec{\beta_2}$$
 ...(i)
 $\therefore \quad \vec{\beta_2} \cdot \vec{\alpha} = 0$
and Let $\vec{\beta_1} = \lambda \vec{\alpha}$
 $\vec{\alpha} \cdot \vec{\beta} = \vec{\alpha} \cdot \vec{\beta_1} - \vec{\alpha} \cdot \vec{\beta_2}$
 $\Rightarrow \quad 5 = \lambda \alpha^2$
 $\Rightarrow \quad 5 = \lambda \times 10$
 $\Rightarrow \quad \lambda = \frac{1}{2}$
 $\therefore \quad \boxed{\vec{\beta_1} = \frac{\vec{\alpha}}{2}}$

Cross product with $\overline{\beta_1}$ in equation (i)

$$\Rightarrow \vec{\beta} \times \vec{\beta_1} = -\vec{\beta_2} \times \vec{\beta_1}$$

$$\Rightarrow \overline{\vec{\beta} \times \vec{\beta_1}} = \vec{\beta_1} \times \vec{\beta_2} = \frac{\vec{\beta_1} \times \vec{\beta_2}}{2}$$

$$\Rightarrow \vec{\beta_1} \times \vec{\beta_2} = \frac{1}{2} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -1 & 3 \\ 3 & 1 & 0 \end{vmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} -3\hat{i} - \hat{j}(-9) + \hat{k}(5) \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} -3\hat{i} + 9\hat{j} + 5\hat{k} \end{bmatrix}$$

- 12. Let $\sum_{k=1}^{10} f(a+k) = 16(2^{10} 1)$, where the function f satisfies f(x + y) = f(x) f(y) for all natural numbers x, y and f(1) = 2. Then the natural number 'a' is :
 - (1) 2(2) 3
 - (2) 0
 - (3) 16
 - (4) 4
- Answer (2)



$$\therefore \text{ Let } \left[f(x) = b^{x} \right]$$

$$\therefore f(1) = 2$$

$$\therefore b' = 2$$

$$\Rightarrow \overline{f(x) = 2^{x}}$$

Now, $\sum_{k=1}^{10} 2^{a+k} = 16(2^{10} - 1)$

$$\Rightarrow 2^{a} \sum_{k=1}^{10} 2^{k} = 16(2^{10} - 1)$$

$$\Rightarrow 2^{a} \times \frac{((2^{10}) - 1) \times 2}{(2 - 1)} = 16 \times (2^{10} - 1)$$

$$\Rightarrow 2^{a} = 8$$

$$\Rightarrow \boxed{a = 3}$$

13. If $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \cdot \dots \cdot \begin{bmatrix} 1 & n - 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 78 \\ 0 & 1 \end{bmatrix},$
then the inverse of $\begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}$ is :
(1) $\begin{bmatrix} 1 & 0 \\ 13 & 1 \end{bmatrix}$ (2) $\begin{bmatrix} 1 & -13 \\ 0 & 1 \end{bmatrix}$
(3) $\begin{bmatrix} 1 & -12 \\ 0 & 1 \end{bmatrix}$ (4) $\begin{bmatrix} 1 & 0 \\ 12 & 1 \end{bmatrix}$
Answer (2)
Sol. $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 4 \\ 0 & 1 \end{bmatrix} \dots \begin{bmatrix} 1 & n - 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 78 \\ 0 & 1 \end{bmatrix}$

$$\begin{bmatrix} 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 & 1+2+3+\ldots+(n-1) \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 78 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow \frac{(n-1)n}{2} = 78$$

$$\Rightarrow n = 13$$
Now, inverse of
$$\begin{bmatrix} 1 & 13 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -13 \\ 0 & 1 \end{bmatrix}$$
14. Let S = { $\theta \in [-2\pi, 2\pi]$; $2\cos^2\theta + 3\sin\theta = 0$ }. Then the sum of the elements of S is :
(1) π (2) 2π

(3)
$$\frac{13\pi}{6}$$
 (4) $\frac{5\pi}{3}$

Answer (2)

- \Rightarrow 2311 0 03110 2 0
- \Rightarrow (2sin θ + 1)(sin θ –2) = 0

$$\Rightarrow \sin\theta = -\frac{1}{2}; \quad \sin\theta = 2 \rightarrow \text{Not Possible}$$

$$\frac{\pi}{6}$$
 $-\frac{\pi}{6}$

Т

 \therefore Sum of all solutions in [-2 π , 2 π] is

$$= \left(\pi + \frac{\pi}{6}\right) + \left(2\pi - \frac{\pi}{6}\right) + \left(-\frac{\pi}{6}\right) + \left(-\pi + \frac{\pi}{6}\right) = 2\pi$$

- 15. Let f(x) = 15 |x 10|; $x \in R$. Then the set of all values of x, at which the function, g(x) = f(f(x)) is not differentiable, is :
 - (1) (10, 15)
 - (2) {5, 10, 15, 20}
 - (3) {10}
 - (4) {5, 10, 15}

Answer (4)

 \Rightarrow

Sol. Given f(x) = 15 - |(10 - x)|

$$\Rightarrow f(f(x)) = 15 - ||10 - x| - 5|$$

 \therefore Non-differentiable at points where

$$10 - x = 0$$
 and $|10 - x| = 5$

- \Rightarrow x = 10 and x 10 = ±5
- \Rightarrow x = 10 and x = 15, 5
- 16. The area (in sq. units) of the region $A = I(x, y) : x^2 < y < x + 21$ is :

$$A = \{(x, y) : x^2 \le y \le x + 2\} \text{ is } :$$

(1)
$$\frac{31}{6}$$
 (2) $\frac{10}{3}$

(3)
$$\frac{9}{2}$$
 (4) $\frac{13}{6}$

Answer (3)





-**1** , , ,

$$= \left(\frac{x^{2}}{2} + 2x - \frac{x^{3}}{3}\right)_{-1}^{2}$$
$$= \left(2 + 4 - \frac{8}{3}\right) - \left(+\frac{1}{2} - 2 + \frac{1}{3}\right)$$
$$= 8 - 3 - \frac{1}{2}$$
$$= 5 - \frac{1}{2} = \frac{9}{2}$$

17. If the line $y = mx + 7\sqrt{3}$ is normal to the hyperbola $\frac{x^2}{24} - \frac{y^2}{18} = 1$, then a value of m is : (1) $\frac{2}{\sqrt{5}}$ (2) $\frac{3}{\sqrt{5}}$

(3)
$$\frac{\sqrt{15}}{2}$$
 (4) $\frac{\sqrt{5}}{2}$

Answer (1)

Sol. $mx - y + 7\sqrt{3}$ is normal to hyperbola $\frac{x^2}{24} - \frac{y^2}{18} = 1$ then $\frac{24}{m^2} - \frac{18}{(-1)^2} = \frac{(24 + 18)^2}{(7\sqrt{3})^2}$ $\Rightarrow \frac{24}{m^2} - 18 = \frac{42 \times 42}{7 \times 7 \times 3}$ $\Rightarrow m = \frac{2}{\sqrt{5}}$ If lx + my + n = 0 is a normal to $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$, then $\frac{a^2}{l^2} - \frac{b^2}{m^2} = \frac{(a^2 + b^2)^2}{n^2}$

18. Let α and β be the roots of the equation $x^2 + x + 1 = 0$. Then for $y \neq 0$ in R,

$$\begin{vmatrix} y+1 & \alpha & \beta \\ \alpha & y+\beta & 1 \\ \beta & 1 & y+\alpha \end{vmatrix}$$
 is equal to :
(1) $y(y^2-1)$ (2) y^3-1
(3) $y(y^2-3)$ (4) y^3

$$\therefore \begin{vmatrix} y + \omega & 1 & \omega^{2} \\ 1 & y + \omega^{2} & \omega \\ \omega^{2} & \omega & 1 + y \end{vmatrix} \text{ operate } c_{1} \rightarrow c_{1} + c_{2} + c_{3} \\ = y \begin{vmatrix} 1 & 1 & \omega^{2} \\ 1 & y + \omega^{2} & \omega \\ 1 & \omega & 1 + y \end{vmatrix} \left(By \begin{array}{c} R_{2} \rightarrow R_{2} - R_{1} \\ R_{3} \rightarrow R_{3} - R_{1} \end{array} \right) \\ = y \begin{vmatrix} 1 & 1 & \omega^{2} \\ 0 & y + \omega^{2} - 1 & \omega - \omega^{2} \\ 0 & \omega - 1 & 1 + y - \omega^{2} \end{vmatrix} \\ = y\{(y + \omega^{2} - 1) (1 + y - \omega^{2}) - \omega(\omega - 1) (1 - \omega)\} \\ = y(y^{2} - (\omega^{2} - 1)^{2}) + y\omega(\omega - 1)^{2} \\ = y^{3} + y(\omega - 1)^{2} (\omega - (\omega + 1)^{2}) = y^{3} \end{aligned}$$

19. The value of $\cos^2 10^\circ - \cos 10^\circ \cos 50^\circ + \cos^2 50^\circ$ is :

(1)
$$\frac{3}{4}$$
 (2) $\frac{3}{2}(1+\cos 20^\circ)$

(3)
$$\frac{3}{2}$$
 (4) $\frac{3}{4} + \cos 20^{\circ}$

Answer (1)

Sol.
$$\left(\frac{1+\cos 20^{\circ}}{2}\right) + \left(\frac{1+\cos 100^{\circ}}{2}\right) - \frac{1}{2}(2\cos 10^{\circ}\cos 50^{\circ})$$

= $1 + \frac{1}{2}(\cos 20^{\circ} + \cos 100^{\circ}) - \frac{1}{2}[\cos 60^{\circ} + \cos 40^{\circ}]$
= $\left(1 - \frac{1}{4}\right) + \frac{1}{2}[\cos 20^{\circ} + \cos 100^{\circ} - \cos 40^{\circ}]$
= $\frac{3}{4} + \frac{1}{2}[2\cos 60^{\circ} \times \cos 40^{\circ} - \cos 40^{\circ}]$
= $\frac{3}{4}$

20. A committee of 11 members is to be formed from 8 males and 5 females. If m is the number of ways the committee is formed with at least 6 males and n is the number of ways the committee is formed with at least 3 females, then :

(1)
$$m = n = 68$$
 (2) $m + n = 68$
(3) $m = n = 78$ (4) $n = m - 8$

Answer (3)



21. Slope of a line passing through P(2, 3) and intersecting the line, x + y = 7 at a distance of 4 units from P, is :

(1)
$$\frac{\sqrt{7}-1}{\sqrt{7}+1}$$
 (2) $\frac{1-\sqrt{7}}{1+\sqrt{7}}$
(3) $\frac{\sqrt{5}-1}{\sqrt{5}+1}$ (4) $\frac{1-\sqrt{5}}{1+\sqrt{5}}$

Answer (2)

- Sol. Point at 4 units from P(2, 3) will be
 - A($4\cos\theta + 2$, $4\sin\theta + 3$) will satisfy x + y = 7

$$\Rightarrow \cos\theta + \sin\theta = \frac{1}{2} \text{ on squaring}$$

$$\Rightarrow \quad \sin 2\theta = \frac{-3}{4} \quad \Rightarrow \quad \frac{2\tan\theta}{1+\tan^2\theta} = -\frac{3}{4}$$

 \Rightarrow 3tan² θ + 8tan θ + 3 = 0

$$\Rightarrow$$
 tan $\theta = \frac{-8 \pm 2\sqrt{7}}{6}$ (Ignoring –ve sign)

$$\Rightarrow \tan \theta = \frac{-8 + 2\sqrt{7}}{6} = \frac{1 - \sqrt{7}}{1 + \sqrt{7}}$$

22. The value of
$$\int_{0}^{\pi/2} \frac{\sin^3 x}{\sin x + \cos x} dx$$
 is :

(1)
$$\frac{\pi - 2}{4}$$
 (2) $\frac{\pi - 2}{8}$
(3) $\frac{\pi - 1}{4}$ (4) $\frac{\pi - 1}{2}$

Answer (3)

Sol.
$$I = \int_{0}^{\frac{\pi}{2}} \frac{\sin^{3} x \, dx}{\sin x + \cos x}$$
$$\Rightarrow I = \int_{0}^{\frac{\pi}{2}} \frac{\cos^{3} x \, dx}{\sin x + \cos x}$$
$$\Rightarrow 2I = \int_{0}^{\frac{\pi}{2}} \left(1 - \frac{1}{2}\sin(2x)\right) dx$$

$$\Rightarrow I = \frac{1}{2} \left[\frac{x + \frac{1}{4} \cos 2x}{2} \right]_{0}$$
$$I = \frac{1}{2} \left(\frac{\pi - 1}{2} \right) = \frac{\pi - 1}{4}$$

23. Let S be the set of all values of x for which the tangent to the curve $y = f(x) = x^3 - x^2 - 2x$ at (x, y) is parallel to the line segment joining the points (1, f(1)) and (-1, f(-1)), then S is equal to :

(1)
$$\left\{\frac{1}{3}, -1\right\}$$
 (2) $\left\{-\frac{1}{3}, 1\right\}$
(3) $\left\{-\frac{1}{3}, -1\right\}$ (4) $\left\{\frac{1}{3}, 1\right\}$

Answer (2)

 \Rightarrow

 \Rightarrow

Sol. $y = f(x) = x^3 - x^2 - 2x$

$$\frac{dy}{dx} = 3x^2 - 2x - 2$$

f(1) = 1 - 1 - 2 = -2, f(-1) = -1 - 1 + 2 = 0
According to question, $3x^2 - 2x - 2$

$$= \frac{f(1) - f(-1)}{1 - (-1)}$$
$$3x^{2} - 2x - 2 = \frac{-2 - 0}{2}$$
$$3x^{2} - 2x - 1 = 0$$

$$\Rightarrow x = \frac{2 \pm 4}{6} = 1, \frac{-1}{3}$$

So,
$$S = \left\{\frac{-1}{3}, 1\right\}$$

24. A plane passing through the points (0, -1, 0)and (0, 0, 1) and making an angle $\frac{\pi}{4}$ with the plane y - z + 5 = 0, also passes through the point :

(1)
$$(\sqrt{2}, 1, 4)$$
 (2) $(\sqrt{2}, -1, 4)$

(3)
$$\left(-\sqrt{2}, -1, -4\right)$$
 (4) $\left(-\sqrt{2}, 1, -4\right)$

Answer (1)



plane is
$$v - z + 5 = 0$$

а

- .

$$\therefore \cos \frac{\pi}{4} = \frac{-1-1}{\sqrt{\frac{1}{a^2} + 1 + 1} \sqrt{2}}$$

$$\Rightarrow a^2 = \frac{1}{2}$$

$$\Rightarrow \frac{1}{a} = \pm \sqrt{2}$$

$$\Rightarrow \frac{\pm \sqrt{2}x - y + z = 1}{\sqrt{2}}$$

$$\therefore (\sqrt{2}, 1, 4) \text{ satisfies } -\sqrt{2}x - y + z = 1$$

25. The solution of the differential equation

$$x \frac{dy}{dx} + 2y = x^{2}(x \neq 0)$$
 with $y(1) = 1$, is :
(1) $y = \frac{4}{5}x^{3} + \frac{1}{5x^{2}}$ (2) $y = \frac{x^{3}}{5} + \frac{1}{5x^{2}}$
(3) $y = \frac{x^{2}}{4} + \frac{3}{4x^{2}}$ (4) $y = \frac{3}{4}x^{2} + \frac{1}{4x^{2}}$

Answer (3)

Sol.
$$\frac{dy}{dx} + \frac{2}{x}y = x$$
 $y(1) = 1$ (given)
 $I \cdot F = e^{\int \frac{2}{x} dx} = x^2$
 $y \times x^2 = \int x^3 dx$
 $\Rightarrow yx^2 = \frac{x^4}{4} + c$
 $\therefore at x = 1 ; y = 1$
 $\Rightarrow c = \frac{3}{4}$
 $\therefore y = \frac{x^2}{4} + \frac{3}{4x^2}$

constant A.P., a_1, a_2, a_3, \dots be 50 n + $\frac{n(n-7)}{2}A$,

where A is a constant. If d is the common difference of this A.P., then the ordered pair (d, a_{50}) is equal to :

Answer (3)

Sol. :
$$S_n = \left(50 - \frac{7A}{2}\right)n + n^2 \times \frac{A}{2}$$

: Common difference $= \frac{A}{2} \times 2 = \boxed{A}$
 $a_{50} = a_1 + 49 \times d$
 $= (50 - 3A) + 49 A$
 $= 50 + 46 A$
So, $(d, a_{50}) = (A, 50 + 46 A)$

27. For any two statements p and q, the negation of the expression p \lor (~ p \land q) is :

(1)
$$\sim p \land \sim q$$
(2) $\sim p \lor \sim q$ (3) $p \land q$ (4) $p \leftrightarrow q$

Answer (1)

Sol.
$$\sim (p \lor (\sim p \land q)) = \sim (\sim p \land q) \land \sim p$$

= $(\sim q \lor p) \land \sim p$
= $\sim p \land (p \lor \sim q)$
= $(\sim q \land \sim p) \lor (p \land \sim p)$
= $(\sim p \land \sim q)$

28. If the tangent to the curve, $y = x^3 + ax - b$ at the point (1, -5) is perpendicular to the line, -x + y + 4 = 0, then which one of the following points lies on curve?

(1) (–2, 1)	(2) (2, -2)
(3) (2, -1)	(4) (-2, 2)

Answer (2)

Sol.
$$f(x) = x^3 + ax - b \implies f'(x) = 3x^2 + a$$

 $f(1) = -5 \qquad \text{and} \qquad f'(1) = 3 + a$
 $\implies 1 + a - b = -5$
 $\implies a - b = -6 \qquad \dots(1)$
Also slope of tangent P(1, -5) = -1 = f'(1)
 $\therefore 3 + a = -1$
 $\implies a = -4$



- \therefore Equation of the out versities $f(x) = x + x^2$
- \therefore (2, -2) lies on the curve
- 29. If the function f : R {1, –1} \rightarrow A defined by

f(x) =
$$\frac{x^2}{1-x^2}$$
, is surjective, then A is equal to :
(1) [0, ∞) (2) R - {-1}

(3) R - (-1, 0) (4) R - [-1, 0)

Answer (4)

Sol.
$$f(x) = \frac{x^2}{1-x^2}$$

$$\Rightarrow f(-x) = \frac{x^2}{1-x^2} = f(x)$$

$$\Rightarrow f'(x) = \frac{2x}{\left(1 - x^2\right)^2}$$

∴ f(x) increases in x ∈ (0, ∞) Also f(0) = 0 $\lim_{x \to \pm \infty} f(x) = -1$

and F(x) is even function \therefore Set A \rightarrow R –[–1, 0)



30. Four persons can hit a target correctly with probabilities $\frac{1}{2}$, $\frac{1}{3}$, $\frac{1}{4}$ and $\frac{1}{8}$ respectively. If all hit at the target independently, then the probability that the target would be hit, is :

(1)
$$\frac{7}{32}$$
 (2) $\frac{25}{192}$
(3) $\frac{1}{192}$ (4) $\frac{25}{32}$

Sol. P(at least one) = 1 – P(none)

$$=1-\frac{1}{2}\times\frac{2}{3}\times\frac{3}{4}\times\frac{7}{8}$$

$$=1-\frac{7}{32}=\frac{25}{32}$$



Answers & Solutions

for

JEE (MAIN)-2019 (Online) Phase-2

(Physics, Chemistry and Mathematics)

Time : 3 hrs.

M.M.: 360

Important Instructions :

- 1. The test is of **3 hours** duration.
- 2. The Test Booklet consists of **90** questions. The maximum marks are **360**.
- 3. There are *three* parts in the question paper A, B, C consisting of **Physics**, **Chemistry** and **Mathematics** having 30 questions in each part of equal weightage.
- 4. Each question is allotted 4 (four) marks for each correct response. ¼ (one-fourth) marks will be deducted for indicating incorrect response of each question. No deduction from the total score will be made if no response is indicated for an item in the answer sheet.
- 5. There is only one correct response for each question.



1. A test particle is moving in a circular orbit in the gravitational field produced by a mass

density $\rho(\mathbf{r}) = \frac{\mathbf{K}}{\mathbf{r}^2}$. Identify the correct relation

between the radius R of the particle's orbit and its period T :

- (1) T/R is a constant (2) TR is a constant
- (3) T/R^2 is a constant (4) T^2/R^3 is a constant

Answer (1)

Sol.
$$M_{(in)} = \int_{0}^{K} 4\pi r^{2} dr \cdot \frac{k}{r^{2}}$$

 $M_{(in)} = 4\pi KR$
 $\therefore \quad G \cdot \frac{4\pi KR}{R^{2}} = \frac{V^{2}}{R} \implies v = \sqrt{4\pi GK}$
 $\therefore \quad T = \frac{2\pi R}{v} = \frac{2\pi \cdot R}{\sqrt{4\pi GK}}$
 $\therefore \quad \frac{T}{R} = \text{constant}$

- 2. The specific heats, C_p and C_V of a gas of diatomic molecules, A, are given (in units of J mol⁻¹ K⁻¹) by 29 and 22, respectively. Another gas of diatomic molecules, B, has the corresponding values 30 and 21. If they are treated as ideal gases, then :
 - (1) A is rigid but B has a vibrational mode.
 - (2) A has a vibrational mode but B has none.
 - (3) Both A and B have a vibrational mode each.

(4) A has one vibrational mode and B has two. Answer (2)

Sol. For (A) $C_P = 29$, $C_V = 22$

For (B) $C_{p} = 30 C_{v} = 21$

$$\therefore \quad \gamma_{A} = \frac{C_{P_{A}}}{C_{V_{A}}} = \frac{29}{22} = 1.31$$

When A has vibrational degree of freedom, then γ_{A} = 9/7 \simeq 1.29.

$$\gamma_{B} = \frac{C_{P_{B}}}{C_{V_{B}}} = \frac{30}{21} = 1.42$$

 \Rightarrow B has no vibrational degree of freedom

3. The position vector of a particle changes with time according to the relation $\vec{r}(t) = 15t^2\hat{i} + (4-20t^2)\hat{j}$. What is the magnitude of the acceleration at t = 1?

Answer (1)

Sol.
$$\vec{r} = 15t^2\hat{i} + 4\hat{j} - 20t^2\hat{j}$$

$$\frac{d\vec{r}}{dt} = 30t\hat{i} - 40t\hat{j}$$
$$\frac{d^2\vec{r}}{dt^2} = 30\hat{i} - 40\hat{j}$$
$$\therefore \quad \left|\frac{d^2\vec{r}}{dt^2}\right| = 50 \text{ m/s}^2$$

- 4. 50 W/m² energy density of sunlight is normally incident on the surface of a solar panel. Some part of incident energy (25%) is reflected from the surface and the rest is absorbed. The force exerted on 1 m² surface area will be close to ($c = 3 \times 10^8$ m/s) :
 - (1) 20 × 10⁻⁸ N
 (2) 35 × 10⁻⁸ N
 (3) 15 × 10⁻⁸ N
 (4) 10 × 10⁻⁸ N

...

Answer (1)

Sol.
$$\therefore$$
 $P = \frac{W}{c}$ and pressure = I/c
 $\therefore \frac{h}{\lambda} + \frac{h}{4\lambda} \Rightarrow \Delta P = \frac{5h}{4\lambda}$ for one photon
 $\therefore \frac{5}{4} \cdot \frac{Nh}{\lambda \Delta tA} = \text{pressure}$
But $\left(\frac{Nhc}{\lambda \cdot \Delta tA}\right) = 50 \text{ W/m}^2$
 $\therefore \frac{5}{4} \times \frac{50}{c} = \text{pressure}$
 $\therefore F_n = \frac{5 \times 50 \times 1 \text{ m}^2}{4 \times c} = 20 \times 10^{-8} \text{ N}$



length. This new wire is now bent and the ends joined to make a circle. If two points on this circle make an angle 60° at the centre, the equivalent resistance between these two points will be :

(1)
$$\frac{5}{3} \Omega$$
 (2) $\frac{5}{2} \Omega$
(3) $\frac{7}{2} \Omega$ (4) $\frac{12}{5} \Omega$

Answer (1)

Sol.

$$\begin{array}{c}
\mathbf{B} \\
\mathbf{R}_{1} \\
\mathbf{R}_{2} \\
\mathbf{R}_{0} = \rho \frac{\mathbf{I}}{\mathbf{A}} = \mathbf{3} \Omega \\
\mathbf{R}_{1} = 2\mathbf{I} \\
\mathbf{Then } \mathbf{A} = \frac{\mathbf{A}}{2} \\
\therefore \quad \mathbf{R} = \frac{\rho 2\mathbf{I} \times 2}{\mathbf{A}} = \mathbf{12} \Omega \\
\mathbf{R}_{1} = \frac{\mathbf{12}}{\mathbf{6}} = \mathbf{2} \Omega \\
\mathbf{R}_{2} = \mathbf{10} \Omega \\
\therefore \quad \frac{\mathbf{1}}{\mathbf{R}_{eq}} = \frac{\mathbf{1}}{2} + \frac{\mathbf{1}}{\mathbf{10}} = \frac{\mathbf{6}}{\mathbf{10}} \\
\therefore \quad \mathbf{R}_{eq} = \frac{\mathbf{5}}{3} \Omega
\end{array}$$

6. A thin smooth rod of length L and mass M is rotating freely with angular speed ω_0 about an axis perpendicular to the rod and passing through its center. Two beads of mass m and negligible size are at the center of the rod initially. The beads are free to slide along the rod. The angular speed of the system, when the beads reach the opposite ends of the rod, will be :

(1)
$$\frac{M\omega_0}{M+3m}$$

(2)
$$\frac{W\omega_0}{M+m}$$

$$(3) \quad \frac{\mathbf{M}\,\omega_{\mathbf{0}}}{\mathbf{M}+\mathbf{6m}}$$

$$(4) \quad \frac{M\omega_0}{M+2m}$$

Momentum

$$\frac{ML^{2}}{12}\omega_{0} = \left(\frac{ML^{2}}{12} + 2\frac{mL^{2}}{4}\right)\omega$$
$$\Rightarrow \omega = \frac{M\omega_{0}}{M + 6m}$$

 Moment of inertia of a body about a given axis is 1.5 kg m². Initially the body is at rest. In order to produce a rotational kinetic energy of 1200 J, the angular acceleration of 20 rad/s² must be applied about the axis for a duration of

Answer (2)

Sol. I = 150;
$$\alpha = 20 \text{ rad/s}^2$$

 $\therefore \omega = \alpha t$
 $E = \frac{1}{2}I\omega^2 = 1200$
 $\frac{1}{2} \times 1.5 \times (20t)^2 = 1200 \text{ J}$
 $\Rightarrow t = 2 \text{ s}$

- 8. The resistance of a galvanometer is 50 ohm and the maximum current which can be passed through it is 0.002 A. What resistance must be connected to it in order to convert it into an ammeter of range 0 - 0.5 A?
 - (1) 0.2 ohm
 - (2) 0.002 ohm
 - (3) 0.5 ohm
 - (4) 0.02 ohm

Answer (1)



We have

$$I_g R_g = (0.5 - I_g) S$$

$$\Rightarrow (0.002) (50) = (0.5 - I_g) S$$

$$\Rightarrow S \simeq \frac{0.002 \times 50}{0.5} = 0.2 \Omega$$


(1) 13.60 eV	(2) 6.04 eV
(3) 48.36 eV	(4) 54.40 eV

Answer (1)

Sol.
$$E_n: \frac{-E_0 z^2}{n^2} = \frac{-E_0 \times 4}{4} = -E_0$$

To ionise it E_0 energy must be supplied.

$$E_0 = 13.6 \text{ eV}.$$

10. Two cars A and B are moving away from each other in opposite directions. Both the cars are moving with a speed of 20 ms⁻¹ with respect to the ground. If an observer in car A detects a frequency 2000 Hz of the sound coming from car B, what is the natural frequency of the sound source in car B?

(speed of sound in air = 340 ms^{-1})

- (1) 2150 Hz (2) 2300 Hz
- (3) 2060 Hz (4) 2250 Hz

Answer (4)

Sol.
$$\underbrace{\begin{array}{c} S \\ 20 \text{ m/s} \end{array}}_{20 \text{ m/s}} \underbrace{\begin{array}{c} 0 \\ 20 \text{ m/s} \end{array}}_{20 \text{ m/s}}$$
$$f = \frac{(v \pm u_0)}{(v \pm u_s)} \cdot f_0 = \frac{(v - 20)}{(v + 20)} \cdot f_0$$
$$\Rightarrow 2000 = \frac{320}{360} \cdot f_0$$
$$\Rightarrow \frac{2000 \times 9}{8} = f_0 = 2250 \text{ Hz}.$$

11. A convex lens of focal length 20 cm produces images of the same magnification 2 when an object is kept at two distances x_1 and x_2 $(x_1 > x_2)$ from the lens. The ratio of x_1 and x_2 is :

(1) 3:1	(2) 2:1
(3) 4:3	(4) 5:3

Answer (1)

Sol.
$$\therefore \quad \frac{1}{v} - \frac{1}{u} = \frac{1}{f}$$

 $\therefore \quad v = (\pm 2u)$
 $\therefore \quad \frac{1}{2u} + \frac{1}{u} = \frac{1}{20} \quad \Rightarrow \frac{3}{2u} = \frac{1}{20}$
 $\therefore \quad u_1 = 30 \text{ cm}$
And $\qquad \frac{1}{u} - \frac{1}{2u} = \frac{1}{20}$

$$\therefore \quad \frac{30}{10} = 3.$$

12. The position of a particle as a function of time t, is given by

 $\mathbf{x(t)} = \mathbf{at} + \mathbf{bt}^2 - \mathbf{ct}^3$

where a, b and c are constants. When the particle attains zero acceleration, then its velocity will be :

(1)
$$a + \frac{b^2}{4c}$$
 (2) $a + \frac{b^2}{3c}$
(3) $a + \frac{b^2}{2c}$ (4) $a + \frac{b^2}{c}$

Answer (2)

Sol. x = at + bt² - ct³
x = a + 2bt - 3ct²
x = 2b - 6ct
For
$$\ddot{x} = 0$$
 $t = +\frac{b}{3c}$
 \therefore $v = \dot{x} = a + 2b\left(\frac{+b}{3c}\right) - 3c\left(\frac{b^2}{3c \times 3c}\right)$
 $\Rightarrow v = -\frac{b^2}{3c} + \frac{2b^2}{3c} + a = a + \frac{b^2}{3c}$

13. The parallel combination of two air filled parallel plate capacitors of capacitance C and nC is connected to a battery of voltage, V. When the capacitors are fully charged, the battery is removed and after that a dielectric material of dielectric constant K is placed between the two plates of the first capacitor. The new potential difference of the combined system is :

(1)
$$\frac{(n+1)V}{(K+n)}$$
 (2) $\frac{V}{K+n}$

Answer (1)

Sol. Initially

$$Q = CV(1 + n)$$

$$\therefore C_{eq} = (K + n)C$$

$$\therefore V = \frac{CV(1+n)}{(K+n)C} = \frac{V(1+n)}{(K+n)}$$
nCV

(4) $\frac{nV}{K+n}$



are:

- (1) Inversely proportional to modulation frequency
- (2) Inversely proportional to carrier frequency
- (3) Proportional to carrier frequency
- (4) Independent of both carrier and modulation frequency

Answer (2)

- Sol. Size of antenna depends on wavelength of carrier wave.
- 15. A wedge of mass M = 4m lies on a frictionless plane. A particle of mass m approaches the wedge with speed v. There is no friction between the particle and the plane or between the particle and the wedge. The maximum height climbed by the particle on the wedge is given by :

(1)
$$\frac{v^2}{g}$$
 (2) $\frac{2v^2}{5g}$
(3) $\frac{2v^2}{7g}$ (4) $\frac{v^2}{2g}$

Answer (2)

Sol. mv = (4m + m)v'

$$\therefore \quad \text{Common speed } \mathbf{v}' = \frac{\mathbf{v}}{5}$$

$$\text{mgh} + \frac{1}{2}5\text{m} \cdot \frac{\mathbf{v}^2}{25} = \frac{1}{2}\text{mv}^2$$

$$\Rightarrow \quad \text{mgh} = \frac{1}{2}\text{mv}^2 \left(1 - \frac{1}{5}\right) = \frac{1}{2}\text{mv}^2 \cdot \frac{4}{5}$$

$$\therefore \quad \mathbf{h} = \frac{2\text{mv}^2}{5 \times \text{mg}} = \frac{2\text{v}^2}{5\text{g}}$$

16. A wooden block floating in a bucket of water

has $\frac{4}{5}$ of its volume submerged. When certain amount of an oil is poured into the bucket, it is found that the block is just under the oil surface

with half of its volume under water and half in oil. The density of oil relative to that of water is

(1) 0.5	(2)	0.8
---------	-----	-----

(3) 0.7 (4) 0.6

Answer (4)

$$\begin{split} \mathbf{V} \sigma \mathbf{g} &= \frac{\mathbf{v}}{2} \rho_{\omega} \mathbf{g} + \frac{\mathbf{v}}{2} \rho_{0} \mathbf{g} \\ \Rightarrow & \left(\frac{\rho_{\omega}}{2} + \frac{\rho_{oil}}{2} \right) = \frac{4}{5} \rho_{\omega} \\ \Rightarrow & \frac{\rho_{oil}}{2} = \rho_{\omega} \left(\frac{4}{5} - \frac{1}{2} \right) = \frac{3}{10} \rho_{\omega} \\ \Rightarrow & \rho_{oil} = \frac{3}{5} \rho_{\omega} = \mathbf{0.6} \rho_{\omega} \end{split}$$

17. Two materials having coefficients of thermal conductivity '3K' and 'K' and thickness 'd' and '3d', respectively, are joined to form a slab as shown in the figure. The temperatures of the outer surfaces are ' θ_2 ' and ' θ_1 ' respectively, ($\theta_2 > \theta_1$). The temperature at the interface is:

$$\begin{array}{c|cccc} d & 3d \\ \theta_2 & 3K & K \\ \end{array} \\ (1) \quad \frac{\theta_1}{6} + \frac{5\theta_2}{6} \\ (2) \quad \frac{\theta_1}{10} + \frac{9\theta_2}{10} \\ (3) \quad \frac{\theta_1}{3} + \frac{2\theta_2}{3} \\ \end{array} \\ (4) \quad \frac{\theta_2 + \theta_1}{2} \end{array}$$

Answer (2)

Sol.
$$\mathbf{H} = \frac{\mathbf{3KA}}{\mathbf{d}} (\theta_2 - \theta) = \frac{\mathbf{KA}}{\mathbf{3d}} (\theta - \theta_1)$$

$$\Rightarrow \theta = \frac{\mathbf{9}\theta_2}{\mathbf{10}} + \frac{\theta_1}{\mathbf{10}}$$

18. The area of a square is 5.29 cm². The area of 7 such squares taking into account the significant figures is :

(1) 37.03 cm^2 (2)	2)	37.0	cm ²
------------------------------	----	------	-----------------

(3)	37.030	cm ²	(4)	37	cm ²
-----	--------	-----------------	-----	----	-----------------

Answer (2)

Sol. $5.29 \times 7 = 37.0 \text{ cm}^2$

Answer should be in 3 significant digits.

- 19. Diameter of the objective lens of a telescope is 250 cm. For light of wavelength 600 nm. coming from a distant object, the limit of resolution of the telescope is close to :
 - (1) 1.5×10^{-7} rad
 - (2) 3.0×10^{-7} rad
 - (3) 2.0×10^{-7} rad
 - (4) 4.5×10^{-7} rad



Sol.
$$\theta = \frac{1.22 \lambda}{D}$$

 $\Rightarrow \theta = \frac{1.22 \times 600 \times 10^{-9}}{250} \times 100 = 2.92 \times 10^{-7}$
 $\Rightarrow \theta = 3 \times 10^{-7} \text{ rad}$

20. A particle 'P' is formed due to a completely inelastic collision of particles 'x' and 'y' having de-Broglie wavelengths ' λ_x ' and ' λ_y ' respectively. If x and y were moving in opposite directions, then the de-Broglie wavelength of 'P' is

(1)
$$\frac{\lambda_x \lambda_y}{|\lambda_x - \lambda_y|}$$
 (2) $\lambda_x - \lambda_y$
(3) $\lambda_x + \lambda_y$ (4) $\frac{\lambda_x \lambda_y}{\lambda_x + \lambda_y}$

Answer (1)

Sol.
$$P_1 = \frac{h}{\lambda_x}$$

 $P = P_1 - P_2 = h\left(\frac{1}{\lambda_x} - \frac{1}{\lambda_y}\right)$
 $\therefore P = \frac{h}{\lambda}$
 $\frac{h}{\lambda} = h\left(\frac{1}{\lambda_x} - \frac{1}{\lambda_y}\right)$
 $\frac{1}{\lambda} = \frac{|\lambda_y - \lambda_x|}{\lambda_x \lambda_y}$
 $\lambda = \frac{\lambda_x \cdot \lambda_y}{|\lambda_x - \lambda_y|}$

21. A moving coil galvanometer has a coil with 175 turns and area 1 cm². It uses a torsion band of torsion constant 10^{-6} N-m/rad. The coil is placed in a magnetic field B parallel to its plane. The coil deflects by 1° for a current of 1 mA. The value of B (in Tesla) is approximately

(1) 10⁻⁴ (2) 10⁻²

(3) 10⁻¹ (4) 10⁻³

Answer (4)

Sol. NIAB = KQ

175 × 1 × 10⁻³ × 1 × 10⁻⁴ × B =
$$\frac{10^{-6} \times \pi}{180}$$

⇒ B = $\frac{\pi}{180} \times \frac{10}{175} \approx 9.97 \times 10^{-4}$ T.
⇒ B = 10⁻³ T.

mean free time is 25 fs (femto second), it's approximate resistivity is

Answer (4)

Sol. J =
$$\eta e v_d$$
 \therefore $v_d = \frac{e E \tau}{m}$

$$\therefore \quad \mathbf{J} = \frac{\eta \mathbf{e} \cdot \mathbf{e} \mathbf{E} \cdot \tau}{\mathbf{m}} = \sigma \vec{\mathbf{E}}$$
$$\mathbf{J} = \frac{\eta \mathbf{e}^2 \tau}{\mathbf{m}} \cdot \mathbf{E} = \sigma \vec{\mathbf{E}}$$
$$\therefore \quad \rho = \frac{\mathbf{m}}{\eta \mathbf{e}^2 \tau}$$

$$= \frac{9.1 \times 10^{-31}}{8.5 \times 10^{28} \times (1.6 \times 10^{-19})^2 \times 25 \times 10^{-15}} \Omega m$$
$$= \frac{9.1}{8.5 \times 2.56 \times 25} \times 10^{(-59+53)} = 1.67 \times 10^{-8} \Omega m$$

23. A thin convex lens L (refractive index = 1.5) is placed on a plane mirror M. When a pin is placed at A, such that OA = 18 cm, its real inverted image is formed at A itself, as shown in figure. When a liquid of refractive index μ_l is put between the lens and the mirror, the pin has to be moved to A', such that OA' = 27 cm, to get its inverted real image at A' itself. The value of μ_l will be



Answer (2)

Sol.
$$f_L = 18$$
 cm

$$\frac{1}{18} = 0.5 \times \frac{2}{R} \implies R = 18 \text{ cm}$$
$$\frac{1}{f_2} = (\mu_1 - 1) \left(-\frac{1}{18} \right)$$



$$\Rightarrow 2 = 3(2 - \mu_{l}) = 6 - 3 \mu_{l}$$
$$\Rightarrow \mu_{l} = \frac{4}{3}$$

24. Two coils 'P' and 'Q' are separated by some distance. When a current of 3 A flows through coil 'P', a magnetic flux of 10⁻³ Wb passes through Q. No current is passed through 'Q'. When no current passes through 'P' and a current of 2 A passes through 'Q', the flux through 'P' is

(1) 6.67 × 10 ⁻³ Wb	(2) 6.67 × 10 ⁻⁴ Wb
(3) 3.67 × 10 ⁻³ Wb	(4) 3.67 × 10 ^{−4} Wb

Sol.
$$\phi_Q = Mi$$

 $\Rightarrow 10^{-3} = M(3)$
 $\phi_P = M(2)$
 $\therefore \frac{10^{-3}}{3} = \frac{\phi_P}{2}$
 $\phi_P = \frac{20}{3} \times 10^{-4} = 6.67 \times 10^{-4}$ Wb.

25. A massless spring (k = 800 N/m), attached with a mass (500 g) is completely immersed in 1 kg of water. The spring is stretched by 2 cm and released so that it starts vibrating. What would be the order of magnitude of the change in the temperature of water when the vibrations stop completely? (Assume that the water container and spring receive negligible heat and specific heat of mass = 400 J/kg K, specific heat of water = 4184 J/kg K)

(1)	10 ^{–5} K	(2)	10 ⁻¹ K
(3)	10 ⁻³ K	(4)	10 ⁻⁴ K

Sol.
$$\frac{1}{2} k\Delta x^{2} = E(\text{dissipated})$$

$$\therefore \quad \frac{1}{2} \times 800 \times \left(\frac{2 \times 2}{100 \times 100}\right) = \frac{16}{100} J$$

$$\frac{16}{100} = \frac{1}{2} \times 400 \times \Delta T + 1 \times 4184 \times \Delta T$$

$$\Rightarrow \quad \frac{16}{100} = (200 + 4184) \Delta T = 4384 \Delta T$$

$$\therefore \quad \Delta T = \frac{16}{4384 \times 100} = 3.6 \times 10^{-5} \text{ K}$$

speed 'v' in the same direction. After collision, the first mass is stopped completely while the second one splits into two particles each of mass 'm', which move at angle 45° with respect to the original direction.

The speed of each of the moving particle will be

(1)
$$v/(2\sqrt{2})$$
 (2) $v/\sqrt{2}$

(3) $\sqrt{2}v$ (4) $2\sqrt{2}v$

Answer (4)

Sol. Initial momentum $\cdot P_i = 2mv + 2mv = 4 mv$

Let v' be the speed of I particle



- 27. A string 2.0 m long and fixed at its ends is driven by a 240 Hz vibrator. The string vibrates in its third harmonic mode. The speed of the wave and its fundamental frequency is
 - (1) 320 m/s, 120 Hz
 - (2) 320 m/s, 80 Hz
 - (3) 180 m/s, 80 Hz
 - (4) 180 m/s, 120 Hz

Answer (2)

Sol.
$$\frac{\lambda}{2} = \frac{2}{3}$$

 $\Rightarrow \lambda = \frac{4}{3}m$
 $\therefore v = \frac{4}{3} \times 240 = 320 \text{ m/s}$
 3^{rd} harmonic
 $f_n = nf_0$
 $f_0 = \frac{240}{3} = 80 \text{ Hz}$





- (1) OR
- (2) NAND
- (3) AND
- (4) NOR

Answer (1)

Sol. $y = \overline{\overline{A} \cdot \overline{B}} = A + B$

 $\mathbf{y}=\mathbf{A}+\mathbf{B}$

- \Rightarrow OR gate
- 29. A very long solenoid of radius R is carrying current I(t) = $kte^{-\alpha t}$ (k > 0), as a function of time (t \ge 0). Counter clockwise current is taken to be positive. A circular conducting coil of radius 2R is placed in the equatorial plane of the solenoid and concentric with the solenoid. The current induced in the outer coil is correctly depicted, as a function of time, by:







$$\therefore \quad \phi = \mathbf{k}_1 \mathbf{i} = \mathbf{k}_1 \mathbf{t} \mathbf{e}^{-\alpha t} \qquad [\mathbf{k}_1 = \mathbf{k} \pi (2\mathbf{R})^2]$$
$$\therefore \quad \mathbf{E} = -\left(\frac{\mathbf{d}\phi}{\mathbf{d}t}\right) = -\mathbf{k}_1 \mathbf{e}^{-\alpha t} + \mathbf{k}_1 \alpha \mathbf{t} \mathbf{e}^{-\alpha t}$$
$$= -\mathbf{k}_1 \mathbf{e}^{-\alpha t} (1 - \alpha t)$$
$$\therefore \quad \text{Induced current } \mathbf{I} = \frac{\mathbf{E}}{\mathbf{r}} = -\mathbf{k}_1 \mathbf{e}^{-\alpha t} (1 - \alpha t)$$

30. For point charges -q, +q, +q and -q are placed on y-axis at y = -2d, y = -d, y = +d and y =+2d, respectively. The magnitude of the electric field E at a point on the x-axis at x = D, with D >> d, will behave as:

(1)
$$E \propto \frac{1}{D^3}$$

(2) $E \propto \frac{1}{D}$
(3) $E \propto \frac{1}{D^4}$
(4) $E \propto \frac{1}{D^2}$

Answer (3)





1. The major product of the following reaction is :





Answer (2)



Acid catalysed intramolecular esterification



- 2. The peptide that gives positive ceric ammonium nitrate and carbylamine tests is
 - (1) Ser Lys
 (2) Lys Asp
 (3) Gln Asp
 (4) Asp Gln

Answer (1)

- Sol. Ceric ammonium nitrate test is given by alcohol. Only serine(ser) contain –OH group.
- 3. Which of the following compounds is a constituent of the polymer

- (1) Formaldehyde (2) Ammonia
- (3) Methylamine

(4) N-Methyl urea

Answer (1)

Sol. $\left\{ \mathbf{NH} = \mathbf{C} = \mathbf{NH} = \mathbf{CH}_2 \right\}$ -urea-formaldehyde resin

... Monomers : Urea and formaldehyde

4. During compression of a spring the work done is 10 kJ and 2 kJ escaped to the surroundings as heat. The change in internal energy, ΔU (in kJ) is

Answer (3)

 $\Delta U = q + w = 10 - 2 = 8 \text{ kJ}$

- What would be the molality of 20% (mass/mass) aqueous solution of KI? (molar mass of KI = 166 g mol⁻¹)
 - (1) 1.48 (2) 1.51
 - (3) 1.08 (4) 1.35

Answer (2)

..

Sol. 20% W/W KI solution

- i.e. 100 g solution contains 20 g KI
- ∴ Mass of solvent = 100 20 = 80 g

$$Molality = \frac{20 \times 1000}{166 \times 80}$$

$$\simeq$$
 1.51 molar

- 6. A solution of $Ni(NO_3)_2$ is electrolysed between platinum electrodes using 0.1 Faraday electricity. How many mole of Ni will be deposited at the cathode?
 - (1) 0.20
 - (2) 0.15
 - (3) 0.10
 - (4) 0.05

Answer (4)

- Sol. 0.1 F of electricity is passed through $\mathrm{Ni}(\mathrm{NO}_3)_2$ solution
 - ... Amount of Ni deposited = 0.1 eq

:.. Moles = $\frac{0.1}{2}$ = 0.05



- (1) Strongest hydrogen bonding
- (2) Lowest dissociation enthalpy
- (3) Strongest van der Waals' interactions
- (4) Lowest ionic character

Answer (1)

- Sol. HF has highest boiling point among the hydrogen halides due to strong H-bonding between HF molecules.
- 8. The one that is not a carbonate ore is
 - (1) Bauxite (2) Calamine
 - (3) Siderite (4) Malachite

Answer (1)

Sol. Bauxite ightarrow AlO_x (OH)_{3–2x}

 $\textbf{Calamine} \rightarrow \textbf{ZnCO}_{\textbf{3}}$

- $\textbf{Siderite} \rightarrow \textbf{FeCO}_{\textbf{3}}$
- Siderite \rightarrow FeCO₃

Malachite \rightarrow CuCO₃ \cdot Cu(OH)₂

- 9. Noradrenaline is a / an
 - (1) Neurotransmitter (2) Antihistamine
 - (3) Antacid (4) Antidepressant

Answer (1)

- Sol. Noradrenaline is neurotransmitter.
- 10. Among the following species, the diamagnetic molecule is

(1)	CO	(2)	NO
(3)	O ₂	(4)	B ₂

- Answer (1)
- Sol. MoleculeNo. of unpaired electronsNO1COZero O_2 2 B_2 2
 - ... Diamagnetic species is CO
- 11. The correct statements among I to III regarding group 13 element oxides are,
 - (I) Boron trioxide is acidic.
 - (II) Oxides of aluminium and gallium are amphoteric.
 - (III) Oxides of indium and thallium are basic.
 - (1) (II) and (III) only (2) (I) and (II) only
 - (3) (I), (II) and (III) (4) (I) and (III) only

 $\mathbf{D}_2 \mathbf{D}_3$ is all actual online

 AI_2O_3 and Ga_2O_3 are amphoteric oxide

 In_2O_3 and TI_2O are basic oxide

12. The major products A and B for the following reactions are, respectively



Answer (4)



13. Increasing order of reactivity of the following compounds for $\rm S_{N}1$ substitution is







(3) (B) < (A) < (D) < (C)

(d) (d)
$$<$$
 (d) $<$ (

Answer (3)

Sol. S_N1 reaction proceeds via formation of carbocation.



On comparing (A) and (B), in (A) there is formation of tertiary carbocation $CH_3 - c$

after rearrangement while (B) is primary.

So, (C) > (D) > (A) > (B).

14. Which of the following potential energy (PE) diagrams represents the S_N1 reaction?



- (1^{st} step) is rate determining step (RDS)
 - \therefore Correct graph is given in option-4.
- 15. Molal depression constant for a solvent is 4.0 K kg mol⁻¹. The depression in the freezing point of the solvent for 0.03 mol kg⁻¹ solution of K_2SO_4 is

(Assume complete dissociation of the electrolyte)

- (1) 0.36 K
- (2) 0.18 K
- (3) 0.12 K
- (4) 0.24 K

Answer (1)

Sol.
$$K_2SO_4 \longrightarrow 2K^+ + SO_4^{2-}$$

i (Van't Hoff Factor) = 3
 $\therefore \Delta T_f = iK_fm$
= 3 × 4 × 0.03
= 0.36 K

16. Consider the given plot of enthalpy of the following reaction between A and B.

 $A + B \rightarrow C + D$

Identify the incorrect statement.



- (1) Activation enthalpy to form C is 5 kJ mol⁻¹ less than that to form D
- (2) D is kinetically stable product
- (3) Formation of A and B from C has highest enthalpy of activation
- (4) C is the thermodynamically stable product

Answer (1)

Sol. Activation enthalpy to form C is 5 kJ more than that to form D.



strength. Which of the following correctly shows the change of pH of the titration mixture in this experiment?



Answer (1)

- **Sol.** The pH of NaOH is more than 7 and during the titration it decreases so graph (1) is correct
- 18. Hinsberg's reagent is
 - (1) $C_6H_5SO_2CI$ (2) $(COCI)_2$
 - (3) C_6H_5COCI (4) $SOCI_2$

Answer (1)

Sol. Hinsberg's reagent is benzenesulphonyl chloride



- 19. The layer of atmosphere between 10 km to 50 km above the sea level is called as
 - (1) Stratosphere (2) Mesosphere
 - (3) Thermosphere (4) Troposphere

Answer (1)

- Sol. Between 10-50 km above sea level lies stratosphere.
- 20. Assertion : For the extraction of iron, haematite ore is used.

Reason : Haematite is a carbonate ore of iron.

- (1) Only the reason is correct
- (2) Only the assertion is correct

for the assertion

(4) Both the assertion and reason are correct, but the reason is not the correct explanation for the assertion

Answer (2)

Sol. For the extraction of iron, haematite ore in used.

Haematite = Fe_2O_3

21. At a given temperature T, gases Ne, Ar, Xe and Kr are found to deviate from ideal gas behaviour. Their equation of state is given as

$$P = \frac{RT}{V-b}$$
 at T.

Here, b is the van der Waal's constant. Which gas will exhibit steepest increase in the plot of Z (compression factor) vs P?

- (1) Kr
- (2) Ar
- (3) Xe
- (4) Ne
- Answer (3)

Sol. P =
$$\frac{RT}{V_m - b}$$

 $\Rightarrow PV_m - Pb = RT$
 $\Rightarrow \frac{PV_m}{RT} = 1 + \frac{Pb}{RT}$
 $\Rightarrow Z = 1 + \frac{Pb}{RT}$

Slope of Z vs P curve (straight line) = $\frac{b}{PT}$

- \therefore Higher the value of b, more steep will be the curve and b \propto size of gas molecules
- 22. The amorphous form of silica is
 - (1) Quartz
 - (2) Tridymite
 - (3) Kieselguhr
 - (4) Cristobalite

Answer (3)

Sol. Quartz, tridymite and cristobalite are crystalline forms of silica.

Kieselguhr is an amorphous form of silica.



- color exhibited by transition metal complexes.
- (II) Valence bond theory can predict quantitatively the magnetic properties of transition metal complexes.
- (III) Valence bond theory cannot distinguish ligands as weak and strong field ones.
- (1) (II) and (III) only (2) (I), (II) and (III)
- (3) (I) and (II) only (4) (I) and (III) only

Answer (4)

- Sol. Valence bond theory cannot predict quantitatively the magnetic properties of transition metal complex.
- 24. In the following reaction

carbonyl compound + MeOH _____ acetal

Rate of the reaction is the highest for

- (1) Acetone as substrate and methanol in excess
- (2) Propanal as substrate and methanol in stoichiometric amount
- (3) Propanal as substrate and methanol in excess
- (4) Acetone as substrate and methanol in stoichiometric amount

Answer (3)

Generally, aldehydes are more reactive than ketones in nucleophilic addition reactions.

∴ Rate of reaction with alcohol to form acetal and ketal is

$$CH_3-CH_2-CHO > CH_3-C-CH_3$$

- 25. The structures of beryllium chloride in the solid state and vapour phase, respectively, are
 - (1) Chain and dimeric (2) Dimeric and dimeric
 - (3) Dimeric and chain (4) Chain and chain

Answer (1)

Sol. BeCl₂ in vapour phase exist as dimer (below 1200 K temperature)

BeCl₂ in solid state has chain structure.



Answer (4)



Product will formed as per -OH group (+M group)

27. The maximum possible denticities of a ligand given below towards a common transition and inner-transition metal ion, respectively, are



Answer (1)

- **Sol.** The maximum possible denticities of the given ligand towards transition metal ion is 6 and towards inner transition metal ion is 8.
- 28. 10 mL of 1 mM surfactant solution forms a monolayer covering 0.24 cm² on a polar substrate. If the polar head is approximated as a cube, what is its edge length?

(1) 2	.0 pm	(2)	2.0 nm
(3) 0	.1 nm	(4)	1.0 pm



Sol. No. of surfactant molecule $6 \times 10^{23} \times \frac{10}{1000} \times 10^{-3}$

 \Rightarrow 6 × 10¹⁸ molecule

Let edge length = a cm

Total surface area of surfactant = $6 \times 10^{18} a^2$

 \Rightarrow 0.24

 $a = 2 \times 10^{-10} \text{ cm} = 2 \text{ pm}$

- 29. Which one of the following about an electron occupying the 1s orbital in a hydrogen atom is incorrect? (The Bohr radius is represented by a₀)
 - (1) The probability density of finding the electron is maximum at the nucleus
 - (2) The electron can be found at a distance $2a_0$ from the nucleus
 - (3) The magnitude of the potential energy is double that of its kinetic energy on an average
 - (4) The total energy of the electron is maximum when it is at a distance a₀ from the nucleus

1 s orbital.

- 30. The maximum number of possible oxidation states of actinoides are shown by
 - (1) Berkelium (Bk) and californium (Cf)
 - (2) Neptunium (Np) and plutonium (Pu)
 - (3) Actinium (Ac) and thorium (Th)
 - (4) Nobelium (No) and lawrencium (Lr)

Answer (2)

Sol. Actinoids	Oxidation state shown
Th	+4
Ac	+3
Pu	+3, +4, +5, +6, +7
Np	+3, +4, +5, +6, +7
Bk	+3, +4
Cm	+3, +4
Lr	+3

... Maximum oxidation state is shown by (Np and Pu)

- Two poles standing on a horizontal ground are 1. of heights 5 m and 10 m respectively. The line joining their tops makes an angle of 15° with the ground. Then the distance (in m) between the poles, is:
 - (1) $5(2+\sqrt{3})$ (2) 10(√<u>3</u>−1)

(3)
$$5(\sqrt{3}+1)$$
 (4) $\frac{5}{2}(2+\sqrt{3})$

Answer (1)

Answer (4)

Sol.



$$\tan 15^\circ = \frac{5}{d} \implies d = \frac{5}{\tan 15^\circ} = \frac{5(\sqrt{3}+1)}{\sqrt{3}-1}$$
$$= \frac{5(4+2\sqrt{3})}{2}$$
$$= 5(2+\sqrt{3})$$

2. If the system of equations 2x + 3y - z = 0, x + ky - 2z = 0 and 2x - y + z = 0 has a

non-trivial solution (x, y, z), then $\frac{x}{y} + \frac{y}{z} + \frac{z}{x} + k$ is equal to:

(1)
$$\frac{1}{2}$$
 (2) -4
(3) $\frac{3}{4}$ (4) -

Answer (1)



∴ Equations are
$$2x + 3y - z = 0$$
 ...(i)
 $2x - y + z = 0$...(ii)
 $2x - y + z = 0$...(iii)
 $2x + 9y - 4z = 0$...(iii)
By (i) – (ii) $2y = z$... $z = -4x$ and $2x + y = 0$
 $\frac{1}{2}$

3. If m is chosen in the quadratic equation $(m^2 + 1) x^2 - 3x + (m^2 + 1)^2 = 0$ such that the sum of its roots is greatest, then the absolute difference of the cubes of its roots is:

(1)	8√3	(2)	10√5

(3) $4\sqrt{3}$ (4) $8\sqrt{5}$

Answer (4)

Sol. Sum of roots = $\frac{3}{m^2 + 1}$

For maximum m = 0

Hence equation becomes $x^2 - 3x + 1 = 0$

$$\alpha + \beta = \mathbf{3}, \quad \alpha\beta = \mathbf{1}, \quad |\alpha - \beta| = \sqrt{5}$$
$$|\alpha^3 - \beta^3| = |(\alpha - \beta)(\alpha^2 + \beta^2 + \alpha\beta)| = \sqrt{5}(\mathbf{9} - \mathbf{1}) = \mathbf{8}\sqrt{5}$$

4. The vertices B and C of a $\triangle ABC$ lie on the line,

 $\frac{x+2}{3} = \frac{y-1}{0} = \frac{z}{4}$ such that BC = 5 units. Then the

area (in sq. units) of this triangle, given that the point A(1, -1, 2), is:

- (1) $5\sqrt{17}$ (2) $\sqrt{34}$
- (3) 6 (4) $2\sqrt{34}$

Answer (2)



Given BC = 5 so we need perpendicular distance of A from line BC.

Let a point D on BC = $(3\lambda - 2, 1, 4\lambda)$

 $\overrightarrow{AD} = (3\lambda - 3)\hat{i} + 2\hat{j} + (4\lambda - 2)\hat{k}$

Also $\overrightarrow{AD} \& \overrightarrow{BC}$ should be perpendicular $\overrightarrow{AD} \cdot \overrightarrow{BC} = 0$ $(3\lambda - 3)3 + 2(0) + (4\lambda - 2)4 = 0$ $9\lambda - 9 + 16\lambda - 8 = 0 \implies \lambda = \frac{17}{25}$ Hence, D = $\left(\frac{1}{25}, 1, \frac{68}{25}\right)$ $|\overline{AD}| = \sqrt{\left(\frac{1}{25} - 1\right)^2 + (2)^2 + \left(\frac{68}{25} - 2\right)^2}$ $= \sqrt{\left(\frac{-24}{25}\right)^2 + 4 + \left(\frac{18}{25}\right)^2}$ $=\sqrt{\frac{(24)^2+4(25)^2+(18)^2}{25^2}}$ $= \sqrt{\frac{576 + 2500 + 324}{25^2}}$ $=\sqrt{\frac{3400}{25^2}}$ $=\frac{\sqrt{34}\cdot 10}{25}=\frac{2\sqrt{34}}{5}$ Area of triangle = $\frac{1}{2} \times |\overrightarrow{BC}| \times |\overrightarrow{AD}|$ $=\frac{1}{2}\times5\times\frac{2\sqrt{34}}{5}=\sqrt{34}$ 5. If $\cos x \frac{dy}{dx} - y \sin x = 6x$, $\left(0 < x < \frac{\pi}{2} \right)$ and $y\left(\frac{\pi}{3}\right) = 0$, then $y\left(\frac{\pi}{6}\right)$ is equal to : (1) $-\frac{\pi^2}{2}$ (2) $-\frac{\pi^2}{4\sqrt{3}}$ (4) $-\frac{\pi^2}{2\sqrt{3}}$ (3) $\frac{\pi^2}{2\sqrt{3}}$ Answer (4)



$$\Rightarrow \int d(y\cos x) = \int 6x \, dx$$

$$\Rightarrow y\cos x = 3x^2 + C$$

As $y\left(\frac{\pi}{3}\right) = 0 \Rightarrow (0) \times \left(\frac{1}{2}\right) = \frac{3\pi^2}{9} + C \Rightarrow C = \frac{-\pi^2}{3}$

$$\Rightarrow y\cos x = 3x^2 - \frac{\pi^2}{3}$$

For $y\left(\frac{\pi}{6}\right)$
 $y\frac{\sqrt{3}}{2} = \frac{3\pi^2}{36} - \frac{\pi^2}{3}$
 $\frac{\sqrt{3}y}{2} = \frac{-3\pi^2}{12} \Rightarrow y = \frac{-\pi^2}{2\sqrt{3}}$

6. If $\int e^{\sec x} (\sec x \tan x f(x) + \sec x \tan x + \sec^2 x)) dx$

$$= e^{secx}f(x) + C$$
, then a possible choice of f(x) is:

(1) $\sec x - \tan x - \frac{1}{2}$ (2) $\sec x + \tan x + \frac{1}{2}$ (3) $\sec x + x \tan x - \frac{1}{2}$ (4) $x \sec x + \tan x + \frac{1}{2}$

Answer (2)

Sol.
$$\int e^{\sec x} (\sec x \tan x f(x) + (\sec x \tan x + \sec^2 x)) dx$$

= $e^{\sec x} f(x) + C$

··· We know that

$$\int e^{g(x)} ((g'(x)f(x)) + f'(x)) dx = e^{g(x)} \times f(x) + C$$

$$\therefore f(x) = \int ((\sec x \tan x) + \sec^2 x) dx$$

$$\therefore f(x) = \sec x + \tan x + C$$

7. If some three consecutive coefficients in the binomial expansion of $(x + 1)^n$ in powers of x are in the ratio 2:15:70, then the average of these three coefficients is:

(1) 625	(2) 964
(3) 232	(4) 227

$$\frac{{}^{n}C_{r-1}}{{}^{n}C_{r}} = \frac{2}{15} & \& \frac{{}^{n}C_{r}}{{}^{n}C_{r+1}} = \frac{15}{70}$$

$$\Rightarrow \frac{r}{n-r+1} = \frac{2}{15} & \& \frac{r+1}{n-r} = \frac{3}{14}$$

$$\Rightarrow 15r = 2n - 2r + 2 & \& 14r + 14 = 3n - 3r$$

$$\Rightarrow 17r = 2n + 2 & \& 17r = 3n - 14$$
i.e., $2n + 2 = 3n - 14 \Rightarrow n = 16 & \& r = 2$
Mean $= \frac{{}^{16}C_{1} + {}^{16}C_{2} + {}^{16}C_{3}}{3} = \frac{16 + 120 + 560}{3}$

$$= \frac{696}{3} = 232$$

8. A rectangle is inscribed in a circle with a diameter lying along the line 3y = x + 7. If the two adjacent vertices of the rectangle are (-8, 5) and (6, 5), then the area of the rectangle (in sq. units) is :

Answer (2)

Sol. Given situation



Perpendicular bisector of AB will pass from centre.

:. Equation of perpendicular bisector x = -1Hence centre (-1, 2)

Let

$$D = (\alpha, \beta) \implies \frac{\alpha + 6}{2} = -1 & \frac{\beta + 5}{2} = 2$$

$$\alpha = -8 & k \beta = -1 \qquad D \equiv (-8, -1)$$

$$|AD| = 6 & |AB| = 14$$

Area = 6 × 14 = 84



 $\hat{\mathbf{j}}$ and $\theta \in (\mathbf{0}, \pi)$ with $\hat{\mathbf{k}}$, then a value of θ is :

(1)	$\frac{5\pi}{12}$	(2)	$\frac{2\pi}{3}$
(3)	$\frac{\pi}{4}$	(4)	$\frac{5\pi}{6}$

Sol. Let $\cos \alpha$, $\cos \beta$, $\cos \gamma$ be direction cosines of \ddot{a} Hence, by given data

$$\cos \alpha = \cos \frac{\pi}{3}, \ \cos \beta = \cos \frac{\pi}{4} \& \cos \gamma = \cos \theta$$

$$\therefore \cos^2 \frac{\pi}{3} + \cos^2 \frac{\pi}{4} + \cos^2 \theta = 1$$
$$\cos^2 \theta = \frac{1}{4} \Rightarrow \cos \theta = \pm \frac{1}{2}, \quad \theta = \frac{\pi}{3} \text{ or } \frac{2\pi}{3}$$

10. If the function
$$f(x) = \begin{cases} a|\pi - x| + 1, & x \le 5 \\ b|x - \pi| + 3, & x > 5 \end{cases}$$
 is

continuous at x = 5, then the value of a - b is:

(1)
$$\frac{2}{\pi-5}$$
 (2) $\frac{-2}{\pi+5}$
(3) $\frac{2}{\pi+5}$ (4) $\frac{2}{5-\pi}$

Answer (4)

Sol. L.H.L
$$\lim_{x\to 5} b|\pi-5|+3=(5-\pi)b+3$$

f(5) = R.H.L.
$$\lim_{x\to 5} a|5-\pi|+1 = a(5-\pi)+1$$

For continuity LHL = RHL

$$(5-\pi)\mathbf{b} + \mathbf{3} = (5-\pi)\mathbf{a} + \mathbf{1}$$
$$\Rightarrow \mathbf{2} = (\mathbf{a} - \mathbf{b})(5-\pi)$$
$$\Rightarrow \mathbf{a} - \mathbf{b} = \frac{\mathbf{2}}{5-\pi}$$

11. Let $z \in C$ be such that |z| < 1. If $\omega = \frac{5+3z}{5(1-z)}$,

then

(1)
$$5 \operatorname{Re}(\omega) > 4$$
 (2) $5 \operatorname{Re}(\omega) > 1$

(3)
$$4 \text{ Im}(\omega) > 5$$
 (4) $5 \text{ Im}(\omega) < 1$

Answer (2)

$$\Rightarrow 5\omega - 5 = z(3 + 5\omega)$$

$$\Rightarrow z = \frac{5(\omega - 1)}{3 + 5\omega}$$
Given $|z| < 1$

$$\Rightarrow 5|\omega - 1| < |3 + 5\omega|$$

$$\Rightarrow 25(\omega\overline{\omega} - \omega - \overline{\omega} + 1) < 9 + 25\omega\overline{\omega} + 15\omega + 15\overline{\omega}$$

$$(using |z|^2 = z\overline{z})$$

$$\Rightarrow 16 < 40\omega + 40\overline{\omega}$$

$$\Rightarrow \quad \omega + \overline{\omega} > \frac{2}{5}$$
$$\Rightarrow \quad 2\operatorname{Re}(\omega) > \frac{2}{5} \quad \Rightarrow \quad \operatorname{Re}(\omega) > \frac{1}{5}$$

12. Let P be the plane, which contains the line of intersection of the planes, x + y + z - 6 = 0 and 2x + 3y + z + 5 = 0 and it is perpendicular to the xy-plane. Then the distance of the point (0, 0, 256) from P is equal to :

(1)
$$63\sqrt{5}$$
 (2) $\frac{17}{\sqrt{5}}$
(3) $205\sqrt{5}$ (4) $\frac{11}{\sqrt{5}}$

Answer (4)

Sol. Let the plane be

$$P = (2x + 3y + z + 5) + \lambda(x + y + z - 6) = 0$$

As the above plane is perpendicular to xy plane

$$\Rightarrow ((2+\lambda)\hat{i} + (3+\lambda)\hat{j} + (1+\lambda)\hat{k}).\hat{k} = 0$$

$$\Rightarrow \lambda = -1$$

$$P = x + 2y + 11 = 0$$

Distance from (0, 0, 256)

$$\left|\frac{0+0+11}{\sqrt{5}}\right| = \frac{11}{\sqrt{5}}$$

- 13. If $f(x) = [x] \left[\frac{x}{4}\right]$, $x \in R$, where [x] denotes the greatest integer function, then :
 - (1) $\lim_{x\to 4^+} f(x)$ exists but $\lim_{x\to 4^-} f(x)$ does not exist
 - (2) f is continuous at x = 4
 - (3) $\lim_{x\to 4^-} f(x)$ exists but $\lim_{x\to 4^+} f(x)$ does not exist
 - (4) Both $\lim_{x\to 4^-} f(x)$ and $\lim_{x\to 4^+} f(x)$ exist but are not equal



Sol. L.H.L
$$\lim_{x \to 4^{-}} [x] - \left\lfloor \frac{x}{4} \right\rfloor = 3 - 0 = 3$$
$$\left(x < 4 \Rightarrow [x] = 3 & \frac{x}{4} < 1 \Rightarrow \left\lfloor \frac{x}{4} \right\rfloor = 0 \right)$$
R.H.L
$$\lim_{x \to 4^{+}} [x] - \left\lfloor \frac{x}{4} \right\rfloor = 4 - 1 = 3$$
$$\left(x > 4 \Rightarrow [x] = 4 & \frac{x}{4} > 1 \Rightarrow \left\lfloor \frac{x}{4} \right\rfloor = 1 \right)$$
f(4) = [4] - $\left\lfloor \frac{4}{4} \right\rfloor = 4 - 1 = 3$

LHL = f(4) = RHL

Hence f(x) is continuous at x = 4

- 14. The common tangent to the circles $x^2 + y^2 = 4$ and $x^{2} + y^{2} + 6x + 8y - 24 = 0$ also passes through the point :
 - (1) (-6, 4) (2) (-4, 6) (4) (6, -2) (3) (4, -2)

Answer (4)

Sol. In given situation $\mathbf{d}_{\mathbf{c}_1\mathbf{c}_2} = |\mathbf{r}_1 - \mathbf{r}_2|$



Common tangent

 $S_1 - S_2 = 0$ $6x + 8y - 20 = 0 \Rightarrow 3x + 4y - 10 = 0$ Hence (6, -2) lies on it

15. The domain of the definition of the function

$$f(x) = \frac{1}{4 - x^2} + \log_{10}(x^3 - x) \text{ is :}$$

$$(1) \ (-1, \ 0) \cup (1, \ 2) \cup (3, \ \infty)$$

$$(2) \ (-1, \ 0) \cup (1, \ 2) \cup (2, \ \infty)$$

$$(3) \ (1, \ 2) \cup (2, \ \infty)$$

$$(4) \ (-2, \ -1) \cup (-1, \ 0) \cup (2, \ \infty)$$
Answer (2)

Hence domain is intersection of (1) & (2) i.e.

$$x \in (-1, 0) \cup (1, 2) \cup (2, \infty)$$

16. The sum of the series $1 + 2 \times 3 + 3 \times 5 + 4 \times 7 + ...$ upto 11th term is :

Answer (2)

Sol. 1 + 2.3 + 3.5 + 4.7 +

Lets break the sequence as shown

We find
$$s_{10} = \sum_{n=1}^{10} (n+1)(2n+1)$$

$$=\sum_{n=1}^{10}(2n^2+3n+1)$$

$$=\frac{2n(n+1)(2n+1)}{6}+\frac{3n(n+1)}{2}+n(n=10)$$

$$=\frac{2.10.11.21}{6}+\frac{3.10.11}{2}+10$$

Hence required sum = 1 + 945 = 946

17. The area (in sq. units) of the smaller of the two circles that touch the parabola, $y^2 = 4x$ at the point (1, 2) and the x-axis is :

(1)
$$4\pi(3+\sqrt{2})$$

(2) $8\pi(2-\sqrt{2})$
(3) $8\pi(3-2\sqrt{2})$
(4) $4\pi(2-\sqrt{2})$

Answer (3)





The circle and parabola will have common tangent at P(1, 2)

... Equation of tangent to parabola

$$\equiv \mathbf{y} \times (\mathbf{2}) = \mathbf{4} \frac{(\mathbf{x} + \mathbf{1})}{\mathbf{2}} \Longrightarrow \mathbf{2}\mathbf{y} = \mathbf{2}\mathbf{x} + \mathbf{2}$$
$$\mathbf{y} = \mathbf{x}$$

Let equation of circle be (using family of circles)

+ 1

$$(x - x_1)^2 + (y - y_1)^2 + \lambda T = 0$$

$$\Rightarrow c \equiv (x - 1)^2 + (y - 2)^2 + \lambda(x - y + 1) = 0$$

Also circle touches x-axis \Rightarrow y-coordinate of centre = radius

$$\Rightarrow c \equiv x^{2} + y^{2} + (\lambda - 2)x + (-\lambda - 4)y + (\lambda + 5) = 0$$

$$\frac{\lambda + 4}{2} = \sqrt{\left(\frac{\lambda - 2}{2}\right)^{2} + \left(\frac{-\lambda - 4}{2}\right)^{2} - (\lambda + 5)}$$

$$\Rightarrow \frac{\lambda^{2} - 4\lambda + 4}{4} = \lambda + 5 \Rightarrow \lambda^{2} - 4\lambda + 4 = 4\lambda + 20$$

$$\Rightarrow \lambda^{2} - 8\lambda - 16 = 0$$

$$\Rightarrow \lambda = \frac{8 \pm \sqrt{64 + 64}}{2}$$

$$= 4 \pm 4\sqrt{2}$$

$$\lambda = 4 - 4\sqrt{2} \text{ (Other value forms bigger circle)}$$
Hence centre of circle $(2\sqrt{2} - 2, 4 - 2\sqrt{2})$

Radius = $4 - 2\sqrt{2}$ Area = $\pi (4 - 2\sqrt{2})^2 = 8\pi (3 - 2\sqrt{2})$

18. The total number of matrices

$$A = \begin{pmatrix} 0 & 2y & 1 \\ 2x & y & -1 \\ 2x & -y & 1 \end{pmatrix}, (x, y \in R, x \neq y) \text{ for }$$

Which $A^T A = 3I_3$ is :

Answer (3)

$$= \begin{bmatrix} 8x^2 & 0 & 0\\ 0 & 6y^2 & 0\\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} 3 & 0 & 0\\ 0 & 3 & 0\\ 0 & 0 & 3 \end{bmatrix}$$

$$\Rightarrow 8x^2 = 3, 6y^2 = 3$$

$$x = \pm \sqrt{\frac{3}{8}}, y = \pm \sqrt{\frac{1}{2}}$$

Total combinations of $(x, y) = 2 \times 2 = 4$

19. If $p \Rightarrow (q \lor r)$ is false, then truth values of p, q, r are respectively :

(1)	T, F, F	(2)	F,	F,	F
(3)	T, T, F	(4)	F,	Т,	Т

Answer (1)

Sol. For
$$p \rightarrow q \lor r$$
 to be F

r must be F & p \rightarrow q must be F for p \rightarrow q to be F p \rightarrow T & q \rightarrow F p, q, r = T, F, F

20. The mean and the median of the following ten numbers in increasing order

10, 22, 26, 29, 34, x, 42, 67, 70, y are 42 and 35 respectively, then $\frac{y}{x}$ is equal to :

(1)
$$\frac{7}{3}$$

(2) $\frac{8}{3}$
(3) $\frac{7}{2}$
(4) $\frac{9}{4}$
Answer (1)
Sol. Mean $= \frac{\sum x_i}{n} = \frac{x + y + 300}{10} = 42 \Rightarrow x + y = 120$
Median $= \frac{T_5 + T_6}{2} = 35 = \frac{34 + x}{2} \Rightarrow x = 36 \& y = 84$
Hence $\frac{y}{x} = \frac{84}{36} = \frac{7}{3}$



consists of one ball, the second row consists of two balls and so on. If 99 more identical balls are added to the total number of balls used in forming the equilateral triangle, then all these balls can be arranged in a square whose each side contains exactly 2 balls less than the number of balls each side of the triangle contains. Then the number of balls used to form the equilateral triangle is :

	(4)	467	(2)	225
۱	(''	137	(2)	223

(3) 262 (4) 190

Answer (4)

Sol. Balls used in equilateral triangle = $\frac{n(n+1)}{2}$

Here side of equilateral triangle has n-balls No. of balls in each side of square is = (n - 2)

Given
$$\frac{n(n+1)}{2} + 99 = (n-2)^2$$

 $\Rightarrow n^2 + n + 198 = 2n^2 - 8n + 8$
 $\Rightarrow n^2 - 9n - 190 = 0$
 $\Rightarrow n^2 - 19n + 10n - 190 = 0$
 $\Rightarrow (n - 19) (n + 10) = 0$
 $\Rightarrow n = 19$

Balls used to form triangle

$$=\frac{n(n+1)}{2}=\frac{19\times 20}{2}=190$$

- 22. Two newspapers A and B are published in a city. It is known that 25% of the city population reads A and 20% reads B while 8% reads both A and B. Further, 30% of those who read A but not B look into advertisements and 40% of those who read B but not A also look into advertisements, while 50% of those who read both A and B look into advertisements. Then the percentage of the population who look into advertisements is :
 - (1) 13.9
 - (2) 13
 - (3) 12.8
 - (4) 13.5



n(A only) = 25 - 8 = 17% n(B only) = 20 - 8 = 12%

% of people from A only who read advertisement = $17 \times 0.3 = 5.1\%$

% of people from B only who read advertisement = 12 × 0.4 = 4.8%

% of people from A&B both who read advertisement = $8 \times 0.5 = 4\%$

Total % of people who read advertisement

= 5.1 **+** 4.8 **+** 4 **=** 13.9%

23. A water tank has the shape of an inverted right circular cone, whose semi-vertical angle is

 $\tan^{-1}\left(\frac{1}{2}\right)$. Water is poured into it at a constant

rate of 5 cubic metre per minute. Then the rate (in m/min.), at which the level of water is rising at the instant when the depth of water in the tank is 10 m; is :

(1)
$$\frac{2}{\pi}$$
 (2) $\frac{1}{15\pi}$
(3) $\frac{1}{10\pi}$ (4) $\frac{1}{5\pi}$

Answer (4)



 $V = \frac{1}{3}\pi r^2 h$...(i) (where r is radius and h is height at any time)



Differentiate eqⁿ. (i), we get

$$\frac{dV}{dt} = \frac{1}{3} \left(\pi 2r \frac{dr}{dt} h + \pi r^2 \frac{dh}{dt} \right) = \frac{1}{3} \left(100\pi \frac{1}{2} + 25\pi \right) \frac{dh}{dt}$$

at h = 10, r = 5
$$5 = \frac{75\pi}{3} \frac{dh}{dt}$$
$$\Rightarrow \frac{dh}{dt} = \frac{1}{5\pi} m / mim$$

24. The value of the integral

$$\int_{0}^{1} x \cot^{-1}(1-x^{2}+x^{4}) dx \text{ is}$$
(1) $\frac{\pi}{4} - \log_{e} 2$
(2) $\frac{\pi}{2} - \log_{e} 2$
(3) $\frac{\pi}{2} - \frac{1}{2}\log_{e} 2$
(4) $\frac{\pi}{4} - \frac{1}{2}\log_{e} 2$

Answer (4)

Sol.
$$\int_{0}^{1} x \cot^{-1} (1 - x^{2} + x^{4}) dx = \int_{0}^{1} x \tan^{-1} \left(\frac{1}{1 + x^{4} - x^{2}} \right)$$

$$\Rightarrow \int_{0}^{1} x \tan^{-1} \left(\frac{x^{2} - (x^{2} - 1)}{1 + x^{2} (x^{2} - 1)} \right) dx$$

$$\Rightarrow \int_{0}^{1} x \tan^{-1} x^{2} dx - \int_{0}^{1} x \tan^{-1} (x^{2} - 1) dx$$

Put $x^{2} = t \Rightarrow 2x dx = dt$, (For 1st integration)
put $x^{2} - 1 = k \Rightarrow 2x dx = dk$ (For 2nd integration)

$$\Rightarrow \frac{1}{2} \int_{0}^{1} 1 \tan^{-1} t dt - \frac{1}{2} \int_{-1}^{0} 1 \tan^{-1} k dk$$

$$\Rightarrow \frac{1}{2} \left(\int_{0}^{1} t \tan^{-1} t \int_{0}^{1} - \int_{0}^{1} \frac{t}{1 + t^{2}} dt \right) - \frac{1}{2}$$

$$\left(k \tan^{-1} k \int_{-1}^{0} - \int_{-1}^{0} \frac{k}{1 + k^{2}} dk \right)$$

$$2\left(4\left(2\left(-\frac{\pi}{4}\right)_{0}^{2}\right) + 2\right)$$

$$\left(0 - \frac{\pi}{4} - \left(\frac{1}{2}\ln(1+k^{2})\right)_{-1}^{0}\right)$$

$$\Rightarrow \left(\frac{\pi}{8} - \frac{1}{4}\ln^{2}\right) - \left(\frac{-\pi}{8} - \frac{1}{4}\ln^{2}\right)$$

$$\Rightarrow \frac{\pi}{4} - \frac{1}{2}\ln^{2}$$

25. The area (in sq. units) of the region

$$A = \left\{ (x, y) : \frac{y^2}{2} \le x \le y + 4 \right\} \text{ is}$$

(1) 18
(2) 16
(3) $\frac{53}{3}$
(4) 30

Answer (1)



Hence area =
$$\int_{-2}^{4} x dy$$

$$=\int_{-2}^{4}\left(y+4-\frac{y^{2}}{2}\right)dy$$

$$=\frac{y^{2}}{2}+4y-\frac{y^{3}}{6}\int_{-2}^{4}=\left(8+16-\frac{64}{6}\right)-\left(2-8+\frac{8}{6}\right)$$
$$=\left(24-\frac{32}{3}\right)-\left(-6+\frac{4}{3}\right)=\frac{40}{3}+\frac{14}{3}=\frac{54}{3}=18$$



of their point of intersection from the origin is

(1)
$$\sqrt{\frac{2}{5}}$$
 (2) $\frac{\sqrt{2}}{5}$
(3) $\frac{2}{5}$ (4) $\frac{2}{\sqrt{5}}$

Answer (1)

Sol. For perpendicular $m_1m_2 = -1$

$$\left(\frac{-1}{a-1}\right)\left(\frac{-2}{a^2}\right) = -1$$

$$\Rightarrow 2 = a^2 (1-a)$$

$$\Rightarrow a^3 - a^2 + 2 = 0$$

$$\Rightarrow (a+1) (a^2 + 2a + 2) = \frac{1}{a = -1}$$

Hence lines are x - 2y = 1 and 2x + y = 1

0

 $\therefore \quad \text{Intersection point}\left(\frac{3}{5},\frac{-1}{5}\right)$

Distance from origin $=\sqrt{\frac{9}{25} + \frac{1}{25}} = \sqrt{\frac{10}{25}} = \sqrt{\frac{2}{5}}$

27. The value of sin10° sin30° sin50° sin70° is

(1)
$$\frac{1}{18}$$
 (2) $\frac{1}{32}$
(3) $\frac{1}{16}$ (4) $\frac{1}{36}$

Answer (3)

Sol. $\sin(60^\circ + A)$. $\sin(60^\circ - A) \sin A = \frac{1}{4} \sin 3A$

Hence, sin10° sin50° sin70°

= sin10° sin(60° – 10°) sin(60° + 10°) =
$$\frac{1}{4}$$
 sin30°

Hence, sin10° sin30° sin50° sin70°
$$=\frac{1}{4}$$
 sin²30°

$$=\frac{1}{16}$$

 x^2 + $2y^2$ = 1, then α is equal to

(1) $\sqrt{2} - 1$ (2) $\sqrt{2} + 1$ (3) $2\sqrt{2} + 1$ (4) $2\sqrt{2} - 1$

Answer (2)

Sol. Let tangent in terms of m to parabola and ellipse

i.e
$$y = mx + \frac{1}{4m}$$
 for parabola at point $\left(\frac{1}{4m^2}, \frac{-1}{2m}\right)^2$
and $y = mx \pm \sqrt{m^2 + \frac{1}{2}}$ for ellipse on comparing
 $\Rightarrow \frac{1}{4m} = \pm \sqrt{m^2 + \frac{1}{2}} \Rightarrow \frac{1}{16m^2} = m^2 + \frac{1}{2}$
 $\Rightarrow 16m^4 + 8m^2 - 1 = 0$
 $m^2 = \frac{-8 \pm \sqrt{64 + 64}}{2(16)} = \frac{-8 \pm 8\sqrt{2}}{2(16)} = \frac{\sqrt{2} - 1}{4}$
 $\alpha = \frac{1}{4m^2} = \frac{1}{4\frac{\sqrt{2} - 1}{4}} = \sqrt{2} + 1$

29. If $f: R \rightarrow R$ is a differentiable function and

f(2) = 6, then
$$\lim_{x \to 2} \int_{6}^{f(x)} \frac{2t \, dt}{(x-2)}$$
 is
(1) 0 (2) $2f'(2)$

(a)
$$12f'(2)$$
 (b) $(2f'(2))$ (c) $(2f'(2))$

Answer (3)

Sol. Using L' Hospital rule and Leibnitz theorem

$$\lim_{x \to 2} \frac{\int_{6}^{f(x)} 2t dt}{(x-2)}$$
$$\lim_{x \to 2} \frac{2f(x)f'(x) - 0}{1}$$
$$2f(2)f'(2) = 12f'(2)$$



value of its 11th term is $T_1 = 7$ (1) -36 (2) 25 $T_2 = 11$ \Rightarrow $T_{11} = T_1 + 10d = 7 + 10(4) = 47$ (3) -25 (4) -35 $T_{3} = 15$ Answer (3) Sol. Let terms be a – d, a, a + d if d = – 4 \Rightarrow 3a = 33 \Rightarrow 11 $T_1 = 15$ **Product of terms** (a – d) a (a + d) = 11 (121 – d²) = 1155 $T_{3} = 7$ \Rightarrow 121 - d² = 105 \Rightarrow d = ± 4



Answers & Solutions

for

JEE (MAIN)-2019 (Online) Phase-2

(Physics, Chemistry and Mathematics)

Time : 3 hrs.

M.M.: 360

Important Instructions :

- 1. The test is of **3 hours** duration.
- 2. The Test Booklet consists of **90** questions. The maximum marks are **360**.
- 3. There are *three* parts in the question paper A, B, C consisting of **Physics**, **Chemistry** and **Mathematics** having 30 questions in each part of equal weightage.
- 4. Each question is allotted 4 (four) marks for each correct response. ¼ (one-fourth) marks will be deducted for indicating incorrect response of each question. No deduction from the total score will be made if no response is indicated for an item in the answer sheet.
- 5. There is only one correct response for each question.



1. Two coaxial discs, having moments of inertia I_1 and $\frac{I_1}{2}$, are rotating with respective angular

velocities ω_1 and $\frac{\omega_1}{2}$ about their common axis. They are brought in contact with each other and thereafter they rotate with a common angular velocity. If E_f and E_i are the final and initial total energies, then $(E_f - E_i)$ is :

(1)
$$-\frac{l_1\omega_1^2}{12}$$
 (2) $\frac{3}{8}l_1\omega_1^2$
(3) $\frac{l_1\omega_1^2}{6}$ (4) $-\frac{l_1\omega_1^2}{24}$

Answer (4)

Sol. By applying conservation of angular momentum

$$(I_{1} + I_{2}) \omega_{common} = I_{1}\omega_{1} + I_{2}\omega_{2}$$

$$\omega_{common} = \frac{I_{1}\omega_{1} + \frac{I_{1}\omega_{1}}{4}}{I_{1} + \frac{I_{1}}{2}} = \left(\frac{5}{4} \times \frac{2}{3}\right)\omega_{1}$$

$$\omega_{c} = \frac{5\omega_{1}}{6}$$

$$\therefore \text{ Loss in KE} = \left(\frac{1}{2}I_{1}\omega_{1}^{2} + \frac{1}{2}I_{2}\omega_{2}^{2}\right) - \frac{1}{2}(I_{1} + I_{2})\omega_{c}^{2}$$

$$\therefore \Delta KE = -\frac{I_{1}\omega_{1}^{2}}{24}$$

 A ray of light AO in vacuum is incident on a glass slab at angle 60° and refracted at angle 30° along OB as shown in the figure. The optical path length of light ray from A to B is :



Answer (4)

Sol. From the given figure



3. A ball is thrown upward with an initial velocity V_0 from the surface of the earth. The motion of the ball is affected by a drag force equal to $m\gamma v^2$ (where m is mass of the ball, v is its instantaneous velocity and γ is a constant). Time taken by the ball to rise to its zenith is :

(1)
$$\frac{1}{\sqrt{\gamma g}} \sin^{-1} \left(\sqrt{\frac{\gamma}{g}} V_0 \right)$$

(2)
$$\frac{1}{\sqrt{\gamma g}} \ln \left(1 + \sqrt{\frac{\gamma}{g}} V_0 \right)$$

(3)
$$\frac{1}{\sqrt{\gamma g}} \tan^{-1} \left(\sqrt{\frac{\gamma}{g}} V_0 \right)$$

(4)
$$\frac{1}{\sqrt{2\gamma g}} \tan^{-1} \left(\sqrt{\frac{2\gamma}{g}} V_0 \right)$$

Answer (3)

Sol. Retardation of the particle

$$a = -(g + \gamma v^{2})$$

$$\int_{v_{0}}^{0} \frac{-dv}{g + \gamma v^{2}} = \int_{0}^{t} dt \qquad [for H_{max} v = 0]$$

$$\frac{1}{\sqrt{\gamma g}} \tan^{-1} \left(\frac{\sqrt{\gamma} v_{0}}{\sqrt{g}}\right) = t$$



 $\mathbf{x} = \mathbf{x}_0 + \mathbf{a} \cos \omega_1 \mathbf{t}$

 $y = y_0 + b \sin \omega_2 t$

The torque, acting on the particle about the origin, at t = 0 is :

(1)
$$-m(x_0b\omega_2^2-y_0a\omega_1^2)\hat{k}$$

- (2) $m(-x_0b+y_0a)\omega_1^2\hat{k}$
- (3) $+ my_0 a \omega_1^2 \hat{k}$
- (4) Zero

Answer (3)

Sol.
$$x = x_0 + a \cos \omega_1 t$$

 $y = y_0 + b \sin \omega_2 t$
 $\Rightarrow v_x = -a\omega_1 \sin (\omega_1 t), \quad v_y = b\omega_2 \cos (\omega_2 t)$
 $a_x = -a\omega_1^2 \cos(\omega_1 t), \quad a_y = -b\omega_2^2 \sin(\omega_2 t)$
At $t = 0, \quad x = x_0 + a, \quad y = y_0$
 $a_x = -a\omega_1^2, \quad a_y = 0$
 $\therefore \quad \vec{\tau} = m(-a\omega_1^2) \times y_0(-\hat{k})$
 $= + my_0 a\omega_1^2 \hat{k}.$

5. In a photoelectric effect experiment the threshold wavelength of light is 380 nm. If the wavelength of incident light is 260 nm, the maximum kinetic energy of emitted electrons will be :

Given E (in eV) =
$$\frac{1237}{\lambda(\text{in nm})}$$

(1) 4.5 eV (2) 15.1 eV
(3) 3.0 eV (4) 1.5 eV

Answer (4)

Sol. Wavelength of incident wave (λ) = 260 nm

Cut off (threshold) wavelength (λ_0) = 380 nm

Then
$$KE_{max} = \frac{hc}{\lambda} - \frac{hc}{\lambda_0}$$

= $1237 \left[\frac{1}{260} - \frac{1}{380} \right]$
= $\frac{1237 \times 120}{380 \times 260} = 1.5 \text{ eV}$

range). As shown in the figure, it is a straight line.



One may conclude that :

(1)
$$R(T) = \frac{R_0}{T^2}$$

(2) $R(T) = R_0 e^{-T_0^2/T^2}$
(3) $R(T) = R_0 e^{-T^2/T_0^2}$
(4) $R(T) = R_0 e^{T^2/T_0^2}$
Answer (2)

Sol.
$$\ln R(T) = a - \frac{a}{b} - \frac{1}{T^2}$$

a, b are constant

$$\mathbf{R}(\mathbf{T}) = \mathbf{R}_0 \mathbf{e}^{\frac{-\mathbf{T}_0^2}{\mathbf{T}^2}}$$

- 7. An npn transistor operates as a common emitter amplifier, with a power gain of 60 dB. The input circuit resistance is 100 Ω and the output load resistance is 10 k Ω . The common emitter current gain β is :
 - (1) 10⁴
 - (2) 6×10^2
 - (3) 10²
 - (4) 60

Answer (3)

Sol.
$$P_{gain} = \beta^2 \left(\frac{R_{out}}{R_{in}} \right) \& I_{gain} = \beta$$

$$\therefore \quad 10^6 = \beta^2 \left(\frac{10000}{100} \right)$$

$$\beta = 100$$



between them is d. A third wire C carrying a current I is to be kept parallel to them at a distance x from A such that the net force acting on it is zero. The possible values of x are :



Answer (1)

Sol. $\Sigma \vec{F} = 0$

$$\frac{\mu_0 I_1}{2\pi x} = \frac{\mu_0 I_2}{2\pi (x-d)} \quad \text{since } (x > d)$$

$$I_1 x - I_1 d = I_2 x$$

$$x = \frac{I_1 d}{I_1 - I_2}$$

9. A stationary source emits sound waves of frequency 500 Hz. Two observers moving along a line passing through the source detect sound to be of frequencies 480 Hz and 530 Hz. Their respective speeds are, in ms⁻¹,

(Given speed of sound = 300 m/s)

(1)	12, 16	(2)	16,	14
(3)	8, 18	(4)	12,	18

Answer (4)

Sol. Frequency of sound source $(f_0) = 500 \text{ Hz}$

When observer is moving away from the source

Apparent frequency
$$f_1 = 480 = f_0 \left(\frac{v - v'_0}{v} \right) \dots (i)$$

And when observer is moving towards the

source
$$f_2 = 530 = f_0 \left(\frac{\nu + \nu'_0}{\nu} \right)$$
 ...(ii)

$$480 = 500 \left(\frac{300 - v'_0}{300} \right)$$

 $v'_0 = 12 \text{ m/s}$
From equation (ii)

$$530 = 500 \left(1 + \frac{\nu_0''}{\nu}\right)$$

 $\therefore v_0'' = 18 \text{ m/s}$

- 10. A transformer consisting of 300 turns in the primary and 150 turns in the secondary gives output power of 2.2 kW. If the current in the secondary coil is 10 A, then the input voltage and current in the primary coil are :
 - (1) 440 V and 5 A (2) 220 V and 20 A
 - (3) 220 V and 10 A (4) 440 V and 20 A

Answer (1)

...

Sol. Power output $(V_2I_2) = 2.2 \text{ kW}$

$$V_2 = \frac{2.2 \text{ kW}}{(10\text{ A})} = 220 \text{ volts}$$

... Input voltage for step-down transformer

$$\frac{V_1}{V_2} = \frac{N_1}{N_2} = 2$$
$$V_{input} = 2 \times V_{output} = 2 \times 220$$
$$= 440 \text{ V}$$

Also
$$\frac{I_1}{I_2} = \frac{N_2}{N_1}$$

:
$$I_1 = \frac{1}{2} \times 10 = 5 \text{ A}$$

A uniformly charged ring of radius 3a and total charge q is placed in xy-plane centred at origin. A point charge q is moving towards the ring along the z-axis and has speed v at z = 4a. The minimum value of v such that it crosses the origin is :

(1)
$$\sqrt{\frac{2}{m}} \left(\frac{1}{15} \frac{q^2}{4\pi\epsilon_0 a}\right)^{1/2}$$
 (2) $\sqrt{\frac{2}{m}} \left(\frac{1}{5} \frac{q^2}{4\pi\epsilon_0 a}\right)^{1/2}$
(3) $\sqrt{\frac{2}{m}} \left(\frac{4}{15} \frac{q^2}{4\pi\epsilon_0 a}\right)^{1/2}$ (4) $\sqrt{\frac{2}{m}} \left(\frac{2}{15} \frac{q^2}{4\pi\epsilon_0 a}\right)^{1/2}$





Potential at any point of the charged ring

$$V_{P} = \frac{KQ}{\sqrt{R^{2} + x^{2}}}$$

The minimum velocity (v_0) should just sufficient to reach the point charge at the center, therefore

$$\frac{1}{2}mv_0^2 = Q[V_C - V_P]$$
$$= Q\left[\frac{KQ}{3a} - \frac{KQ}{5a}\right]$$
$$v_0^2 = \frac{4KQ^2}{15 \text{ ma}} = \frac{4}{15}\frac{1}{4\pi\epsilon_0}\frac{q^2}{\text{ ma}}$$
$$\therefore \quad v_0 = \sqrt{\frac{2}{m}}\left(\frac{2q^2}{15 \times 4\pi\epsilon_0a}\right)^{\frac{1}{2}}$$

12. n moles of an ideal gas with constant volume heat capacity C_v undergo an isobaric expansion by certain volume. The ratio of the work done in the process, to the heat supplied is :

(1)
$$\frac{4nR}{C_V + nR}$$
 (2) $\frac{nR}{C_V + nR}$
(3) $\frac{4nR}{C_V - nR}$ (4) $\frac{nR}{C_V - nR}$

Answer (Bonus)

Sol. For Isobaric process Work done (W) = $nR\Delta T$ and Heat given (Q) = $nC_{P}\Delta T$

$$\therefore \quad \frac{\mathsf{W}}{\mathsf{Q}} = \frac{\mathsf{R}}{\mathsf{C}_{\mathsf{P}}} = \frac{\mathsf{R}}{\mathsf{C}_{\mathsf{V}} + \mathsf{R}}$$

13. A 25 × 10^{-3} m³ volume cylinder is filled with 1 mol of O_2 gas at room temperature (300 K). The molecular diameter of O_2 , and its root mean square speed, are found to be 0.3 nm and 200 m/s, respectively. What is the average collision rate (per second) for an O_2 molecule?

(1)	~10 ¹²	(2)	~10 ¹⁰
(3)	~10 ¹¹	(4)	~10 ¹³

JUI. $v = 23 \times 10^{-111}$, $N = 1^{-111010}$ of O_2

T = 300 K
V_{rms} = 200 m/s

$$\therefore \lambda = \frac{1}{2}$$

$$\sqrt{2N\pi r^2}$$

Average time
$$\frac{1}{\tau} = \frac{\langle \mathbf{v} \rangle}{\lambda} = 200 \cdot N\pi r^2 \cdot \sqrt{2}$$

$$=\frac{\sqrt{2}\times200\times6.023\times10^{23}}{25\times10^{-3}}\cdot\pi\times10^{-18}\times0.09$$

Average no. of collision $\approx 10^{10}$

14. Two radioactive materials A and B have decay constants 10 λ and λ , respectively. If initially they have the same number of nuclei, then the ratio of the number of nuclei of A to that of B will be 1/e after a time :

(1)
$$\frac{1}{10\lambda}$$
 (2) $\frac{11}{10\lambda}$
(3) $\frac{1}{9\lambda}$ (4) $\frac{1}{11\lambda}$

Answer (3)

Sol. Number of nuclei present at any time t

$$N = N_0 e^{-\lambda t}$$

$$\therefore \quad \frac{N_A}{N_B} = e^{(\lambda_B - \lambda_A)t} = \frac{1}{e}$$

$$(\lambda_A - \lambda_B) \cdot t = 1$$

$$\therefore \quad t = \frac{1}{-\lambda + 10\lambda} = \frac{1}{9\lambda}$$

Figure shows charge (q) versus voltage (V) 15. graph for series and parallel combination of two given capacitors. The capacitances are :





con Equivalent capacitance for series combination

$$\mathbf{C}' = \frac{\mathbf{C}_1 \mathbf{C}_2}{\mathbf{C}_1 + \mathbf{C}_2}$$

For parallel combination $C'' = C_1 + C_2$

$$C_1 + C_2 = \frac{500}{10} = 50 \ \mu\text{F}$$

and
$$\frac{C_1C_2}{C_1+C_2} = \frac{80}{10} = 8\,\mu\text{F}$$

∴ C₁C₂ = 400 μF

Solving C₁ = 40 μ F C₂ = 10 μ F

16. A message signal of frequency 100 MHz and peak voltage 100 V is used to execute amplitude modulation on a carrier wave of frequency 300 GHz and peak voltage 400 V. The modulation index and difference between the two side band frequencies are :

(1) 4;
$$2 \times 10^8$$
 Hz (2) 4; 1×10^8 Hz

(3) 0.25;
$$1 \times 10^8$$
 Hz (4) 0.25; 2×10^8 Hz

Answer (4)

- **Sol.** Range of frequency = $(f_c f_m)$ to $(f_c + f_m)$
 - $\therefore \text{ Band width} = 2f_m = 2 \times 100 \times 10^6 \text{ Hz}$ $= 2 \times 10^8 \text{ Hz}$

and Modulation index = $\frac{A_m}{A_c} = \frac{100}{400} = 0.25$

17. Two particles, of masses M and 2M, moving, as shown, with speeds of 10 m/s and 5 m/s, collide elastically at the origin. After the collision, they move along the indicated directions with speeds v_1 and v_2 , respectively. The values of v_1 and v_2 are nearly :



- (3) 6.5 m/s and 3.2 m/s
- (4) 6.5 m/s and 6.3 m/s

X and Y direction for the system then

 $M(10\cos 30^\circ) + 2M(5\cos 45^\circ) = 2M (v_1\cos 30^\circ)$

+ M(v₂cos45°)

$$5\sqrt{3} + 5\sqrt{2} = \sqrt{3} v_1 + \frac{v_2}{\sqrt{2}}$$
 ...(1)

Also

2M(5sin45°) - M(10sin30°) = 2Mv₁sin30°

- Mv₂sin45°

$$5\sqrt{2} - 5 = v_1 - \frac{v_2}{\sqrt{2}} \qquad \dots (2)$$

Solving equation (1 and 2)

$$(\sqrt{3}+1)v_1 = 5\sqrt{3}+10\sqrt{2}-5 \implies v_1 = 6.5 \text{ m/s}$$

 $v_2 = 6.3 \text{ m/s}$

- 18. A moving coil galvanometer allows a full scale current of 10^{-4} A. A series resistance of 2 M Ω is required to convert the above galvanometer into a voltmeter of range 0 5 V. Therefore the value of shunt resistance required to convert the above galvanometer into an ammeter of range 0–10 mA is :
 - **(1) 200** Ω
 - **(2) 500** Ω
 - **(3) 100** Ω
 - **(4) 10** Ω

Answer (Bonus)

- Sol. Data contradictory
- 19. In the given circuit, an ideal voltmeter connected across the 10 Ω resistance reads 2 V. The internal resistance r, of each cell is :





Sol. For the given circuit

$$15 \Omega$$

$$2 \Omega$$

$$A 10 \Omega B$$

$$r$$

$$r$$

$$1.5 V$$

$$1.5 V$$
Given that $V = 2 V$

Given that $V_{AB} = 2 V$

$$\therefore \quad \mathbf{I} = \frac{2}{15} + \frac{2}{10} = \frac{1}{3} \mathbf{A}$$

Also I(2r + 2) = 1.5 + 1.5 - V_{AB}

⇒ 2r + 2 = (3–2)3

- \Rightarrow $\mathbf{r} = \frac{1}{2} \Omega$
- 20. The value of acceleration due to gravity at Earth's surface is 9.8 ms⁻². The altitude above its surface at which the acceleration due to gravity decreases to 4.9 ms⁻², is close to : (Radius of earth = 6.4×10^6 m)
 - (1) 9.0×10^6 m (2) 6.4×10^6 m (3) 1.6×10^6 m (4) 2.6×10^6 m

Answer (4)

Sol. Given

$$g_{height} = \frac{g_{surface}}{2} = 4.9 \text{ m/s}^2$$
As
$$g_h = g \left(1 + \frac{h}{R_e} \right)^{-2}$$

$$h = R_e \left(\sqrt{2} - 1 \right)$$

$$h = 6400 \times 0.414$$

$$h = 2649.6 \text{ km}$$

$$h = 2.6 \times 10^6 \text{ m}$$

21. The ratio of surface tensions of mercury and water is given to be 7.5 while the ratio of their densities is 13.6. Their contact angles, with glass, are close to 135° and 0°, respectively. It is observed that mercury gets depressed by an amount h in a capillary tube of radius r_1 , while water rises by the same amount h in a capillary tube of radius r_2 . The ratio, (r_1/r_2) , is then close to :

(1) 4/5	(2) 2/3
(3) 3/5	(4) 2/5

our natio or surface tension

$$\frac{S_{Hg}}{S_{Water}} = 7.5$$

$$\frac{\rho_{Hg}}{\rho_w} = 13.6 \& \frac{\cos \theta_{Hg}}{\cos \theta_W} = \frac{\cos 135^\circ}{\cos 0^\circ} = \frac{1}{\sqrt{2}}$$

$$\frac{R_{Hg}}{R_{Water}} = \left(\frac{S_{Hg}}{S_W}\right) \left(\frac{\rho_W}{\rho_{Hg}}\right) \left(\frac{\cos \theta_{Hg}}{\cos \theta_W}\right)$$

$$= 7.5 \times \frac{1}{13.6} \times \frac{1}{\sqrt{2}} = 0.4 = \frac{2}{5}$$

22. A thin disc of mass M and radius R has mass per unit area $\sigma(r) = kr^2$ where r is the distance from its centre. Its moment of inertia about an axis going through its centre of mass and perpendicular to its plane is :

(1)
$$\frac{MR^2}{3}$$
 (2) $\frac{MR^2}{6}$
(3) $\frac{MR^2}{2}$ (4) $\frac{2MR^2}{3}$

Answer (4)

Sol. Surface mass density (σ) = kr²

Mass of disc
$$M = \int_{0}^{R} (kr^2) 2\pi r dr$$

$$=2\pi k\frac{R^4}{4}=\frac{\pi kR^4}{2}$$

... Moment of inertia about the axis of the disc.

$$I = \int dI = \int (dm)r^2 = \int \sigma dAr^2$$
$$= \int (Kr^2)(2\pi rdr)r^2$$
$$= \int_0^R 2\pi k r^5 dr = \frac{\pi kR^6}{3} = \frac{2}{3}MR^2$$

- 23. A cylinder with fixed capacity of 67.2 lit contains helium gas at STP. The amount of heat needed to raise the temperature of the gas by 20° C is : [Given that R = 8.31 J mol⁻¹K⁻¹]
 - (1) 700 J
 - (2) 350 J
 - (3) 374 J
 - (4) 748 J



Sol. No. of moles of He at STP $=\frac{67.2}{22.4}=3$

As the volume is constant \rightarrow Isochoric proces

$$\mathbf{Q} = \mathbf{nC}_{\mathbf{v}} \Delta \mathbf{T} = \mathbf{3} \times \frac{\mathbf{3R}}{2} \times \mathbf{20} = \mathbf{90R} = \mathbf{90} \times \mathbf{8.31} \simeq \mathbf{748} \text{ J}.$$

24. A proton, an electron, and a Helium nucleus, have the same energy. They are in circular orbits in a plane due to magnetic field perpendicular to the plane. Let r_p , r_e and r_{He} be their respective radii, then,

(1)
$$r_e < r_p < r_{He}$$
 (2) $r_e > r_p = r_{He}$
(3) $r_e < r_p = r_{He}$ (4) $r_e > r_p > r_{He}$

Answer (3)

Sol. Radius of circular path (r) in a perpendicular

uniform magnetic field
$$=\frac{mv}{qB}=\frac{\sqrt{2mK}}{qB}$$

For proton, electron and α -particle,

$$m_{\alpha} = 4m_{p}$$
 and $m_{p} >> m_{e}$

Also $q_{\alpha} = 2q_{p}$ and $q_{p} = q_{e}$

 \therefore As KE of all the particles is same then,

$$\mathbf{r} \propto \frac{\sqrt{\mathbf{m}}}{\mathbf{q}}$$

- \therefore $\mathbf{r}_{\alpha} = \mathbf{r}_{p} > \mathbf{r}_{e}$
- 25. One plano-convex and one plano-concave lens of same radius of curvature 'R' but of different materials are joined side by side as shown in the figure. If the refractive index of the material of 1 is μ_1 and that of 2 is μ_2 , then the focal length of the combination is :



$$\mathbf{f_1} = \frac{\mathbf{R}}{(\boldsymbol{\mu_1} - \mathbf{1})}$$

Focal length of plano concave lens-

$$\boldsymbol{f_2} = \frac{-\boldsymbol{R}}{\left(\boldsymbol{\mu_2} - \boldsymbol{1}\right)}$$

For the combination of two lens-

$$\frac{1}{f_{eq}} = \frac{1}{f_1} + \frac{1}{f_2} = \frac{\mu_1 - 1}{R} - \frac{\mu_2 - 1}{R}$$
$$= \frac{\mu_1 - \mu_2}{R}$$
$$\therefore \quad f_{eq} = \frac{R}{\mu_1 - \mu_2}$$

26. The electric field of a plane electromagnetic wave is given by

$$\vec{\mathbf{E}} = \mathbf{E}_{\mathbf{0}}\hat{\mathbf{i}}\cos(\mathbf{kz})\cos(\mathbf{\omega t})$$

The corresponding magnetic field $\vec{\textbf{B}}$ is then given by :

(1)
$$\vec{B} = \frac{E_0}{C}\hat{j} \sin(kz)\cos(\omega t)$$

(2) $\vec{B} = \frac{E_0}{C}\hat{j}\cos(kz)\sin(\omega t)$
(3) $\vec{B} = \frac{E_0}{C}\hat{j}\sin(kz)\sin(\omega t)$

(4)
$$\vec{B} = \frac{E_0}{C}\hat{k} \sin(kz)\cos(\omega t)$$

Answer (3)

Sol.
$$\frac{\mathbf{E}_0}{\mathbf{B}_0} = \mathbf{C}$$

 $\therefore \quad \mathbf{B}_0 = \frac{\mathbf{E}_0}{\mathbf{C}}$

Given that $\vec{E} = E_0 \cos(kz)\cos(\omega t)\hat{i}$

$$\vec{\mathbf{E}} = \frac{\mathbf{E}_0}{2} \left[\cos(\mathbf{k}\mathbf{z} - \omega \mathbf{t}) \,\hat{\mathbf{i}} - \cos(\mathbf{k}\mathbf{z} + \omega \mathbf{t}) \,\hat{\mathbf{i}} \right]$$

Correspondingly

$$\vec{B} = \frac{B_0}{2} \Big[\cos(kz - \omega t) \hat{j} - \cos(kz + \omega t) \hat{j} \Big]$$
$$\vec{B} = \frac{B_0}{2} \times 2 \sin kz \sin \omega t$$
$$\vec{B} = \Big(\frac{E_0}{C} \sin kz \sin \omega t \Big) \hat{j}$$



table are shown in figure.



Which of the readings is inconsistent ?

(1) 3	(2) 2
(3) 1	(4) 4

Answer (4)

Sol.
$$\frac{R}{X} = \frac{I}{100-I}$$

Using the above expression

$$\mathbf{X} = \frac{\mathbf{R}(100 - \mathbf{I})}{\mathbf{I}}$$

for case (a) $x = \frac{100 \times 40}{60} = \frac{2000}{3} \Omega$

for case (b)
$$x = \frac{100 \times 87}{13} = \frac{8700}{13} \Omega$$

for case (c)
$$x = \frac{10 \times 98.5}{1.5} = \frac{1970}{3} \Omega$$

for case (d)
$$x = \frac{1 \times 99}{1} = 99 \Omega$$

Clearly we can see that the value of x calculate in case (d) is inconsistent than other cases.

- 28. A current of 5 A passes through a copper conductor (resistivity = $1.7 \times 10^{-8} \Omega$ m) of radius of cross-section 5 mm. Find the mobility of the charges if their drift velocity is 1.1×10^{-3} m/s.
 - (1) 1.3 m²/Vs
 - (2) 1.8 m²/Vs
 - (3) 1.5 m²/Vs
 - (4) $1.0 \text{ m}^2/\text{Vs}$

Sol. Mobility
$$(\mu) = \frac{\mathbf{v}_{d}}{\mathbf{E}}$$

and resistivity $(\rho) = \frac{\mathbf{E}}{\mathbf{j}} = \frac{\mathbf{EA}}{\mathbf{I}}$
 $\therefore \quad \mu = \frac{\mathbf{v}_{d}\mathbf{A}}{i\rho}$
 $= \frac{1.1 \times 10^{-3} \times \pi \times (5 \times 10^{-3})^{2}}{5 \times 1.7 \times 10^{-8}}$
 $\mu = 1.0 \frac{m^{2}}{Vs}$

- 29. Given below in the left column are different modes of communication using the kinds of waves given in the right column.
 - A. Optical Fibre P. Ultrasound Communication
 - B. Radar Q. Infrared Light
 - C. Sonar R. Microwaves
 - D. Mobile Phones S. Radio Waves From the options given below, find the most

appropriate match between entries in the left and the right column.

- (1) A-Q, B-S, C-P, D-R
- (2) A-Q, B-S, C-R, D-P
- (3) A-S, B-Q, C-R, D-P
- (4) A-R, B-P, C-S, D-Q

Answer (1)

- Sol. Optical Fibre Communication Infrared Light Radar – Radio Waves Sonar – Ultrasound
 - Mobile Phones Microwaves
- 30. The displacement of a damped harmonic oscillator is given by $x(t) = e^{-0.1t} \cos(10\pi t + \phi)$. Here t is in seconds. The time taken for its amplitude of vibration to

drop to half of its initial value is close to :

- (1) 7 s
- (2) 27 s
- (3) 13 s

(4) 4 s Answer (1)

Sol. Amplitude at (t = 0) $A_0 = e^{-0.1 \times 0} = 1$

$$\therefore \text{ at } t = t \qquad \text{if } A = \frac{A_0}{2}$$

$$\Rightarrow \frac{1}{2} = e^{-0.1t}$$

$$t = 10 \ln 2 \simeq 7 \text{ s}$$



PART-B : CHEMISTRY

- 1. Consider the following statements
 - (a) The pH of a mixture containing 400 mL of 0.1 M H₂SO₄ and 400 mL of 0.1 M NaOH will be approximately 1.3.
 - (b) Ionic product of water is temperature dependent.
 - (c) A monobasic acid with $K_a = 10^{-5}$ has a pH = 5. The degree of dissociation of this acid is 50%.
 - (d) The Le Chatelier's principle is not applicable to common-ion effect.

The correct statements are :

- (1) (a), (b) and (d)
- (2) (b) and (c)
- (3) (a) and (b)
- (4) (a), (b) and (c)

Answer (4)

 H_2SO_4 + NaOH \rightarrow NaHSO₄ + H_2O Sol. (a)

Initial moles 0.04 0.04 0

0.04 0.04

 $NaHSO_{4} \rightarrow Na^{+} + H^{+} + SO_{4}^{2-}$

$$[\text{H}^+] = \frac{0.04}{0.80} = 0.05 \text{ M}; \text{pH} = 1.3$$

(b) lonic product of water increases with increase of temperature because ionisation of water is endothermic.

0

(c) HA
$$\rightleftharpoons$$
 H⁺ + A⁻
 $C(1 - \alpha)$ C α C α pH = 5 & K_a = 10⁻⁵
 $10^{-5} = \frac{C\alpha^2}{1 - \alpha}$; C = 2 × 10⁻⁵ and α = 0.5

- 2. The oxoacid of sulphur that does not contain bond between sulphur atoms is :
 - (1) $H_2S_4O_6$
 - (2) $H_2S_2O_4$
 - (3) $H_2S_2O_7$
 - (4) $H_2S_2O_3$

```
Answer (3)
```



 $H_2S_2O_7$ does not have S – S linkage

The graph between $|\psi|^2$ and r(radial distance) is 3. shown below. This represents :



- (1) 1s orbital (2) 2s orbital
- (3) 2p orbital (4) 3s orbital

Answer (2)

Sol. The given probability density curve is for 2s orbital because it has only one radial node. Among other given orbitals, 1s and 2p do not have any radial node and 3s has two radial nodes.



- 4. Which of the following is a condensation polymer?
 - (1) Nylon 6, 6
 - (2) Teflon
 - (3) Buna S
 - (4) Neoprene

Answer (1)

Sol. Nylon 6, 6 is obtained by condensation polymerisation of hexamethylenediamine and adipic acid



$$\longrightarrow \left\{ \mathsf{NH}_{(\mathsf{CH}_2)_6} - \mathsf{NH}_{\mathsf{CO}_{-}}(\mathsf{CH}_2)_4 - \mathsf{CO}_{\mathsf{n}} \right\}$$
Nylon 6, 6

So, Nylon 6, 6 is a condensation polymer. Other polymers given, i.e., Buna-S, Teflon and Neoprene are addition polymers.

- 5. The principle of column chromatography is :
 - (1) Differential adsorption of the substances on the solid phase.
 - (2) Gravitational force.
 - (3) Differential absorption of the substances on the solid phase.
 - (4) Capillary action.

Answer (1)

- Sol. In column chromatograph a solid adsorbent is packed in a column and a solution containing number of solute particles is allowed to flow down the column. The solute molecules get adsorbed on the surface of adsorbent. So it is differential adsorption of the substances on the solid phase.
- 6. The major product of the following reaction is :

$$(1) CH_{3} - CH_{2}CH_{2}CH_{3} - CH_{3}CH_{3}CH_{3}OH \rightarrow H Br$$

$$(1) CH_{3} - CH_{2}CH_{2}CH_{3} (2) CH_{3} - CH_{1}CH_{3}$$

$$(2) CH_{3} - CH_{2}CH_{3} H OCH_{3}$$

$$(3) CH_{3} CH_{2} CH_{3} CH_{3} -CH_{2} (4) CH_{3} -CH_{2} CH_{3} CH_{3} -CH_{2} CH_{3} CH_{3} -CH_{3} CH_{3} CH_{3} -CH_{3} CH_{3} C$$

Answer (1)

Sol.
$$CH_3 - CH - CH - CH_3 \xrightarrow{CH_3OH} CH_3 - CH_3 - CH_3 + Br^{-1}$$

 $CH_3 = Br^{-1}$
 $CH_3 - CH_3 - CH_3$

ċн₃

сн

- decrease in the concentration of electrolyte.
- S2 : Molar conductivity always increases with decrease in the concentration of electrolyte.

The correct option among the following is :

- (1) S1 is wrong and S2 is correct
- (2) S1 is correct and S2 is wrong
- (3) Both S1 and S2 are wrong
- (4) Both S1 and S2 are correct

Answer (1)

- Sol. Conductivity of an electrolyte is the conductance of 1 cm³ of the given electrolyte. So, it increases with the increase of concentration of electrolyte. Molar conductivity (λ_m) is the conductance of a solution containing 1 mole of the electrolyte. It increases with the decrease of concentration (i) due to increase in interionic attraction for strong electrolytes and (ii) due to decrease in degree of ionisation for weak electrolytes. Therefore, (S_1) is wrong and (S_2) is correct.
- 8. Consider the following table :

Gas	a/(k Pa dm ⁶ mol ^{−1})	b/(dm ³ mol ⁻¹)
Α	642.32	0.05196
В	155.21	0.04136
С	431.91	0.05196
D	155.21	0.4382

a and b are van der Waals constants. The correct statement about the gases is :

- Gas C will occupy lesser volume than gas A; gas B will be lesser compressible than gas D
- (2) Gas C will occupy more volume than gas A; gas B will be more compressible than gas D
- (3) Gas C will occupy lesser volume than gasA; gas B will be more compressible than gas D
- (4) Gas C will occupy more volume than gas A; gas B will be lesser compressible than gas D

Answer (2)

Sol. If two gases have same value of 'b' but different values of 'a', then the gas having a larger value of 'a' will occupy lesser volume. This is because the gas having larger value of



If two gases have same value of 'a' but different values of 'b', then the smaller value of 'b' will occupy lesser volume and hence will be more compressible.

- 9. Amylopectin is composed of
 - (1) β -D-glucose, C₁-C₄ and C₂-C₆ linkages
 - (2) α -D-glucose, C₁-C₄ and C₂-C₆ linkages
 - (3) β -D-glucose, C₁ C₄ and C₁ C₆ linkages
 - (4) α -D-glucose, C₁-C₄ and C₁-C₆ linkages

Answer (4)

- Sol. Starch is a polymer of α -D-glucose. It has two components, namely
 - (i) Amylose and
 - (ii) Amylopectin

Amylose has only α -1,4-glycosidic linkage and is a linear polymer

Amylopectin has α -1, 6-glycosidic linkage in addition to α -1,4-glycosidic linkage and is a cross-linked polymer.

10. A bacterial infection in an internal wound grows as $N'(t) = N_0 exp(t)$, where the time t is in hours. A dose of antibiotic, taken orally, needs 1 hour to reach the wound. Once it reaches there, the bacterial population goes down as

$$\frac{dN}{dt}$$
 = - 5N². What will be the plot of $\frac{N_0}{N}$ vs. t

after 1 hour ?





Sol. When drug is administered bacterial growth is

given by
$$\frac{dN}{dt} = -5N^2$$

 $\Rightarrow \frac{N_0}{N_t} = 1+5t N_0$. Thus $\frac{N_0}{N_t}$ increases linearly
with t.



Answer (4)



The acylation of $\rm NH_2$ group takes place and not of OH group due to lower electronegativity of N-atom.

- 12. Consider the hydrated ions of Ti²⁺, V²⁺, Ti³⁺, and Sc³⁺. The correct order of their spin-only magnetic moments is :
 - (1) $Sc^{3+} < Ti^{3+} < Ti^{2+} < V^{2+}$
 - (2) $Ti^{3+} < Ti^{2+} < Sc^{3+} < V^{2+}$
 - (3) $Sc^{3+} < Ti^{3+} < V^{2+} < Ti^{2+}$
 - (4) $V^{2+} < Ti^{2+} < Ti^{3+} < Sc^{3+}$

Answer (1)

Sol. Electronic configuration of the given transition metal ions are

 V^{2+} (Z = 23) $1s^22s^22p^63s^23p^63d^3$



increasing order of magnetic moment is $Sc^{3+} < Ti^{3+} < Ti^{2+} < V^{2+}$

0 1 2 3 unpaired electrons

13. The isoelectronic set of ions is :

(1) N^{3-} , Li⁺, Mg²⁺ and O²⁻

(2) Li⁺, Na⁺, O²⁻ and F⁻

(3) N³⁻, O²⁻, F⁻ and Na⁺

Answer (3)

- Sol. Atomic numbers of N, O, F and Na are 7, 8, 9 and 11 respectively. Therefore, total number of electrons in each of N³⁻, O²⁻, F⁻ and Na⁺ is 10 and hence they are isoelectronic.
- 14. The alloy used in the construction of aircrafts is :

(1)	Mg - Zn	(2)	Mg - Sn
(3)	Mg - Mn	(4)	Mg - Al

Answer (4)

- Sol. An alloy of Mg and Al called magnalium is used in manufacturing of aircraft due to its light weight and high strength.
- 15. Ethylamine $(C_2H_5NH_2)$ can be obtained from N-ethylphthalimide on treatment with :

(1)	NH ₂ NH ₂	(2)	$NaBH_4$
(3)	H ₂ O	(4)	CaH ₂

Answer (1)

Sol. N-ethyl phthalimide on treatment with NH₂—NH₂ gives ethylamine.



- 16. A process will be spontaneous at all temperatures if :
 - (1) $\Delta H < 0$ and $\Delta S > 0$
 - (2) $\Delta H > 0$ and $\Delta S < 0$
 - (3) $\Delta H > 0$ and $\Delta S > 0$

(4) $\Delta H < 0$ and $\Delta S < 0$

Answer (1)

Sol. A reaction is spontaneous if ${\scriptstyle \Delta G}_{_{\scriptscriptstyle SVS}}$ is negative.

 $\Delta \mathbf{G}_{\mathrm{sys}} = \Delta \mathbf{H}_{\mathrm{sys}} - \mathbf{T} \Delta \mathbf{S}_{\mathrm{sys}}$

A reaction will be spontaneous at all temperatures if ΔH_{sys} is negative and ΔS_{sys} = +ve

17. Increasing rate of S_N 1 reaction in the following compounds is :



Answer (1)

Sol. The rate of $S_N 1$ is decided by the stability of carbocation formed in the rate determining step.







Carbocation (D) is most stable due to +R effect of CH_3O group, (C) is stabilised by +I and +H effect of CH_3 group; (B) is least stable due to -I effect of MeO group. So increasing order of rate of $S_N 1$ is (B) < (A) < (C) < (D)

18. The major product of the following reaction is :



- (ch chance i, 2-diamine, 0x 0xalate)
- (1) [Zn(en)Cl₂]
- (2) [Pt(en)Cl₂]
- (3) [Cr(en)₂(ox)]⁺
- (4) [Pt(en)₂Cl₂]²⁺

Answer (4)



Cis-trans isomerism is possible with $[Pt(en)_2Cl_2]^{2+}$. $[Cr(en)_2Ox]^+$ shows optical isomerism but not geometrical isomerism. The other two complexes, i.e. $[Pt(en)Cl_2]$ and $[Zn(en)Cl_2]$ do not show stereoisomerism.

20. Major products of the following reaction are :





$$\begin{array}{c} & & \\ & &$$



- 21. The correct order of catenation is :
 - (1) C > Si > Ge \approx Sn
 - (2) C > Sn > Si \approx Ge
 - (3) Si > Sn > C > Ge
 - (4) Ge > Sn > Si > C

Answer (1)

Sol. The order of catenation property amongst 14th group elements is based on bond enthalpy values of identical atoms of the same element. The decreasing order of bond enthalpy values is

$$\begin{array}{c} \textbf{C-C} > \textbf{Si-Si} > \textbf{Ge-Ge} \approx \textbf{Sn-Sn} \\ \textbf{Bond enthalpy} \begin{array}{c} {}^{348}_{\text{kJ/mol}} & {}^{297}_{\text{kJ/mol}} > {}^{260}_{\text{kJ/mol}} \\ \end{array} \end{array}$$

 \therefore Decreasing order of catenation is

 $\textbf{C > Si > Ge} \approx \textbf{Sn}$

22. The increasing order of the reactivity of the following compounds towards electrophilic aromatic substitution reactions is :



Answer (4)

Sol. CH₃ group when bonded to benzene increases the electron density of benzene by +I and hyper conjugation effects and hence makes the compound more reactive towards EAS. CI group decreases the electron density of benzene by -I effect, and CH₃CO group strongly decreases the electron density of benzene by -I and -R effects. Therefore, correct increasing order the given compounds towards EAS is



23. During the change of O_2 to O_2^- , the incoming electron goes to the orbital :

(1) $\pi 2 p_x$ (2) $\pi^* 2 p_x$ (3) $\pi 2 p_y$ (4) $\sigma^* 2 p_z$

Answer (2)

Sol. Electronic configuration of O₂ is

$$\sigma_{1s}^2 \sigma_{1s}^{\star^2} \sigma_{2s}^2 \sigma_{2s}^{\star^2} \sigma_{2p_z}^2 \pi_{2p_x}^2 = \pi_{2p_y}^2 \pi_{2p_x}^{\star 1} = \pi_{2p_y}^{\star 1}$$

When O_2 gains an electron to form O_2^- , the

incoming electron goes to $\pi^{\star}_{2p_{u}}$ or $\pi^{\star}_{2p_{u}}$

- 24. The regions of the atmosphere, where clouds form and where we live, respectively, are :
 - (1) Troposphere and Troposphere
 - (2) Stratosphere and Troposphere
 - (3) Troposphere and Stratosphere
 - (4) Stratosphere and Stratosphere

Answer (1)

- **Sol.** The lowest region of atmosphere in which human beings live is troposphere. It extends up to a height of 10 km from sea level. Clouds are also formed in this layer.
- 25. At room temperature, a dilute solution of urea is prepared by dissolving 0.60 g of urea in 360 g of water. If the vapour pressure of pure water at this temperature is 35 mmHg, lowering of vapour pressure will be : (molar mass of urea = 60 g mol⁻¹)

(1)	0.031	mmHg	(2)	0.017	mmHg
-----	-------	------	-----	-------	------

(3) 0.028 mmHg (4) 0.027 mmHg

Answer (2)

Sol. Relative lowering of VP is given by

$$\frac{\textbf{P}_{B}^{\circ}-\textbf{P}_{B}}{\textbf{P}_{B}^{\circ}}=\textbf{x}_{A}=\frac{\textbf{n}_{A}}{\textbf{n}_{A}+\textbf{n}_{B}}\ \simeq\frac{\textbf{n}_{A}}{\textbf{n}_{B}}$$

$$\frac{P_{B}^{\circ}-P_{B}}{35}=\frac{0.6\times18}{60\times360}=\frac{1}{2000}$$

On solving, $\Delta P_B = P_B^\circ - P_B = 0.017$



adsorption isotherm equation

$$\frac{\mathbf{x}}{\mathbf{m}} = \mathbf{k}\mathbf{p}^{0.6}$$

Adsorption of the gas increases with :

- (1) Increase in p and decrease in T
- (2) Decrease in p and decrease in T
- (3) Increase in p and increase in T
- (4) Decrease in p and increase in T

Answer (1)

Sol. Freundlich adsorption is applicable for physical adsorption. The variation of extent of adsorption with (i) Pressure and (ii) Temp is given by the following curves.



So, extent of adsorption increases with increase of pressure and decrease of temperature.

- 27. The synonym for water gas when used in the production of methanol is :
 - (1) fuel gas
 - (2) syn gas
 - (3) laughing gas
 - (4) natural gas

Answer (2)

Sol. When steam is passed over red hot coke, an equimolar mixture of CO and H₂ is obtained

 $H_2O(g) + C \longrightarrow CO + H_2$

Steam Red hot

The gaseous mixture thus obtained is called water gas or syn. gas.

28. At 300 K and 1 atmospheric pressure, 10 mL of a hydrocarbon required 55 mL of O₂ for complete combustion, and 40 mL of CO₂ is formed. The formula of the hydrocarbon is :

(1)	C_4H_{10}	(2)	C_4H_8
(3)	C ₄ H ₆	(4)	C₄H ₇ CI

Sol. CxHy +
$$\left(x + \frac{y}{4}\right)O_2 \longrightarrow xCO_2 + \frac{y}{2}H_2O$$

10 ml 55 ml

0 55 - 10
$$\left(x + \frac{y}{4} \right)$$
 10 x

Vol. of CO_2 , 10x = 40; x = 4

$$55 - 10\left(\mathbf{x} + \frac{\mathbf{y}}{4}\right) = 0 ; \qquad \mathbf{y} = 6$$

- \therefore Hydrocarbon is C₄H₆
- 29. Three complexes, $[CoCI(NH_3)_5]^{2+}(I),$ $[Co(NH_3)_5H_2O]^{3+}(II)$ and $[Co(NH_3)_6]^{3+}(III)$ absorb light in the visible region. The correct order of the wavelength of light absorbed by them is :
 - (1) (I) > (II) > (III)(2) (II) > (I) > (III)

(3) (III) > (I) > (II)(4) (III) > (II) > (I)

Answer (1)

- Sol. In a co-ordination compound, the strong field ligand causes higher splitting of the d-orbitals. Wavelength of the energy absorbed by the coordination compound is inversely proportional to ligand field strength of the given coordination compound. The decreasing order of ligand field strength is $NH_3 > H_2O > CI$. Therefore decreasing order of wavelength absorbed is (I) > (II) > (III).
- 30. Match the refining methods (Column I) with metals (Column II).

Column I	Column II
(Refining methods)	(Metals)
(I) Liquation	(a) Zr
(II) Zone Refining	(b) Ni
(III) Mond Process	(c) Sn
(IV) Van Arkel Method	(d) Ga
(1) (I)-(c); (II)-(a); (III)-(b)); (IV)-(d)
(2) (I)-(b); (II)-(d); (III)-(a); (IV)-(c)	
(3) (I)-(c); (II)-(d); (III)-(b); (IV)-(a)	
(4) (I)-(b); (II)-(c); (III)-(d)); (IV)-(a)
Answer (3)	
Sol. Mond's process is used Arkel method is used fo for Sn and zone refining	for refining of Ni, Van or Zr, Liquation is used is used for Ga.

So, correct match is

(I)-(c); (II)-(d); (III)-(b); (IV)-(a)


PART-C : MATHEMATICS

given $|z| = \sqrt{\frac{2}{5}}$ 1. All the pairs (x, y) that satisfy the inequality $2^{\sqrt{\sin^2 x - 2\sin x + 5}} \cdot \frac{1}{4\sin^2 y} \le 1$ also satisfy the equation : so $\sqrt{\frac{2}{5}} = \frac{2}{\sqrt{1+a^2}}$ from equation (i) (1) $\sin x = |\sin y|$ (square both side) (2) $\sin x = 2 \sin y$ $\Rightarrow \frac{2}{5} = \frac{4}{1+a^2}$ (3) $2 \sin x = \sin y$ (4) $2|\sin x| = 3 \sin y$ \Rightarrow 1 + a² = 10 Answer (1) $2^{\sqrt{\sin^2 x - 2\sin x + 5}} < 2^{2\sin^2 y}$ Sol. $a^2 = 9$ $\Rightarrow \sqrt{\sin^2 x - 2\sin x + 5} < 2\sin^2 y$ \Rightarrow a ± 3 \therefore (a > 0) \therefore a = 3 Hence $z = \frac{1+i^2+2i}{3-i} = \frac{2i}{3-i} = \frac{2i(3+i)}{10} = \frac{-1+3i}{5}$ $\Rightarrow \sqrt{(\sin x - 1)^2 + 4} \leq 2 \sin^2 y$ it is true when sinx = 1 $\overline{z} = \frac{-1}{5} - \frac{3}{5}i$ |siny| = 1so sinx = |siny|The number of 6 digit numbers that can be 3. formed using the digits 0, 1, 2, 5, 7 and 9 which 2. If a > 0 and $z = \frac{(1+i)^2}{a-i}$, has magnitude $\sqrt{\frac{2}{5}}$, are divisible by 11 and no digit is repeated, is : (1) 72 (2) 48 then \overline{z} is equal to : (4) 36 (3) 60 (1) $-\frac{1}{5}+\frac{3}{5}i$ (2) $-\frac{3}{5}-\frac{1}{5}i$ Answer (3) (3) $\frac{1}{5} - \frac{3}{5}i$ **Sol.** $a_1 a_2 a_3 a_4 a_5 a_6$ digit 0, 1, 2, 5, 7, 9 (4) $-\frac{1}{5}-\frac{3}{5}i$ $(a_1 + a_2 + a_5) - (a_2 + a_4 + a_6) = 11 \text{ K}$ Answer (4) so (1, 2, 9) (0, 5, 7) **Sol.** $z = \frac{(1+i)^2}{2-i} \times \frac{a+i}{2+i}$ Now number of ways to arranging them $= 3! \times 3! + 3! \times 2 \times 2$ $= 6 \times 6 + 6 \times 4$ $z = \frac{(1-1+2i)(a+i)}{a^2+1} = \frac{2ai-2}{a^2+1}$...(i) = 6 × 10 = 60 $|z| = \sqrt{\left(\frac{-2}{a^2+1}\right)^2 + \left(\frac{2a}{a^2+1}\right)^2} = \sqrt{\frac{4+4a^2}{(a^2+1)^2}}$ The value of $\int_{1}^{2\pi} \left[\sin 2x(1+\cos 3x)\right] dx$, where [t] 4. denotes the greatest integer function, is : $=\sqrt{\frac{4(1+a^2)}{(1+a^2)^2}}=\frac{2}{\sqrt{1+a^2}}$ **(1)** π **(2)** –π **(3)** –2π **(4)** 2π



Sol.
$$I = \int_{0}^{2\pi} \left[\sin 2x (1 + \cos 3x) \right] dx \qquad \dots(i)$$

$$\therefore \int_{0}^{\pi} f(x) = \int_{0}^{\pi} f(a - x) dx$$

$$\therefore I = \int_{0}^{2\pi} \left[-\sin 2x (1 + \cos 3x) \right] dx \qquad \dots(ii)$$

By (i) + (ii)

$$2I = \int_{0}^{2\pi} (-1) dx$$

$$2I = -(x)_{0}^{2\pi}$$

$$\Rightarrow I = -\pi$$

5. If $\Delta_{1} = \begin{vmatrix} x & \sin \theta & \cos \theta \\ -\sin \theta & -x & 1 \\ \cos \theta & 1 & x \end{vmatrix} and$

$$\Delta_{2} = \begin{vmatrix} x & \sin 2\theta & \cos 2\theta \\ -\sin 2\theta & -x & 1 \\ \cos 2\theta & 1 & x \end{vmatrix} , x \neq 0; then$$

for all $\theta \in \left(0, \frac{\pi}{2}\right)$:
(1) $\Delta_{1} + \Delta_{2} = -2x^{3}$
(2) $\Delta_{1} - \Delta_{2} = -2x^{3}$
(3) $\Delta_{4} + \Delta_{2} = -2(x^{3} + x - 1)$
(4) $\Delta_{1} - \Delta_{2} = x(\cos 2\theta - \cos 4\theta)$
Answer (1)
Sol. $\Delta_{4} = \begin{vmatrix} x & \sin \theta & \cos \theta \\ -\sin \theta & -x & 1 \\ \cos \theta & 1 & x \end{vmatrix}$

$$= x(-x^{2} - 1) - \sin \theta(-x \sin \theta - \cos \theta) + \cos \theta(-\sin \theta + x \cos \theta)$$

$$= -x^{3} - x + x \sin^{2}\theta + \sin \theta \cos \theta - \cos \theta \sin \theta + x \cos^{2}\theta$$

$$= -x^{3} - x + x = -x^{3}$$

Similarly $\Delta_{4} = -x^{3} + x = -2x^{3}$

6. If
$$f(x) = \begin{cases} x , x < 0 \\ q , x = 0 \\ \frac{\sqrt{x + x^2} - \sqrt{x}}{x^{3/2}} , x > 0 \end{cases}$$

is continuous at x = 0, then the ordered pair (p, q) is equal to :

(1)
$$\left(\frac{5}{2}, \frac{1}{2}\right)$$
 (2) $\left(-\frac{3}{2}, \frac{1}{2}\right)$
(3) $\left(-\frac{3}{2}, -\frac{1}{2}\right)$ (4) $\left(-\frac{1}{2}, \frac{3}{2}\right)$

Answer (2)

Sol.
$$f(x) = \begin{cases} \frac{\sin(p+1)x + \sin x}{x} & x < 0 \\ q & x = 0 \\ \sqrt{x^2 + x} - \sqrt{x} & x > 0 \end{cases}$$

is continuous at $x = 0$
So $f(0^-) = f(0) = f(0^+)$...(1)
 $f(0^-) = \frac{h}{h \to 0} f(0^-h)$
 $= \frac{h}{h \to 0} \frac{\sin(p+1)(-h) + \sin(-h)}{-h}$
 $= \frac{h}{h \to 0} \frac{\sin(p+1)h}{h(p+1)} \times (p+1) + \frac{h}{h \to 0} \frac{\sinh h}{h}$
 $= (p+1) + 1 = p + 2$...(2)
Now $f(0^+) = \lim_{h \to 0} f(0^-h) = \lim_{h \to 0} \frac{\sqrt{h^2 + h} - \sqrt{h}}{h^{3/2}}$
 $= \lim_{h \to 0} \frac{\sqrt{h+1} - 1}{h} \times \frac{\sqrt{h+1} + 1}{\sqrt{h+1} + 1}$
 $= \lim_{h \to 0} \frac{\sqrt{h+1} - 1}{h(\sqrt{h+1} + 1)}$
 $= \lim_{h \to 0} \frac{1}{\sqrt{h+1} + 1} = \frac{1}{1+1} = \frac{1}{2}$...(3)



p + 2 = q =
$$\frac{1}{2}$$

So, q = $\frac{1}{2}$ and p = $\frac{1}{2}$ -2 = $\frac{-3}{2}$
(p, q) = $\left(-\frac{3}{2}, \frac{1}{2}\right)$

7. If the length of the perpendicular from the point $(\beta, 0, \beta)$ $(\beta \neq 0)$ to the line, $\frac{x}{1} = \frac{y-1}{0} = \frac{z+1}{-1} \text{ is } \sqrt{\frac{3}{2}} \text{ , then } \beta \text{ is equal to :}$ (1) -1 (2) -2 (3) 1 (4) 2

Answer (1)

Sol.
$$\frac{x}{1} = \frac{y-1}{0} = \frac{z+1}{-1} = p$$

$$P(\beta, 0, \beta)$$
any point on line A = (p, 1, -p - 1)
Now, DR of AP = \beta, 1 - 0, -p - 1 -
Which is perpendicular to line so
(p - β). 1 + 0.1 - 1(-p - 1 - β) = 0
 $\Rightarrow p - \beta + p + 1 + \beta = 0$
 $p = \frac{-1}{2}$
Point A $\left(\frac{-1}{2}, 1 - \frac{1}{2}\right)$
Now, distance AP = $\sqrt{\frac{3}{2}}$
 $\Rightarrow AP^2 = \frac{3}{2}$
 $\Rightarrow \left(\beta + \frac{1}{2}\right)^2 + 1 + \left(\beta + \frac{1}{2}\right)^2 = \frac{3}{2}$
 $2\left(\beta + \frac{1}{2}\right)^2 = \frac{1}{2}$
 $\Rightarrow \left(\beta + \frac{1}{2}\right)^2 = \frac{1}{4}$
 $\Rightarrow \beta = 0, -1, (\beta \neq 0)$
 $\therefore \overline{\beta = -1}$

to:
(1)
$$\frac{4}{3}(2)^{\frac{4}{3}}$$
(2) $\frac{3}{4}(2)^{\frac{4}{3}} - \frac{3}{4}$
(3) $\frac{4}{3}(2)^{\frac{3}{4}}$
(4) $\frac{3}{4}(2)^{\frac{4}{3}} - \frac{4}{3}$

Answer (2)

Sol.
$$\lim_{n \to \infty} \frac{(n+1)^{\frac{1}{3}} + (n+2)^{\frac{1}{3}} + \ldots + (n+n)^{\frac{1}{3}}}{n(n)^{\frac{1}{3}}}$$

$$= \lim_{n \to \infty} \sum_{r=1}^{n} \frac{(n+r)^{\frac{1}{3}}}{n \cdot n^{\frac{1}{3}}} \qquad \frac{r}{n} \to x \text{ and } \frac{1}{n} \to dx$$

$$= \int_{0}^{1} (1+x)^{\frac{1}{3}} dx$$

$$= \left[\frac{3}{4}(1+x)^{\frac{4}{3}}\right]_{0}^{1} = \frac{3}{4}(2)^{\frac{4}{3}} - \frac{3}$$

9. The line x = y touches a circle at the point (1, 1). If the circle also passes through the point (1, -3), then its radius is :

(1) <u>3√2</u>	(2) 2
(2) $\sim \sqrt{2}$	(4) 2

(3) $2\sqrt{2}$ (4) 3

Answer (3)

β>

Sol. Equation of circle = $(x - 1)^2 + (y - 1)^2 + \lambda(y - x) = 0$ Which passes through (1, -3)



Now equation of circle $(x - 1)^2 + (y - 1)^2 + 4y - 4x = 0$ $\Rightarrow x^2 + y^2 - 6x + 2y + 2 = 0$ radius = $\sqrt{9 + 1 - 2} = 2\sqrt{2}$



(1)
$$(p \lor q) \lor (p \lor \neg q)$$

(2) $(p \land q) \lor (p \land \neg q)$
(3) $(p \lor q) \land (p \lor \neg q)$
(4) $(p \lor q) \land (p \lor \neg q)$
Answer (1)
Sol. $(p \lor q) \lor (p \lor \neg q)$
 $= p \lor (q \lor p) \lor \neg q$
 $= (p \lor p) \lor (q \lor \neg q)$
 $= p \lor T$
 $= T$ so first statement is tautology
11. If $\lim_{x \to 1} \frac{x^4 - 1}{x - 1} = \lim_{x \to k} \frac{x^3 - k^3}{x^2 - k^2}$, then k is :
(1) $\frac{4}{3}$
(2) $\frac{3}{2}$
(3) $\frac{8}{3}$
(4) $\frac{3}{8}$
Answer (3)
Sol. If $\lim_{x \to 1} \frac{x^4 - 1}{x - 1} = \lim_{x \to K} \left(\frac{x^3 - k^3}{x^2 - k^2}\right)$
L·H·S·

origin and passing through the point $(4, -2\sqrt{3})$ is $5x = 4\sqrt{5}$ and its eccentricity is e, then : (1) $4e^4 + 8e^2 - 35 = 0$ (2) $4e^4 - 24e^2 + 35 = 0$ (3) $4e^4 - 12e^2 - 27 = 0$ (4) $4e^4 - 24e^2 + 27 = 0$

Answer (2)

Sol.
$$x = \frac{4}{\sqrt{5}}$$

$$\therefore \quad \boxed{\frac{a}{e} = \frac{4}{\sqrt{5}}}$$

$$(\frac{a}{e}, 0) \quad \odot (4, -2\sqrt{3})$$

Equation of hyperbola

$$\frac{x^{2}}{a^{2}} - \frac{y^{2}}{b^{2}} = 1 \text{ it passes through } (4, -2\sqrt{3})$$

$$\therefore e^{2} = 1 + \frac{b^{2}}{a^{2}}$$

$$\Rightarrow \boxed{a^{2}e^{2} - a^{2} = b^{2}}$$

$$\Rightarrow \frac{16}{a^{2}} - \frac{12}{a^{2}e^{2} - a^{2}} = 1$$

$$\Rightarrow \frac{4}{a^{2}} \left[\frac{4}{1} - \frac{3}{e^{2} - 1} \right] = 1$$

$$\Rightarrow 4e^{2} - 4 - 3 = (e^{2} - 1) \left(\frac{a^{2}}{4} \right)$$

$$\Rightarrow 4(4e^{3} - 7) = (e^{2} - 1) \left(\frac{4e}{\sqrt{5}} \right)^{2}$$

$$\Rightarrow 4e^{4} - 24e^{2} + 35 = 0$$

13. If the line x - 2y = 12 is tangent to the ellipse

$$\frac{x^{2}}{a^{2}} + \frac{y^{2}}{b^{2}} = 1 \text{ at the point } \left(3, \frac{-9}{2} \right), \text{ then the length of the latus rectum of the ellipse is :}$$

(1) 5 (2) $8\sqrt{3}$

(3) $12\sqrt{2}$ (4) 9



 $\operatorname{Lt}_{x \to 1} \frac{x^4 - 1}{x - 1} = \left(\frac{0}{0} \text{ form}\right)$

Now, $\lim_{x\to K} \frac{x^3 - k^3}{x^2 - k^2} = 4$

 $\Rightarrow \lim_{x \to K} \frac{3x^2}{2x} = 4$

 $\Rightarrow \frac{3}{2} k = 4$

 $k = \frac{8}{3}$

 $\operatorname{Lt}_{x\to 1}\frac{4x^3}{1}=4$

Sol. Equation of tangent at $\left(3, -\frac{9}{2}\right)$ to $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is $\frac{3x}{a^2} - \frac{y9}{2b^2} = 1$ which is equivalent to x - 2y = 12 $\frac{3}{a^2} = \frac{-9}{2b^2 \cdot (-2)} = \frac{1}{12}$ (On comparing) $a^2 = 3 \times 12$ and $b^2 = \frac{9 \times 12}{4}$ $\boxed{a = 6} \Rightarrow b = 3\sqrt{3}$

So latus rectum = $\frac{2b^2}{a} = \frac{2 \times 27}{6} = 9$

14. Let A(3, 0 –1), B(2, 10, 6) and C(1, 2, 1) be the vertices of a triangle and M be the mid point of AC. If G divides BM in the ratio, 2 : 1 then $\cos(\angle \text{GOA})$ (O being the origin) is equal to :

(1)
$$\frac{1}{6\sqrt{10}}$$

(2) $\frac{1}{\sqrt{30}}$
(3) $\frac{1}{2\sqrt{15}}$
(4) $\frac{1}{\sqrt{15}}$

Answer (4)

Sol. G is the centroid of $\triangle ABC$



 $OG = \sqrt{4+16+4}, \quad OA = \sqrt{9+1}$

 $AG = \sqrt{1+16+9}$

$$=\frac{8}{2\sqrt{8\times3\times2\times5}}$$
$$=\frac{4}{4\sqrt{15}}=\frac{1}{\sqrt{15}}$$

15. If the system of linear equations

x + y + z = 5x + 2y + 2z = 6**x** + 3**y** + λ **z** = μ , (λ , $\mu \in \mathbf{R}$), has infinitely many solutions, then the value of λ + μ is : (1) 10 (2) 12 (3) 7 (4) 9 Answer (1) **Sol.** x + y + z = 5x + 2y + 2z = 6 $x + 3y + \lambda z = \mu$ have infinite solution $\Delta = 0$, $\Delta x = \Delta y = \Delta z = 0$ 1 1 1 $\Delta = \begin{vmatrix} \mathbf{1} & \mathbf{2} & \mathbf{2} \\ \mathbf{1} & \mathbf{3} & \lambda \end{vmatrix} = \mathbf{0}$ \Rightarrow 1(2 λ -6) - 1(λ - 2) + 1 (3 - 2) = 0 \Rightarrow 2 λ - 6 - λ + 2 + 1 = 0 $\lambda = 3$ Now, $\Delta \mathbf{x} = \begin{vmatrix} 5 & 1 & 1 \\ 6 & 2 & 2 \\ 11 & 3 & 3 \end{vmatrix} = 0, \ \Delta \mathbf{y} = \begin{vmatrix} 1 & 5 & 1 \\ 1 & 6 & 2 \\ 1 & 1 & 3 \end{vmatrix} = 0$ $\Rightarrow \begin{vmatrix} 1 & 5 & 1 \\ 0 & 1 & 1 \\ 0 & \mu - 5 & 2 \end{vmatrix} = 0$ \Rightarrow 1(2 – μ + 5) = 0 $\mu = 7$ $\Delta z = \begin{vmatrix} 1 & 1 & 5 \\ 1 & 2 & 6 \\ 1 & 3 & \mu \end{vmatrix} = \begin{vmatrix} 1 & 1 & 5 \\ 0 & -1 & -1 \\ 0 & 2 & \mu - 5 \end{vmatrix}$ \Rightarrow 1 (5 – μ + 2) = 0 $\Rightarrow \mu = 7$ **So**, $\lambda + \mu = 10$



children each, then the conditional probability that all children are girls given that at least two are girls is :

(1)
$$\frac{1}{11}$$
 (2) $\frac{1}{12}$
(3) $\frac{1}{10}$ (4) $\frac{1}{17}$

Answer (1)

Sol. A = At least two girls

B = All girls

$$P\left(\frac{B}{A}\right) = \frac{P(B \cap A)}{P(A)}$$

$$= \frac{P(B)}{P(A)} = \frac{\left(\frac{1}{4}\right)^4}{1 - {}^4C_0\left(\frac{1}{2}\right)^4 - {}^4C_1\left(\frac{1}{2}\right)^4}$$

$$= \frac{1}{16 - 1 - 4} = \frac{1}{11}$$

17. If for some $x \in R$, the frequency distribution of the marks obtained by 20 students in a test is :

Marks	2	3	5	7
Frequency	$(x + 1)^{2}$	2x – 5	x ² – 3x	x

Then the mean of the marks is :

(1)	3.2	(2)	3.0
(3)	2.5	(4)	2.8

Answer (4)

Sol. Number of students

$$\Rightarrow (x + 1)^{2} + (2x - 5) + (x^{2} - 3x) + x = 20$$

$$\Rightarrow 2x^{2} + 2x - 4 = 20$$

$$x^{2} + x - 12 = 0$$

$$(x + 4) (x - 3) = 0$$

$$x = 3$$

So, Marks 2 3 5 7
No. of students 16 1 0 3

Average marks = $\frac{32+3+21}{20} = \frac{56}{20} = 2.8$

18. If the circles $x^2 + y^2 + 5Kx + 2y + K = 0$ and $2(x^2 + y^2) + 2Kx + 3y - 1 = 0$, (K \in R), intersect at the points P and Q, then the line 4x + 5y - K = 0 passes through P and Q, for :

- (3) Exactly two values of K
- (4) No value of K

Answer (4)

Sol.
$$S_1 \equiv x^2 + y^2 + 5Kx + 2y + K = 0$$

 $S_2 \equiv x^2 + y^2 + Kx + \frac{3}{2}y - \frac{1}{2} = 0$

Equation of common chord is

$$S_1 - S_2 = 0$$

$$\Rightarrow 4Kx + \frac{y}{2} + K + \frac{1}{2} = 0 \qquad \dots (1)$$

$$4x + 5y - K = 0 \qquad \dots (2) \text{ (given)}$$

On comparing (1) and (2)

$$\frac{4K}{4} = \frac{1}{10} = \frac{2K+1}{-2K}$$

$$\Rightarrow \quad \boxed{K = \frac{1}{10}} \text{ and } - 2K = 20K+10$$

$$\Rightarrow \quad 22K = -10$$

$$\boxed{K = \frac{-5}{11}}$$

- \therefore No value of K exists
- 19. The region represented by $|\mathbf{x} \mathbf{y}| \le 2$ and
 - $|\mathbf{x} + \mathbf{y}| \le 2$ is bounded by a :
 - (1) Square of side length $2\sqrt{2}$ units
 - (2) Square of area 16 sq. units
 - (3) Rhombus of side length 2 units
 - (4) Rhombus of area $8\sqrt{2}$ sq. units

Answer (1)

Sol.
$$C_1 : |y - x| \le 2$$

 $C_2 : |y + x| \le 2$
Now region is square





equal to :

 (1) 98
 (2) 38

 (3) 64
 (4) 76

Answer (4)

Sol. $3(a_1 + a_{16}) = 114$

- $a_1 + a_{16} = 38$ Now $a_1 + a_6 + a_{11} + a_{16} = 2(a_1 + a_{16})$ $= 2 \times 38 = 76$
- 21. If Q(0, -1, -3) is the image of the point P in the plane 3x y + 4z = 2 and R is the point (3, -1, -2), then the area (in sq. units) of $\triangle PQR$ is :

(1)
$$\frac{\sqrt{65}}{2}$$
 (2) $\frac{\sqrt{91}}{4}$
(3) $2\sqrt{13}$ (4) $\frac{\sqrt{91}}{2}$

Answer (4)

Sol. Image of Q in plane

$$\frac{(x-0)}{3} = \frac{(y+1)}{-1} = \frac{z+3}{+4} = \frac{-2(1-12-2)}{9+1+16} = 1$$

x = 3, y = -2, z = 1
P(3, -2, 1), Q(0, -1, -3), R(3, -1, -2)
Now area of \triangle PQR is

$$\frac{1}{2} |\overline{PQ} \times \overline{QR}| = \frac{1}{2} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -1 & 4 \\ 3 & 0 & 1 \end{vmatrix} \\$$
$$= \frac{1}{2} |\{\hat{i}(-1) - \hat{j}(3 - 12) + \hat{k}(3)\}|$$
$$= \frac{1}{2} \sqrt{(1 + 81 + 9)}$$
$$= \frac{\sqrt{91}}{2}$$

22. Let $f(x) = x^2$, $x \in R$. For any $A \subseteq R$, define $g(A) = \{x \in R : f(x) \in A\}$. If S = [0, 4], then which one of the following statements is not true ?

(1) $f(g(S)) = S$	(2) $g(f(S)) = g(S)$
(3) q(f(S)) ≠ S	(4) f(q(S)) ≠ f(S)

- $\begin{array}{l} \text{Sol. } \mathbf{I}(\mathbf{x}) = \mathbf{x} & \in \mathbf{R} \\ \text{g}(\mathbf{A}) = \{\mathbf{x} \in \mathbf{R} : \ \mathbf{f}(\mathbf{x}) \in \mathbf{A}\} \quad \mathbf{S} \equiv [0, \, 4] \\ \text{g}(\mathbf{S}) = \{\mathbf{x} \in \mathbf{R} : \ \mathbf{f}(\mathbf{x}) \in \mathbf{S}\} \\ = \{\mathbf{x} \in \mathbf{R} : \ 0 \le \mathbf{x}^2 \le 4\} \\ = \{\mathbf{x} \in \mathbf{R} : \ -2 \le \mathbf{x} \le 2\} \\ \therefore \quad \mathbf{g}(\mathbf{S}) \neq \mathbf{S} \\ \therefore \quad \mathbf{f}(\mathbf{g}(\mathbf{S})) \neq \mathbf{f}(\mathbf{S}) \\ \text{g}(\mathbf{f}(\mathbf{S})) = \{\mathbf{x} \in \mathbf{R} : \ \mathbf{f}(\mathbf{x}) \in \mathbf{f}(\mathbf{S})\} \\ = \{\mathbf{x} \in \mathbf{R} : \ \mathbf{x}^2 \in \mathbf{S}^2\} \\ = \{\mathbf{x} \in \mathbf{R} : \ 0 \le \mathbf{x}^2 \le 16\} \\ = \{\mathbf{x} \in \mathbf{R} : \ -4 \le \mathbf{x} \le 4\} \\ \therefore \quad \mathbf{g}(\mathbf{f}(\mathbf{S})) \neq \mathbf{g}(\mathbf{S}) \end{array}$
 - \therefore g(f(S)) = g(S) is incorrect
- 23. If the coefficients of x^2 and x^3 are both zero, in the expansion of the expression $(1 + ax + bx^2)$ $(1 - 3x)^{15}$ in powers of x, then the ordered pair (a, b) is equal to :
 - (1) (- 54, 315)
 - (2) (28, 861)
 - (3) (-21, 714)
 - (4) 28, 315

Answer (4)

Sol.
$$(1 + ax + bx^2)(1 - 3x)^{15}$$

Co-eff. of $x^2 = 1.^{15}C_2(-3)^2 + a.^{15}C_1(-3) + b.^{15}C_0$

$$= \frac{15 \times 14}{2} \times 9 - 15 \times 3a + b = 0 \text{ (Given)}$$

$$\Rightarrow 945 - 45a + b = 0 \qquad \dots(i)$$
Now co-eff. of $x^3 = 0$

$$\Rightarrow {}^{15}C_3(-3)^3 + a.{}^{15}C_2(-3)^2 + b.{}^{15}C_1(-3) = 0$$

$$\Rightarrow \frac{15 \times 14 \times 13}{3 \times 2} \times (-3 \times 3 \times 3) + a \times \frac{15 \times 14 \times 9}{2}$$

$$-b \times 3 \times 15 = 0$$

$$\Rightarrow 15 \times 3[-3 \times 7 \times 13 + a \times 7 \times 3 - b] = 0$$

$$\Rightarrow 21a - b = 273 \qquad \dots(ii)$$
From (i) and (ii)

$$a = +28, b = 315 \equiv (a, b) \equiv (28, 315)$$



$$\frac{3\times1^{\circ}}{1^{2}} + \frac{5\times(1^{\circ}+2^{\circ})}{1^{2}+2^{2}} + \frac{7\times(1^{\circ}+2^{\circ}+3^{\circ})}{1^{2}+2^{2}+3^{2}} + \dots$$

upto 10th term, is :
(1) 620 (2) 600
(3) 680 (4) 660

Answer (4)

Sol.
$$T_r = \frac{(2r+1)(1^3+2^3+3^3+...+r^3)}{1^2+2^2+3^2+...+r^2}$$

 $T_r = (2r+1)\left(\frac{r(r+1)}{2}\right)^2 \times \frac{6}{r(r+1)(2r+1)}$
 $T_r = \frac{3r(r+1)}{2}$

Now,

$$S = \sum_{r=1}^{10} T_r = \frac{3}{2} \sum_{r=1}^{10} (r^2 + r)$$
$$= \frac{3}{2} \left\{ \frac{10 \times (10 + 1)(2 \times 10 + 1)}{6} + \frac{10 \times 11}{2} \right\}$$
$$= \frac{3}{2} \left\{ \frac{10 \times 11 \times 21}{6} + 5 \times 11 \right\}$$
$$= \frac{3}{2} \times 5 \times 11 \times 8 = 660$$

25. Let $f(x) = e^x - x$ and $g(x) = x^2 - x$, $\forall x \in R$. Then the set of all $x \in R$, where the function h(x)=(fog) (x) is increasing, is :

(1)
$$\left[-1,\frac{-1}{2}\right] \cup \left[\frac{1}{2},\infty\right)$$
 (2) $[0,\infty)$
(3) $\left[0,\frac{1}{2}\right] \cup [1,\infty)$ (4) $\left[\frac{-1}{2},0\right] \cup [1,\infty)$

Answer (3)

·____

Sol. $f(x) = e^x - x$, $g(x) = x^2 - x$

$$f(g(x)) = e^{(x^2-x)} - (x^2 - x)$$

If f(g(x)) is increasing function

$$(f(g(x)))' = e^{(x^2-x)} \times (2x-1) - 2x + 1$$
$$= (2x-1)e^{(x^2-x)} + 1 - 2x$$
$$= (2x-1)[e^{(x^2-x)} - 1]$$
AB

$$\begin{array}{c|c} - +ve -ve + \\ \hline 0 & 1 \\ \hline 2 \end{array}$$

for $(f(g(x)))' \ge 0$,

$$\mathbf{x} \in \left[\mathbf{0}, \frac{\mathbf{1}}{\mathbf{2}}\right] \cup \left[\mathbf{1}, \infty\right)$$

26. If α and β are the roots of the quadratic equation, $x^2 + x \sin\theta - 2\sin\theta = 0$, $\theta \in \left(0, \frac{\pi}{2}\right)$, then $\frac{\alpha^{12} + \beta^{12}}{(\alpha^{-12} + \beta^{-12}) \cdot (\alpha - \beta)^{24}}$ is equal to :

$$(\alpha^{-12} + \beta^{-12}) \cdot (\alpha - \beta)^{24} \quad \text{is optimized}$$

$$(1) \quad \frac{2^{12}}{(\sin \theta - 8)^6} \qquad (2) \quad \frac{2^{12}}{(\sin \theta - 4)^{12}}$$

$$(3) \quad \frac{2^6}{(\sin \theta + 8)^{12}} \qquad (4) \quad \frac{2^{12}}{(\sin \theta + 8)^{12}}$$

Answer (4)

Sol. Given $\alpha + \beta = -\sin\theta$ and $\alpha\beta = -2\sin\theta$

$$\frac{(\alpha^{12} + \beta^{12})\alpha^{12}\beta^{12}}{(\alpha^{12} + \beta^{12})(\alpha - \beta)^{24}} = \frac{(\alpha\beta)^{12}}{(\alpha - \beta)^{24}}$$

$$|\alpha - \beta| = \sqrt{(\alpha + \beta)^2 - 4\alpha\beta} = \sqrt{\sin^2 \theta + 8\sin\theta}$$

Hence required quantity

$$\frac{(\alpha\beta)^{12}}{(\alpha-\beta)^{24}} = \frac{(2\sin\theta)^{12}}{\sin^{12}\theta(\sin\theta+8)^{12}} = \frac{2^{12}}{(\sin\theta+8)^{12}}$$

27. If y = y(x) is the solution of the differential equation $\frac{dy}{dx} = (\tan x - y) \sec^2 x$, $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, such that y(0) = 0, then $y\left(-\frac{\pi}{4}\right)$ is equal to :

(1)
$$\frac{1}{e} - 2$$

(2) $\frac{1}{2} - e$
(3) $e - 2$
(4) $2 + \frac{1}{e}$



Sol. $\frac{dy}{dx}$ + y sec² x = sec² x tanx \rightarrow This is linear differential equation

$$IF = e^{\int sec^2 x dx} = e^{tanx}$$

Now solution is

$$\mathbf{y} \cdot \mathbf{e}^{tanx} = \int \mathbf{e}^{tanx} \sec^2 \mathbf{x} tan \mathbf{x} d\mathbf{x}$$

∴ Let tanx = t

$$ye^{tanx} = \int e^t t dt$$

$$ye^{tanx} = te^{t} - e^{t} + c$$

 $ye^{tanx} = (tanx - 1)e^{tanx} + c$

$$y = (tanx - 1) + c \cdot e^{-tanx}$$

Given y(0) = 0

$$\Rightarrow$$
 0 = -1 + c

$$\mathbf{y}\left(-\frac{\pi}{4}\right) = -1 - 1 + \mathbf{e} = -2 + \mathbf{e}$$

- 28. Let $f : R \rightarrow R$ be differentiable at $c \in R$ and f(c) = 0. If g(x) = |f(x)|, then at x = c, g is :
 - (1) not differentiable if f'(c) = 0
 - (2) differentiable if f'(c) = 0
 - (3) not differentiable
 - (4) differentiable if $f'(c) \neq 0$

Sol.
$$\therefore$$
 $g'(c) = \lim_{x \to c} \frac{g(x) - g(c)}{x - c}$
 $\Rightarrow g'(c) = \lim_{x \to c} \frac{|f(x)| - |f(c)|}{x - c}$
 \therefore $f(c) = 0$
 $\Rightarrow g'(c) = \lim_{x \to c} \frac{|f(x)|}{x - c}$
 $\Rightarrow g'(c) = \lim_{x \to c} \frac{f(x)}{x - c}$ if $f(x) > 0$

$$\Rightarrow g'(c) = f'(c) = -f'(c)$$
$$\Rightarrow 2f'(c) = 0$$
$$\Rightarrow f'(c) = 0$$

29. ABC is a triangular park with AB = AC = 100 metres. A vertical tower is situated at the mid-point of BC. If the angles of elevation of the top of the tower at A and B are $\cot^{-1}(3\sqrt{2})$

and $\csc ec^{-1}(2\sqrt{2})$ respectively, then the height of the tower (in metres) is :

- (1) 20
- (2) 10√5
- (3) 25
- (4) $\frac{100}{3\sqrt{3}}$

Sol.





$$= A\left(\tan^{-1}\left(\frac{x-1}{3}\right) + \frac{f(x)}{x^2 - 2x + 10}\right) + C$$

where C is a constant of integration, then :

(1)
$$A = \frac{1}{81}$$
 and $f(x) = 3(x - 1)$
(2) $A = \frac{1}{54}$ and $f(x) = 3(x - 1)$
(3) $A = \frac{1}{27}$ and $f(x) = 9(x - 1)$
(4) $A = \frac{1}{54}$ and $f(x) = 9(x - 1)^2$

Answer (2)

Sol.
$$\int \frac{dx}{(x^2 - 2x + 10)^2} = \int \frac{dx}{((x - 1)^2 + 9)^2}$$

Let $(x - 1)^2 = 9\tan^2\theta$...(i)
 $\Rightarrow \tan \theta = \frac{x - 1}{3}$

 $2(x-1)dx = 18\tan\theta \sec^2\theta d\theta$ $\therefore I = \int \frac{18\tan\theta \sec^2\theta d\theta}{2\times 3\tan\theta \times 81\sec^4\theta}$ $I = \frac{1}{27}\int \cos^2\theta d\theta = \frac{1}{27} \times \frac{1}{2}\int (1+\cos 2\theta)d\theta$ $I = \frac{1}{54} \left\{ \theta + \frac{\sin 2\theta}{2} \right\} + c$ $I = \frac{1}{54} \left[\tan^{-1} \left(\frac{x-1}{3} \right) + \frac{1}{2} \times \frac{2\left(\frac{x-1}{3} \right)}{1+\left(\frac{x-1}{3} \right)^2} \right] + c$ $I = \frac{1}{54} \left[\tan^{-1} \left(\frac{x-1}{3} \right) + \frac{3(x-1)}{x^2 - 2x + 10} \right] + c$ So $A = \frac{1}{54}$ f(x) = 3(x-1)



Answers & Solutions

for

JEE (MAIN)-2019 (Online) Phase-2

(Physics, Chemistry and Mathematics)

Time : 3 hrs.

M.M.: 360

Important Instructions :

- 1. The test is of **3 hours** duration.
- 2. The Test Booklet consists of **90** questions. The maximum marks are **360**.
- 3. There are *three* parts in the question paper A, B, C consisting of **Physics**, **Chemistry** and **Mathematics** having 30 questions in each part of equal weightage.
- 4. Each question is allotted 4 (four) marks for each correct response. ¼ (one-fourth) marks will be deducted for indicating incorrect response of each question. No deduction from the total score will be made if no response is indicated for an item in the answer sheet.
- 5. There is only one correct response for each question.



1. One mole of an ideal gas passes through a process where pressure and volume obey the

relation
$$\mathbf{P} = \mathbf{P}_0 \left[1 - \frac{1}{2} \left(\frac{\mathbf{V}_0}{\mathbf{V}} \right)^2 \right]$$
. Here \mathbf{P}_0 and \mathbf{V}_0 are

constants. Calculate the change in the temperature of the gas if its volume changes from V_0 to $2V_0$.

(1)
$$\frac{1}{4} \frac{P_0 V_0}{R}$$
 (2) $\frac{5}{4} \frac{P_0 V_0}{R}$
(3) $\frac{1}{2} \frac{P_0 V_0}{R}$ (4) $\frac{3}{4} \frac{P_0 V_0}{R}$

Answer (2)

Sol. If
$$V_1 = V_0 \Rightarrow P_1 = P_0 \left[1 - \frac{1}{2} \right] = \frac{P_0}{2}$$

If $V_2 = 2V_0 \Rightarrow P_2 = P_0 \left[1 - \frac{1}{2} \left(\frac{1}{4} \right) \right] = \left(\frac{7P_0}{8} \right)$
 $\left(T = \frac{PV}{nR} \right) \Rightarrow \Delta T = \left| \frac{P_1 V_1}{nR} - \frac{P_2 V_2}{nR} \right|$
 $\Delta T = \left| \left(\frac{1}{nR} \right) \left(P_1 V_1 - P_2 V_2 \right) \right| = \left(\frac{1}{nR} \right) \left| \left(\frac{P_0 V_0}{2} - \frac{7P_0 V_0}{4} \right) \right|$
 $= \frac{5P_0 V_0}{4nR}$

2. The correct figure that shows, schematically, the wave pattern produced by superposition of two waves of frequencies 9 Hz and 11 Hz, is :



Sol. Beat frequency = $|f_1 - f_2| = 11 - 9 = 2$ Hz

3. A spaceship orbits around a planet at a height of 20 km from its surface. Assuming that only gravitational field of the planet acts on the spaceship, what will be the number of complete revolutions made by the spaceship in 24 hours around the planet?

[Given: Mass of planet = 8×10^{22} kg,

Radius of planet = 2×10^6 m,

Gravitational constant G = $6.67 \times 10^{-11} \text{ Nm}^2/\text{kg}^2$]

Answer (3)

Sol.
$$T = \frac{2\pi r}{v}, v = \sqrt{\frac{GM}{r}}$$

 $T = 2\pi r \sqrt{\frac{r}{GM}} = 2\pi \sqrt{\frac{r^3}{GM}}$
 $T = 2\pi \sqrt{\frac{(202)^3 \times 10^{12}}{6.67 \times 10^{-11} \times 8 \times 10^{22}}}$ sec
 $T = 7812.2 \text{ s}$
 $T \simeq 2.17 \text{hr} \Rightarrow 11 \text{ revolutions}$

4. A plane is inclined at an angle $\alpha = 30^{\circ}$ with respect to the horizontal. A particle is projected with a speed u = 2 ms⁻¹, from the base of the plane, making an angle $\theta = 15^{\circ}$ with respect to the plane as shown in the figure. The distance from the base, at which the particle hits the plane is close to :

$$(Take g = 10 ms^{-2})$$

(1) 18 cm
(3) 14 cm
(4) 26 cm
Answer (2)
Sol. Time of flight (T) =
$$\frac{2u\sin\alpha}{g\cos\beta}$$

T = $\frac{(2)(2\sin15)}{g\cos30} = \frac{4}{10}\frac{\sin15}{\cos30}$





- 5. A bullet of mass 20 g has an initial speed of 1 ms^{-1} , just before it starts penetrating a mud wall of thickness 20 cm. If the wall offers a mean resistance of 2.5×10^{-2} N, the speed of the bullet after emerging from the other side of the wall is close to :
 - (1) 0.4 ms^{-1} (2) 0.7 ms^{-1} (3) 0.3 ms^{-1} (4) 0.1 ms^{-1}

Answer (2)

Sol.
$$v^2 = u^2 - 2aS$$

$$v^{2} = (1)^{2} - (2) \left[\frac{2.5 \times 10^{-2}}{20 \times 10^{-3}} \right] \frac{20}{100}$$
$$v^{2} = 1 - \frac{1}{2}$$
$$\Rightarrow \quad v = \frac{1}{\sqrt{2}} m/s = 0.7 m/s$$

6. The time dependence of the position of a particle of mass m = 2 is given by $\vec{r}(t) = 2t\hat{i} - 3t^2\hat{j}$. Its angular momentum, with respect to the origin, at time t = 2 is :

(1)
$$-34(\hat{k}-\hat{i})$$
 (2) $48(\hat{i}+\hat{j})$
(3) $36\hat{k}$ (4) $-48\hat{k}$

Answer (4)

Sol. $\vec{r} = 2t\hat{i} - 3t^2\hat{j}$

$$\vec{v} = \frac{d\vec{r}}{dt} = 2\hat{i} - 6\hat{t}$$

$$\vec{a} = \frac{dv}{dt} = -6\hat{j}$$

 $\vec{F} = m\vec{a} = -12\hat{j}$

 $\vec{L} = m(\vec{r} \times \vec{v}) = 2(4\hat{i} - 12\hat{j}) \times (2\hat{i} - 12\hat{j}) = -48\hat{k}$

7. The figure represents a voltage regulator circuit using a Zener diode. The breakdown voltage of the Zener diode is 6 V and the load resistance is, $R_L = 4 \text{ k}\Omega$. The series resistance of the circuit is $R_i = 1 \text{ k}\Omega$. If the battery voltage V_B varies from 8 V to 16 V, what are the minimum and maximum values of the current through Zener diode?



(1) 0.5 mA; 6 mA
(2) 0.5 mA; 8.5 mA
(3) 1.5 mA; 8.5 mA
(4) 1 mA; 8.5 mA

Answer (2)



8. A metal coin of mass 5 g and radius 1 cm is fixed to a thin stick AB of negligible mass as shown in the figure. The system is initially at rest. The constant torque, that will make the system rotate about AB at 25 rotations per second in 5 s, is closed to :

(1)
$$4.0 \times 10^{-6}$$
 Nm (2) 7.9×10^{-6} Nm (3) 2.0×10^{-5} Nm (4) 1.6×10^{-5} Nm Answer (3)



$$\Rightarrow 25 \times 2\pi = (\alpha)5$$

$$\alpha = 10\pi$$

$$\Rightarrow \tau = \left(\frac{5}{4} \text{mR}^2\right)\alpha$$

$$\approx \left(\frac{5}{4}\right)(5 \times 10^{-3})(10^{-4})10\pi$$

$$= 2.0 \times 10^{-5} \text{Nm}$$

9. In a Young's double slit experiment, the ratio of the slit's width is 4 : 1. The ratio of the intensity of maxima to minima, close to the central fringe on the screen, will be :

(1)
$$(\sqrt{3}+1)^4$$
: 16 (2) 4: 1
(3) 25: 9 (4) 9: 1

Answer (4)

Sol.
$$I_1 = 4 I_0$$

$$I_2 = I_0$$

$$\frac{I_{max}}{I_{min}} = \frac{\left(\sqrt{I_1} + \sqrt{I_2}\right)^2}{\left(\sqrt{I_1} - \sqrt{I_2}\right)^2} = \left(\frac{9}{1}\right)^2$$

10. Two radioactive substances A and B have decay constants 5λ and λ respectively. At t = 0, a sample has the same number of the two nuclei. The time taken for the ratio of the

number of nuclei to become $\left(\frac{1}{e}\right)^2$ will be :

 $\frac{1}{4\lambda}$

 $\frac{2}{\lambda}$

(1)
$$\frac{1}{2\lambda}$$
 (2)
(3) $\frac{1}{\lambda}$ (4)

Answer (1)

Sol.
$$N_x(at t) = N_0 e^{-5\lambda t}$$

 $N_y(at t) = N_0 e^{-\lambda t}$
 $\frac{N_x}{N_y} = \frac{1}{e^2} = e^{-4\lambda t}$
 $\Rightarrow 4\lambda t = 2$
 $\Rightarrow t = \frac{2}{4\lambda} = \left(\frac{1}{2\lambda}\right)$

maximum weight that can be put on the block without fully submerging it under water?

[Take, density of water = 10^3 kg/m^3]

(1) 30.1 kg (2) 46.3 kg (3) 87.5 kg (4) 65.4 kg

Answer (3)

Sol. Given
$$(50)^3 \times \frac{30}{100} \times (1) \times g = M_{cube}g$$
 ...(i)

Let m mass should be placed

Hence
$$(50)^3 \times (1) \times g = (M_{cube} + m)g$$
 ...(ii)

equation (ii) - equation (i)

$$\Rightarrow$$
 mg = (50)³ × g(1 – 0.3) = 125 × 0.7 × 10³ g

12. The graph shows how the magnification m produced by a thin lens varies with image distance v. What is the focal length of the lens used?



Answer (4)

Sol. As the graph between magnification (m) and image distance (v) varies linearly, then

$$m = k_1 v + k_2$$

$$\Rightarrow \frac{v}{u} = k_1 v + k_2$$

$$\Rightarrow \frac{1}{u} = k_1 + \frac{k_2}{v}$$

$$\Rightarrow \frac{k_2}{v} - \frac{1}{u} = k_1$$
Clearly, $k_1 = \frac{1}{f}$ and $k_2 = 1$ here
$$\therefore f = \frac{1}{\text{slope of } m - v \text{ graph}} = \frac{b}{c}$$



ion gets deexcited to the ground state in all possible ways (including intermediate emissions), a total of six spectral lines are observed. What is the value of λ ?

(Given:
$$h = 6.63 \times 10^{-34} \text{ Js}; c = 3 \times 10^8 \text{ ms}^{-1}$$
)

- (1) 11.4 nm
- (2) 12.3 nm
- (3) 9.4 nm
- (4) 10.8 nm

1- -

Answer (4)

Sol.
$$\Delta \mathbf{E} = \frac{\mathbf{nc}}{\lambda}$$

= $(13.4)(3)^2 \left[1 - \frac{1}{16} \right] \mathbf{eV}$
 $\Rightarrow \lambda = \frac{1242 \times 16}{(13.4) \times (9)(15)} \mathbf{nm} \approx 10.8 \mathbf{nm}$

14. Two blocks A and B of masses $m_A = 1$ kg and $m_B = 3$ kg are kept on the table as shown in figure. The coefficient of friction between A and B is 0.2 and between B and the surface of the table is also 0.2. The maximum force F that can be applied on B horizontally, so that the block A does not slide over the block B is:

[Take g = 10 m/s²]
(1) 40 N
(2) 12 N
(3) 16 N
(4) 8 N
Answer (3)
Sol.
$$a_c = \left(\frac{F-f}{M+m}\right)$$

 $a = \frac{F-(0.2)4 \times 10}{4} = \left(\frac{F-8}{4}\right)$
We have $\frac{F-8}{4} \le (0.2)10$
 $\Rightarrow F-8 \le 8$
 $\Rightarrow F \le 16$

same charge and mass 4 μ g is kept at a distance of 1 mm from P. If B is released, then its velocity at a distance of 9 mm from P is :

$$\begin{bmatrix} \text{Take } \frac{1}{4\pi \in_0} = 9 \times 10^9 \text{ Nm}^2 \text{C}^{-2} \end{bmatrix}$$
(1) 2.0 × 10³ m/s
(2) 3.0 × 10⁴ m/s
(3) 1.5 × 10² m/s
(4) 1.0 m/s

Answer (Bonus)

$$\begin{array}{c|c} 1 \mu C & 1 mm \\ \hline A & B & B' & = 4 \mu gram \\ \hline (Fixed) & U_i = \frac{kq_1q_2}{r_i} & U_f = \frac{kq_1q_2}{r_i} \end{array}$$

Conservation of energy

$$\frac{kq_1q_2}{r_1} = \frac{kq_1q_2}{r_2} + \frac{1}{2}mv^2$$

$$v^2 = \frac{2kq_1q_2}{m} \left[\frac{1}{r_1} - \frac{1}{r_2}\right]$$

$$= \frac{2 \times 9 \times 10^9 \times 10^{-12}}{4 \times 10^{-9} \times 10^{-3}} \left[1 - \frac{1}{9}\right] = 4 \times 10^9$$

$$v = \sqrt{40} \times 10^4 \text{ m/s} = 6.32 \times 10^4 \text{ m/s}$$

16. Water from a tap emerges vertically downwards with an initial speed of 1.0 ms^{-1} . The cross-sectional area of the tap is 10^{-4} m^2 . Assume that the pressure is constant throughout the stream of water and that the flow is streamlined. The cross-sectional area of stream, 0.15 m below the tap would be:

(Take g = 10 ms⁻²)
(1)
$$1 \times 10^{-5} m^2$$
 (2) $5 \times 10^{-5} m^2$
(3) $5 \times 10^{-4} m^2$ (4) $2 \times 10^{-5} m^2$

Answer (2)

Sol. Using Bernoullie's equation $v_2 = \sqrt{v_1^2 + 2gh}$

Equation of continuity

 $A_1V_1 = A_2V_2$

$$(1 \text{ cm}^2)(1 \text{ m/s}) = (A_2) \left(\sqrt{(1)^2 + 2 \times 10 \times \frac{15}{100}} \right)$$

$$\Rightarrow A_2(\ln cm^2) = \frac{1}{2}$$
$$\Rightarrow A_2 = 5 \times 10^{-5} m^2$$



temperature increases by ΔT . The heat required to produce the same change in temperature, at a constant pressure is:

(1)
$$\frac{3}{2}Q$$
 (2) $\frac{2}{3}Q$
(3) $\frac{7}{5}Q$ (4) $\frac{5}{3}Q$

Answer (3)

Sol. Heat supplied at constant volume

$$Q = nC_V \Delta T$$

and heat supplied at constant pressure

$$Q_{1} = nC_{p}\Delta I$$

$$\therefore \quad \frac{Q_{1}}{Q} = \frac{C_{p}}{C_{v}}$$

$$\Rightarrow \quad Q_{1} = (Q)\left(\frac{7}{5}\right)$$

18. A simple pendulum of length L is placed between the plates of a parallel plate capacitor having electric field E, as shown in figure. Its bob has mass m and charge q. The time period of the pendulum is given by:



(1)
$$2\pi \sqrt{\frac{L}{\left(g - \frac{qE}{m}\right)}}$$
 (2) $2\pi \sqrt{\frac{L}{\left(g + \frac{qE}{m}\right)}}$

2π

(3)
$$2\pi \sqrt{\frac{L}{\sqrt{g^2 + \left(\frac{qE}{m}\right)^2}}}$$
 (4)

$$\begin{array}{l} \sqrt{g_{eff}} \\ \Rightarrow \quad g_{eff} = \sqrt{g^2 + \left(\frac{gE}{m}\right)^2} \\ \Rightarrow \quad t = 2\pi \sqrt{\frac{L}{\sqrt{g^2 + \left(\frac{qE}{m}\right)^2}}} \\ \end{array}$$

19. A source of sound S is moving with a velocity of 50 m/s towards a stationary observer. The observer measures the frequency of the source as 1000 Hz. What will be the apparent frequency of the source when it is moving away from the observer after crosssing him? (Take velocity of sound in air is 350 m/s)

(1) 857 Hz	(2) 1143 Hz
------------	-------------

(3) 807 Hz (4) 750 Hz

Answer (4)

Sol.
$$f_{app} = f_{act} \left(\frac{V \pm V_0}{V \mp V_s} \right)$$

$$1000 = f_{act} \left(\frac{350 - 0}{350 + (-50)} \right) \text{ and } f' = f_{act} \left(\frac{350}{350 + 50} \right)$$

$$\Rightarrow f_{act} = \frac{1000 \times 300}{400}$$

f_{act} ≈750Hz

20. A 2 mW laser operates at a wavelength of 500 nm. The number of photons that will be emitted per second is:

[Given Planck's constant h = 6.6×10^{-34} Js, speed of light c = 3.0×10^8 m/s]

- (1) 2 × 10¹⁶
- (2) 1.5 × 10¹⁶
- (3) 1 × 10¹⁶
- (4) 5 × 10¹⁵

Answer (4)

Sol.
$$E = \frac{hc}{\lambda}$$

Let no. of photons per sec. is N

$$\Rightarrow N\frac{hc}{\lambda} = 2 mW$$

$$\Rightarrow N = \frac{2 \times \lambda}{hC} = \frac{2 \times 5000 \times 10^{-3}}{12420 \times 1.6 \times 10^{-19}}$$

$$N = 5 \times 10^{15}$$



medium of resistivity $\rho.$ The resistance between the two spheres will be:

(1)
$$\frac{\rho}{4\pi} \left(\frac{1}{a} - \frac{1}{b} \right)$$
 (2) $\frac{\rho}{2\pi} \left(\frac{1}{a} - \frac{1}{b} \right)$
(3) $\frac{\rho}{2\pi} \left(\frac{1}{a} + \frac{1}{b} \right)$ (4) $\frac{\rho}{4\pi} \left(\frac{1}{a} + \frac{1}{b} \right)$

Answer (1)



$$= \left(\frac{\rho}{4\pi}\right) = \int_{a}^{b} \frac{dx}{x^{2}}$$
$$= \left(\frac{\rho}{4\pi}\right) \cdot \left(\frac{1}{2} - \frac{1}{b}\right)$$

22. In an experiment, brass and steel wires of length 1 m each with areas of cross section 1 mm^2 are used. The wires are connected in series and one end of the combined wire is connected to a rigid support and other end is subjected to elongation. The stress required to produce a net elongation of 0.2 mm is,

[Given, the Young's Modulus for steel and brass are, respectively, $120 \times 10^9 \text{ N/m}^2$ and $60 \times 10^9 \text{ N/m}^2$]

(1) 1.8 × 10 ⁶ N/m ²	(2) 1.2 × 10 ⁶ N/m ²
(3) 4.0 × 10 ⁶ N/m ²	(4) 0.2 × 10 ⁶ N/m ²

Answer (Bonus)



$$=8\times10^{6}\frac{N}{m^{2}}$$

(Answer is not matching)

23. A coil of self inductance 10 mH and resistance 0.1 Ω is connected through a switch to a battery of internal resistance 0.9 Ω . After the switch is closed, the time taken for the current to attain 80% of the saturation value is: [take ln5 = 1.6]

(1)	0.002 s	(2)	0.324 s

(3) 0.103 s (4) 0.016 s

Answer (4)

Sol.
$$I = I_{sat} \left(1 - e^{-\frac{Rt}{L}} \right)$$
 Here $R = R_L + r = 1\Omega$
 $0.8I_{sat} = I_{sat} \left(1 - e^{-\frac{t}{.01}} \right)$
 $\Rightarrow \frac{4}{5} = 1 - e^{-100t}$
 $\Rightarrow e^{-100t} = \left(\frac{1}{5} \right)$
 $\Rightarrow 100t = \ln 5$
 $\Rightarrow t = \frac{1}{100} \ln 5$
 $= 0.016 \sec c$

24. The magnitude of the magnetic field at the center of an equilateral triangular loop of side 1 m which is carrying a current of 10 A is:

[Take $\mu_0 = 4\pi \times 10^{-7} \text{ NA}^{-2}$] (1) 18 μ T (2) 1 μ T (3) 3 μ T (4) 9 μ T

Answer (1)





$$= \frac{3\mu_0 I}{4\pi \left(\frac{a}{2\sqrt{3}}\right)} = (2) \left(\frac{\sqrt{3}}{2}\right) \qquad d = \left(\frac{1}{3}\right) (a \sin 60)$$
$$= \frac{9}{2} \left(\frac{\mu_0 I}{\pi a}\right) \qquad d = \frac{a}{3} \times \frac{\sqrt{3}}{2}$$
$$= \frac{9 \times 2 \times 10^{-7} \times 10}{1} \qquad = \left(\frac{a}{2\sqrt{3}}\right)$$
$$= 18 \ \mu T$$

25. A square loop is carrying a steady current I and the magnitude of its magnetic dipole moment is m. If this square loop is changed to a circular loop and it carries the same current, the magnitude of the magnetic dipole moment of circular loop will be :

(1)
$$\frac{2 \text{ m}}{\pi}$$
 (2) $\frac{4 \text{ m}}{\pi}$
(3) $\frac{\text{m}}{\pi}$ (4) $\frac{3 \text{ m}}{\pi}$

Answer (2)

Sol.

$$I \xrightarrow{a} \rightarrow r = \left(\frac{2a}{\pi}\right)$$

$$m = (I) a^{2}$$

$$m_{1} = (I) (\pi) \left(\frac{4a^{2}}{\pi^{2}}\right)$$

$$m_{1} = \frac{4Ia^{2}}{\pi}$$

$$m_{1} = \frac{4m}{\pi}$$

- 26. A submarine experiences a pressure of 5.05×10^6 Pa at a depth of d₁ in a sea. When it goes further to a depth of d₂, it experiences a pressure of 8.08×10^6 Pa. Then d₂ d₁ is approximately (density of water = 10^3 kg/m³ and acceleration due to gravity = 10 ms^{-2}):
 - (1) 600 m (2) 500 m

(3) 300 m (4) 400 m

 $\Rightarrow \Delta H \simeq 300 \text{ m}$

- 27. The elastic limit of brass is 379 MPa. What should be the minimum diameter of a brass rod if it is to support a 400 N load without exceeding its elastic limit?
 - (1) 0.90 mm
 - (2) 1.16 mm
 - (3) 1.00 mm
 - (4) 1.36 mm

Answer (2)

Sol. Stress =
$$\frac{400}{\pi r^2} \le 379 \times 10^6 \text{ N/m}^2$$

$$\Rightarrow r^2 \ge \frac{400}{379 \times 10^6 \pi}$$

 $2r \ge 1.15 \text{ mm}$

28. Light is incident normally on a completely absorbing surface with an energy flux of 25 Wcm⁻². If the surface has an area of 25 cm², the momentum transferred to the surface in 40 min time duration will be :

Answer (3)

Sol. P =
$$\frac{\Delta E}{C}$$

= $\frac{(25 \times 25) \times 40 \times 60}{3 \times 10^8}$ N-s
= 5 × 10⁻³ N-s

29. A solid sphere of mass M and radius R is divided into two unequal parts. The first part

has a mass of $\frac{7 \text{ M}}{8}$ and is converted into a

uniform disc of radius 2R. The second part is converted into a uniform solid sphere. Let I_1 be the moment of inertia of the disc about its axis and I_2 be the moment of inertia of the new sphere about its axis. The ratio I_1/I_2 is given by :

- (1) 140 (2) 185
- (3) 285 (4) 65

Answer (1)





- 30. In the formula X = 5YZ², X and Z have dimensions of capacitance and magnetic field, respectively. What are the dimensions of Y in SI units ?
- (c) [m 2 7 A] Answer (4) Sol. X = 5YZ² $Y \propto \frac{X}{Z^2}$ $X = C = \frac{Q^2}{E} = \frac{[A^2T^2]}{[ML^2T^{-2}]}$ $X = [M^{-1}L^{-2}T^4A^2]$ $Z = B = \frac{F}{IL}$ $Z = [MT^{-2}A^{-1}]$ $Y = \frac{[M^{-1}L^{-2}T^4A^2]}{[MT^{-2}A^{-1}]^2}$ $Y = [M^{-3}L^{-2}T^8A^4]$

PART-B : CHEMISTRY

1. The correct statement is :

- (1) Zincite is a carbonate ore.
- (2) Zone refining process is used for the refining of titanium.
- (3) Aniline is a froth stabilizer.
- (4) Sodium cyanide cannot be used in the metallurgy of silver.

Answer (3)

- Sol. Ti is refined by Van Arkel method. Ag is leached by dilute solution of NaCN. Zincite is ZnO. Aniline is a froth stabilizer.
- 2. The pH of a 0.02 M NH₄Cl solution will be [given $K_b(NH_4OH) = 10^{-5}$ and log2 = 0.301]
 - (1) 2.65
 - (2) 5.35
 - (3) 4.35
 - (4) 4.65
- Answer (2)

Sol.
$$NH_4^+ + H_2O \implies NH_4OH + H^+$$

0.02 - x x x
$$K_{h} = \frac{10^{-14}}{10^{-5}} = 10^{-9}$$

 ≈ 0.02

$$K_{h} = \frac{x^{2}}{0.02}$$

$$10^{-9} \times 2 \times 10^{-2} = x^{2}$$

$$x = \sqrt{20} \times 10^{-6}$$

$$pH = -\log(\sqrt{20} \times 10^{-6})$$

$$pH = 5.35$$

- 3. The INCORRECT statement is :
 - (1) The spin-only magnetic moments of $[Fe(H_2O)_6]^{2+}$ and $[Cr(H_2O)_6]^{2+}$ are nearly similar.
 - (2) The gemstone, ruby, has Cr³⁺ ions occupying the octahedral sites of beryl.
 - (3) The spin-only magnetic moment of $[Ni(NH_3)_4(H_2O)_2]^{2+}$ is 2.83 BM.
 - (4) The color of $[CoCl(NH_3)_5]^{2+}$ is violet as it absorbs the yellow light.

Answer (2)

Sol. Ruby is aluminium oxide (AI_2O_3) containing about 0.5 – 1% Cr³⁺ ions which are randomly distributed in the position normally occupied by AI^{3+} ions.



- (1) 5 & 5(2) 4 & 4
- (3) 5 & 4
- (4) 4 & 5



4 stereogenic centres



5 stereogenic centres

- 5. The difference between ΔH and ΔU ($\Delta H \Delta U$), when the combustion of one mole of heptane(I) is carried out at a temperature T, is equal to :
 - (1) -3RT
 - (2) 4RT
 - (3) 3RT
 - (4) –4RT

Answer (4)

Sol.
$$C_7H_{16} + 11O_2 \xrightarrow{\Delta} 7CO_2 + 8H_2O_{(1)}$$

 $\Delta H - \Delta U = \Delta n_g RT$
 $\therefore \Delta n_g = -4$
 $\therefore \Delta H - \Delta U = -4RT$



Answer (3)

Sol. KCl is more conducting than NaCl



- 7. The crystal field stabilization energy (CFSE) of $[Fe(H_2O)_6]CI_2$ and $K_2[NiCI_4]$, respectively, are :
 - (1) –2.4 Δ_o and –1.2 Δ_t (2) –0.6 Δ_o and –0.8 Δ_t
 - (3) –0.4 Δ_{o} and –0.8 Δ_{t}

(4) –0.4
$$\Delta_o$$
 and –1.2 Δ_t

Answer (3)



 $230_2(g) + 0_2(g) - 230_3(g),$

 Δ H = -57.2 kJ mol⁻¹ and K_c = 1.7 × 10¹⁶.

Which of the following statement is INCORRECT?

- (1) The equilibrium constant is large suggestive of reaction going to completion and so no catalyst is required.
- (2) The addition of inert gas at constant volume will not affect the equilibrium constant.
- (3) The equilibrium will shift in forward direction as the pressure increases.
- (4) The equilibrium constant decreases as the temperature increases.

Answer (1)

Sol. $2SO_2(g) + O_2(g) \implies 2SO_3(g)$

 $K_c = 1.7 \times 10^{16}$ i.e. reaction goes to completion. Equilibrium constant has no relation with catalyst. Catalyst only affects the rate with which a reaction proceeds.

For the given reaction, catalyst V_2O_5 is used to speed up the reaction (Contact process).

- 9. A hydrated solid X on heating initially gives a monohydrated compound Y. Y upon heating above 373 K leads to an anhydrous white powder Z. X and Z, respectively are :
 - (1) Baking soda and dead burnt plaster.
 - (2) Baking soda and soda ash.
 - (3) Washing soda and soda ash.
 - (4) Washing soda and dead burnt plaster.

Answer (3)

Sol.
$$\operatorname{Na_2CO_3} \cdot 10H_2O \longrightarrow \operatorname{Na_2CO_3} \cdot H_2O + 9H_2O$$

 (X)
 $\operatorname{Na_2CO_3} \cdot H_2O \longrightarrow \operatorname{Na_2CO_3} + H_2O$
 (Y)
 (Z)
 (Z)
 (Z)

- X = Washing soda
- Z = Soda ash
- 10. Which of these factors does not govern the stability of a conformation in acyclic compounds?
 - (1) Angle strain
 - (2) Steric interactions
 - (3) Electrostatic forces of interaction
 - (4) Torsional strain

- 11. The highest possible oxidation states of uranium and plutonium, respectively, are :
 - (1) 6 and 7 (2) 7 and 6
 - (3) 6 and 4 (4) 4 and 6

Answer (1)

Sol. Maximum oxidation state shown by

Uranium = + 6

Plutonium = + 7

- 12. The correct option among the following is :
 - (1) Colloidal medicines are more effective because they have small surface area.
 - (2) Colloidal particles in lyophobic sols can be precipitated by electrophoresis.
 - (3) Brownian motion in colloidal solution is faster if the viscosity of the solution is very high.
 - (4) Addition of alum to water makes it unfit for drinking.

Answer (2)

- Sol. Electrophoresis is used to coagulate lyophobic colloids.
- 13. The major product obtained in the given reaction is :



AICI₃



are :

- (1) 20 and 3
- (2) 12 and 3
- (3) 12 and 4
- (4) 20 and 4

Answer (3)

Sol. Pentagons in C_{60} = 12

Triangles in $P_4 = 4$

15. For the reaction of H_2 with I_2 , the rate constant is 2.5 × 10⁻⁴ dm³ mol⁻¹ s⁻¹ at 327°C and 1.0 dm³ mol⁻¹ s⁻¹ at 527°C. The activation energy for the reaction, in kJ mol⁻¹ is :

(R = 8.314 J K⁻¹ mol⁻¹) (1) 150 (2) 59 (3) 72 (4) 166

Answer (4)

Sol. log
$$\frac{K_2}{K_1} = \frac{E_a}{2.303R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$

log $\frac{1}{2.5 \times 10^{-4}} = \frac{E_a}{8.314 \times 2.303} \left(\frac{1}{600} - \frac{1}{800} \right)$
 $3.6 = \frac{E_a}{8.314 \times 2.303} \times \frac{200}{600 \times 800}$
 $E_a = 165.4 \text{ kJ/mol}$
 $\approx 166 \text{ kJ/mol}$

16. The major product 'Y' in the following reaction is :





Answer (3)



18. 1 g of a non-volatile non-electrolyte solute is dissolved in 100 g of two different solvents A and B whose ebullioscopic constants are in the ratio of 1 : 5. The ratio of the elevation in their

boiling points,
$$\frac{\Delta T_{b}(A)}{\Delta T_{b}(B)}$$
, is :

(1) 1:5	(2) 10 : 1
(3) 5:1	(4) 1:0.2

Answer (1)

...

Sol.
$$\Delta T_b = k_b \times m$$

$$\frac{(k_b)_A}{(k_b)_B} = \frac{1}{5}$$

$$\frac{(\Delta I_b)_A}{(\Delta T_b)_B} = \frac{(K_b)_A}{(k_b)_B} = \frac{1}{5}$$



 $B(C_8H_6O_4)$. Anhydride of B is used for the preparation of phenolphthalein. Compound A is:



Answer (2)

Sol.



- 20. The ratio of the shortest wavelength of two spectral series of hydrogen spectrum is found to be about 9. The spectral series are :
 - (1) Paschen and Pfund
 - (2) Brackett and Pfund
 - (3) Lyman and Paschen
 - (4) Balmer and Brackett

Answer (3)

Sol. Shortest wavelength means $n_2 = \infty$

Lyman series
$$\overline{v}_{L} = \frac{1}{\lambda_{L}} = -1312 \times \frac{1}{1^{2}} \times 1^{2}$$

Paschen series $\overline{v}_{P} = \frac{1}{\lambda_{P}} = -1312 \times \frac{1}{3^{2}} \times 1^{2}$
 $\overline{v}_{L} = \lambda_{P} = 0$

$$\frac{1}{\overline{v}_{P}} = \frac{1}{\lambda_{L}} = 3$$

			item=1			
		(a)	High density polythene	(I)	Peroxide catalyst	
	-	(b)	Polyacrylonitrile	(II)	Condensation at high temperature and pressure	
	-	(c)	Novolac	(111)	Ziegler – Natta Catalyst	
		(d)	Nylon 6	(IV)	Acid or base catalyst	
(1) (a) $ ightarrow$ (III), (b) $ ightarrow$ (I), (c) $ ightarrow$ (IV), (d) $ ightarrow$ (II)						
	(2) (a) $ ightarrow$ (III), (b) $ ightarrow$ (I), (c) $ ightarrow$ (II), (d) $ ightarrow$ (IV)					
(3) (a) $ ightarrow$ (IV), (b) $ ightarrow$ (II), (c) $ ightarrow$ (I), (d) $ ightarrow$ (III)						
(4) (a) $ ightarrow$ (II), (b) $ ightarrow$ (IV), (c) $ ightarrow$ (I), (d) $ ightarrow$ (III)						
Ans	wei	r (1))			
Sol.	a.	HC	PE –	 Ziegler-Natta Catalyst 		
	b.	Ро	lyacrylonitrile	nitrile Peroxide Catalyst		

c. Novolac – Catalysed by acid or base d. Nylon-6 – Condensation at High T and P

22. Air pollution that occurs in sunlight is :

- (1) Fog (2) Oxidising smog
- (3) Acid rain (4) Reducing smog

Answer (2)

- Sol. Air pollution caused by sunlight is photochemical smog and it is oxidising.
- 23. The increasing order of nucleophilicity of the following nucleophiles is :
 - (a) $CH_3CO_2^{\odot}$ (b) H_2O
 - (c) $CH_3SO_3^{\odot}$ (d) OH
 - (1) (d) < (a) < (c) < (b)
 - (2) (b) < (c) < (d) < (a)
 - (3) (a) < (d) < (c) < (b)
 - (4) (b) < (c) < (a) < (d)

Answer (4)

Sol. Greater the negative charge Present on a nucleophilic centre greater would be its nucleophilicity.

$$OH > CH_3 - C \stackrel{70}{\swarrow} O = CH_3 - S \stackrel{1}{\longrightarrow} O = C$$



- (1) Mn < Ti < Zn < Ni (2) Ti < Mn < Zn < Ni
- (3) Ti < Mn < Ni < Zn (4) Zn < Ni < Mn < Ti

Answer (3)

Sol. Order for I.E. is

Ti < Mn < Ni < Zn

- 25. The noble gas that does NOT occur in the atmosphere is :
 - (1) Ne (2) Kr

(3) He (4) Ra

Answer (4)

- Sol. Radon is not present in atmosphere.
- 26. Which of the following is NOT a correct method of the preparation of benzylamine from cyanobenzene?

(ii) NaBH₄

- (1) (i) $SnCl_2 + HCl(gas)$ (ii) $NaBH_4$
- (2) H_2/Ni
- (3) (i) LiAlH_4 (ii) H_3O^+
- (4) (i) HCI/H₂O

Answer (4)



27. The minimum amount of $O_2(g)$ consumed per gram of reactant is for the reaction:

(Given atomic mass : Fe = 56, O = 16, Mg = 24, P = 31, C = 12, H = 1)

(1) $2Mg(s) + O_2(g) \rightarrow 2MgO(s)$

(2) 4Fe(s) +
$$3O_2(g) \rightarrow 2Fe_2O_3(s)$$

(3)
$$C_3H_8(g)$$
 + $5O_2(g) \rightarrow 3CO_2(g)$ + $4H_2O(I)$

(4)
$$P_4(s) + 5O_2(g) \rightarrow P_4O_{10}(s)$$

Answer (2)

Sol. (1) 2 Mg +
$$O_2 \longrightarrow 2$$
 MgO

1 g requires
$$\frac{32}{48}$$
 g = 0.66 g of O₂

- (3) $C_3H_8 + 5O_2 \longrightarrow 3CO_2 + 4H_2O$ 1 g of C_3H_8 requires = 3.6 g of O_2
- (4) $P_4 + 5O_2 \longrightarrow P_4O_{10}$ 1 g of P requires = 1.3 g of oxygen
- 28. Points I, II and III in the following plot respectively correspond to

(V_{mp} : most probable velocity)



From curve

$$\begin{aligned} (\mathsf{V}_{mp})_{\mathsf{I}} &< (\mathsf{V}_{mp})_{\mathsf{II}} < (\mathsf{V}_{mp})_{\mathsf{III}} \\ & \left(\mathsf{V}_{mp}\right)_{\mathsf{N}_{2}} \sim \sqrt{\frac{300}{28}}, \left(\mathsf{V}_{mp}\right)_{\mathsf{O}_{2}} \propto \sqrt{\frac{400}{32}}, \left(\mathsf{V}_{mp}\right)_{\mathsf{H}_{2}} \sim \sqrt{\frac{300}{2}} \\ & \therefore \quad \left(\mathsf{V}_{mp}\right)_{\mathsf{N}_{2}} < \left(\mathsf{V}_{mp}\right)_{\mathsf{O}_{2}} < \left(\mathsf{V}_{mp}\right)_{\mathsf{H}_{2}} \\ & \left(\mathsf{Under given Condition}\right) \end{aligned}$$



- (d) duffice hydrides produce h₂ gas when reacted with H₂O.
- (b) Reaction of LiAlH_4 with BF_3 leads to B_2H_6 .
- (c) PH_3 and CH_4 are electron rich and electron - precise hydrides, respectively.
- (d) HF and CH_4 are called as molecular hydrides.
- (1) (a), (c) and (d) only.
- (2) (c) and (d) only.
- (3) (a), (b) and (c) only.
- (4) (a), (b), (c) and (d).

Answer (4)

With water saline hydrides produce H₂ gas Sol. –

$$- \quad 3\text{LiAlH}_{4} + 4\text{BF}_{3} \longrightarrow 2\text{B}_{2}\text{H}_{6} + 3\text{LiF} + 3\text{AIF}_{3}$$

- HF and CH₄ are molecular hydrides
- 30. In chromatography, which of the following statements is INCORRECT for R_f?
 - (1) The value of R_f cannot be more than one.
 - (2) Higher R_f value means higher adsorption.
 - (3) R_f value is dependent on the mobile phase.
 - (4) R_f value depends on the type of chromatography.

Answer (2)

Sol. R_f represents retardation factor in chromatography.

Distance moved by the substance from base line $R_f = \frac{Distance moved by the solvent from baseline$

Higher R_f value means lower adsorpation

С

PART-C : MATHEMATICS

Let $f(x) = \log_{a}(\sin x)$, $(0 < x < \pi)$ and $g(x) = \sin^{-1}$ 1. Sol. .: A, B, C, are in A.P (e^{-x}), (x \ge 0). If α is a positive real number such \Rightarrow 2B = A + C that a = (fog)' (α) and b = (fog)(α), then : (1) $a\alpha^2 - b\alpha - a = 1$ $\Rightarrow B = \frac{\pi}{3}$ (2) $a\alpha^2 + b\alpha + a = 0$ (3) $a\alpha^2 - b\alpha - a = 0$ Area= $\frac{1}{2}(4x)\sin 60^{\circ}$ (4) $a\alpha^2 + b\alpha - a = -2\alpha^2$ Answer (1) $=\sqrt{3}x$ **Sol.** $f(x) = ln(sin x), g(x) = sin^{-1} (e^{-x})$ $f(g(x)) = ln(sin(sin^{-1}e^{-x}))$ Now $\cos 60^\circ = \frac{16 + x^2 - 3x^2}{8x}$ = -x $\Rightarrow -\alpha = b$ \Rightarrow 4x = 16 - 2x² $f'(g(\alpha)) = a$ x = 2 (as -4 is rejected) i.e.,a = -1 \therefore $a\alpha^2 - b\alpha + 1 = -\alpha^2 + \alpha^2 + 1 = -a$ Hence, area = $2\sqrt{3}$ sq. cm The angles A, B and C of a triangle ABC are in 2. 3. The sum of the real roots of the equation A.P. and a : b = 1 : $\sqrt{3}$. If c = 4 cm, then the area (in sq.cm) of this triangle is : 2 -3x x - 3 = 0, is equal to : (1) $\sqrt{3}$ -3 2x x+2 (1) 6 (2) $4\sqrt{3}$ (2) 0 (3) 2√3 (3) -4 (4) $\frac{1}{\sqrt{3}}$ (4) 1 Answer (3) Answer (2)



Sol.
$$\begin{vmatrix} 2 & -3x & x-3 \\ -3 & 2x & x+2 \end{vmatrix} = 0$$

 $\Rightarrow x(-3x^2 - 6x - 2x^2 + 6x) - 6(-3x + 9 - 2x - 4)$
 $-(4x - 9xA) = 0$
 $\Rightarrow x(-5x^2) - 6(-5x + 5) - 4x + 9x = 0$
 $\Rightarrow x^3 - 7x + 6 = 0$
All the roots are real

 \therefore Sum of real roots = $\frac{0}{1} = 0$

4. If z and w are two complex numbers such that

$$|zw| = 1 \text{ and } \arg(z) - \arg(w) = \frac{\pi}{2}, \text{ then }:$$
(1) $z\overline{w} = \frac{1-i}{\sqrt{2}}$
(2) $\overline{z}w = i$
(3) $z\overline{w} = \frac{-1+i}{\sqrt{2}}$
(4) $\overline{z}w = -i$

Answer (4)

Sol. |zw| = 1 ...(i)

$$\arg\left(\frac{z}{w}\right) = \frac{\pi}{2} \qquad \dots (ii)$$
$$\therefore \quad \frac{z}{w} + \frac{\overline{z}}{\overline{w}} = 0 \qquad \Rightarrow z\overline{w} = -\overline{z}w$$

from (i) $z\overline{z}w\overline{w} = 1$

$$(\overline{z}w)^2 = -1 \qquad \Rightarrow \overline{z}w = \pm i$$

from (ii)
$$-\arg(\overline{z}) - \arg w = \frac{\pi}{2}$$

$$\Rightarrow$$
 arg $(\overline{z}w) = \frac{-\pi}{2}$

Hence, $\overline{z}w = -i$

5. The locus of the centres of the circles, which touch the circle, $x^2 + y^2 = 1$ externally, also touch the y-axis and lie in the first quadrant, is :

(1)
$$y = \sqrt{1+2x}, x \ge 0$$
 (2) $x = \sqrt{1+4y}, y \ge 0$
(3) $x = \sqrt{1+2y}, y \ge 0$ (4) $y = \sqrt{1+4x}, x \ge 0$

Answer (1)

Sol. Let centre of required circle is (h, k).

$$\therefore 00' = r + r'$$

$$\Rightarrow \sqrt{h^2 + k^2} = 1 + h$$

h² + k² = 1 + h² + 2h
k² = 1 + 2h

Locus is $y = \sqrt{1+2x}$

6. The sum
$$1 + \frac{1^3 + 2^3}{1 + 2} + \frac{1^3 + 2^3 + 3^3}{1 + 2 + 3} + \dots$$

 $+ \frac{1^3 + 2^3 + 3^3 + \dots + 15^3}{1 + 2 + 3 + \dots + 15} - \frac{1}{2}(1 + 2 + 3 + \dots + 15)$
 is equal to :
(1) 1860 (2) 620
(3) 660 (4) 1240

Answer (2)

Sol.
$$S = 1 + \frac{1^3 + 2^3}{1 + 2} + \frac{1^3 + 2^3 + 3^3}{1 + 2 + 3} + \dots 15$$
 terms

$$T_{n} = \frac{1^{3} + 2^{3} + \dots n^{3}}{1 + 2 + \dots n} = \frac{\frac{n^{2} (n+1)^{2}}{4}}{\frac{n(n+1)}{2}} = \frac{n(n+1)}{2}$$

$$S = \frac{1}{2} \left(\sum_{n=1}^{15} n^2 + \sum_{n=1}^{15} n \right) = \frac{1}{2} \left(\frac{15(16)(31)}{6} + \frac{15(16)}{2} \right)$$
$$= 680$$

$$\Rightarrow 680 - \frac{1}{2} \frac{15(16)}{2} = 680 - 60 = 620$$

7. The smallest natural number n, such that the coefficient of x in the expansion of $\left(x^2 + \frac{1}{x^3}\right)^n$ is ${}^nC_{23}$, is : (1) 58 (2) 35 (3) 38 (4) 23

Answer (3)



General term $T_{r+1} = {}^{n}C_{r} (x^{2})^{n-r} (\frac{1}{x^{3}})^{r}$ ${}^{n}C_{r} \cdot x^{2n-5r}$ for coefficiant of x, 2n - 5r = 1 Given ${}^{n}C_{r} = {}^{n}C_{23}$ r = 23 or n - r = 23 $\Rightarrow n = 58$ or n = 38Minimum value is n = 38

8. If both the mean and the standard deviation of 50 observations $x_1, x_2, \dots x_{50}$ are equal to 16, then the mean of $(x_1 - 4)^2, (x_2 - 4)^2, \dots (x_{50} - 4)^2$ is :

(1)	380	(2)	480
(3)	400	(4)	525

Answer (3)

Sol.
$$\frac{x_1 + x_2 + \dots x_{50}}{50} = 16$$
$$16^2 = \frac{x_1^2 + x_2^2 \dots x_{50}^2}{50} - 16^2$$
$$2(16)^2 50 = x_1^2 + x_2^2 + \dots x_{50}^2$$
Required mean =
$$\frac{(x_1 - 4)^2 + (x_2 - 4)^2 + \dots (x_{50} - 4)^2}{50}$$
$$= \frac{16^2 (100) + 4^2 (50) - 8(16 \times 50)}{50}$$
$$= 16^2 (2) + 16 - 8(16) = 400$$

9. If $\int x^5 e^{-x^2} dx = g(x)e^{-x^2} + c$, where c is a constant of integration, then g(-1) is equal to :

(1) -1 (2)
$$-\frac{1}{2}$$

(3) 1 (4) $-\frac{5}{2}$

Answer (4)

Sol.
$$I = \int x^5 \cdot e^{-x^2} dx$$

Put $-x^2 = t \implies -2xdx = dt$
 $I = \int \frac{t^2 \cdot e^t dt}{(-2)} = \frac{-1}{2} e^t (t^2 - 2t + 2) + c$
 $\therefore \quad g(x) = \frac{-1}{2} (x^4 + 2x^2 + 2)$
 $g(-1) = \frac{-5}{2}$

where a \neq 0 and 0 < r $\leq \frac{1}{2}$. If 3a, 7b and 15c are the first three terms of an A.P., then the 4th term of this A.P. is :

(1)
$$\frac{2}{3}a$$
 (2) a
(3) $\frac{7}{3}a$ (4) 5a

Answer (2)

- Sol. Let b = ar, c = ar² AP : 3a, 7ar, 15ar² 14ar = 3a + 15ar² \Rightarrow 15r² - 14r + 3 = 0 \Rightarrow r = $\frac{1}{3}$ or $\frac{3}{5}$ (rejected) Fourth term = 15ar² + 7ar - 3a = a(15r² + 7r - 3) = a($\frac{15}{9} + \frac{7}{3} - 3$)
 - = a
- 11. The area (in sq. units) of the region bounded by the curves $y = 2^x$ and y = |x + 1|, in the first quadrant is :

(1)
$$\frac{3}{2} - \frac{1}{\log_e 2}$$
 (2) $\frac{1}{2}$
(3) $\log_e 2 + \frac{3}{2}$ (4) $\frac{3}{2}$

Answer (1)





line $\frac{1}{2} = \frac{1}{-1} = \frac{1}{1}$ to the plane x + y + z = 3 such that the foot of the perpendicular Q also lies on the plane x - y + z = 3. Then the co-ordinates of Q are :

(1) (1, 0, 2) (2) (2, 0, 1)(4) (-1, 0, 4)(3) (4, 0, -1)

Answer (2)

Sol.
$$P(2\lambda + 1, \overline{p}^{\lambda - 1, \lambda})$$

 $< 1, 1, 1 >$
d.r's of normal to
 $x + y + z = 3$
 $Q_{(\alpha, \beta, \gamma)}$

Let Q be (α, β, γ)

$$\alpha + \beta + \gamma = 3 \qquad \dots(i)$$

$$\alpha - \beta + \gamma = 3 \qquad \dots(ii)$$

$$\therefore \quad \alpha + \gamma = 3 \text{ and } \beta = 0$$

Equating DB's of PQ :

Equating DR's of PQ :

$$\frac{\alpha - 2\lambda - 1}{1} = \frac{\lambda + 1}{1} = \frac{\gamma - \lambda}{1}$$

$$\Rightarrow \alpha = 3\lambda + 2, \gamma = 2\lambda + 1$$

Substituting in equation (i) we get

 \Rightarrow 5 λ + 3 = 3

 $\lambda = 0$

Point is Q(2, 0, 1)

13. If the tangent to the curve $y = \frac{x}{x^2 - 3}$, $x \in \mathbb{R}$, $(\mathbf{x} \neq \pm \sqrt{3})$, at a point $(\alpha, \beta) \neq (0, 0)$ on it is parallel to the line 2x + 6y - 11 = 0, then : (1) $|6\alpha + 2\beta| = 19$ (2) $|2\alpha + 6\beta| = 19$ (3) $|6\alpha + 2\beta| = 9$ (4) $|2\alpha + 6\beta| = 11$

Answer (1)

Sol.
$$y = \frac{x}{x^2 - 3}$$

$$\frac{dy}{dx} = \frac{(x^2 - 3) - x(2x)}{(x^2 - 3)^2} = \frac{-x^2 - 3}{(x^2 - 3)^2}$$
$$\frac{dy}{dx}\Big|_{(\alpha, \beta)} = \frac{-\alpha^2 - 3}{(\alpha^2 - 3)^2} = -\frac{1}{3}$$
$$3(\alpha^2 + 3) = (\alpha^2 - 3)^2 \qquad \dots (i)$$

Also,
$$\beta = \frac{\alpha}{\alpha^2 - 3} \Rightarrow \alpha^2 - 3 = \frac{\alpha}{\beta} \Rightarrow \frac{\alpha}{\beta} = 6$$

 $\Rightarrow \alpha = \pm 3, \beta = \pm \frac{1}{2}$

Which satisfies $|6\alpha + 2\beta| = 19$

14. Let y = y(x) be the solution of the differential equation, $\frac{dy}{dx} + y \tan x = 2x + x^2 \tan x, x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, such that y(0) = 1. Then : (1) $\mathbf{y}\left(\frac{\pi}{4}\right) + \mathbf{y}\left(-\frac{\pi}{4}\right) = \frac{\pi^2}{2} + 2$ (2) $y\left(\frac{\pi}{2}\right) - y\left(-\frac{\pi}{2}\right) = \sqrt{2}$

(2)
$$\mathbf{y}'\left(\frac{\pi}{4}\right) + \mathbf{y}'\left(-\frac{\pi}{4}\right) = -\sqrt{2}$$

(3) $\mathbf{y}'\left(\frac{\pi}{4}\right) + \mathbf{y}'\left(-\frac{\pi}{4}\right) = -\sqrt{2}$
(4) $\mathbf{y}'\left(\frac{\pi}{4}\right) - \mathbf{y}'\left(-\frac{\pi}{4}\right) = \pi - \sqrt{2}$

Answer (4)

Sol.
$$\frac{dy}{dx} + y \tan x = 2x + x^{2} \tan x$$

$$P = \tan x, Q = 2x + x^{2} \tan x$$

$$I.F. = e^{\int \tan x dx} = e^{\ln|\sec x|} = |\sec x|$$

$$y(\sec x) = \int (2x + x^{2} \tan x) \sec x dx$$

$$= \int x^{2} \tan x \sec x dx + \int 2x \sec x dx$$

$$= x^{2} \sec x - \int 2x \sec x dx + \int 2x \sec x dx$$

$$= x^{2} \sec x + c$$

$$As \ y(0) = 1, \ c = 1$$

$$\therefore \ y = x^{2} + \cos x$$

$$At \ x = \frac{\pi}{4}, \quad y\left(\frac{\pi}{4}\right) = \frac{\pi^{2}}{16} + \frac{1}{\sqrt{2}}$$

$$y\left(-\frac{\pi}{4}\right) = \frac{\pi^{2}}{16} + \frac{1}{\sqrt{2}}$$

$$y\left(\frac{\pi}{4}\right) - y\left(-\frac{\pi}{4}\right) = 0$$

$$\frac{dy}{dx} = 2x - \sin x$$

$$y'\left(\frac{\pi}{4}\right) = \frac{\pi}{2} - \frac{1}{\sqrt{2}}, \ y'\left(-\frac{\pi}{4}\right) = -\frac{\pi}{2} + \frac{1}{\sqrt{2}}$$

 $\mathbf{y}'\left(\frac{\pi}{\mathbf{4}}\right) - \mathbf{y}'\left(-\frac{\pi}{\mathbf{4}}\right) = \pi - \sqrt{2}$

maximises the product $a_1 a_4 a_5$, is : (1) $\frac{2}{3}$ (2) $\frac{8}{5}$ (4) $\frac{6}{5}$ (3) $\frac{3}{2}$ Answer (1) Sol. a + 5d = 2 Let $A = a_1 a_4 a_5 = a(a + 3d)(a + 4d)$ = a(2 - 2d)(2 - d) $A = (2 - 5d)(4 - 6d + 2d^2)$ $\frac{dA}{dd} = 0$ $(2-5d)(-6+4d) + (4-6d+2d^2)(-5) = 0$ \Rightarrow 15d² - 34d + 16 = 0 $d = \frac{8}{5}, \frac{2}{2}$ For $d = \frac{2}{3}$, $\frac{d^2 A}{dd^2} < 0$ Hence $d = \frac{2}{3}$

16. If the plane 2x - y + 2z + 3 = 0 has the distances $\frac{1}{3}$ and $\frac{2}{3}$ units from the planes $4x - 2y + 4z + \lambda = 0$ and $2x - y + 2z + \mu = 0$, respectively, then the maximum value of $\lambda + \mu$ is equal to :

(1) 13	(2) 15
(3) 5	(4) 9

Answer (1)

Sol. $P_1 : 2x - y + 2z + 3 = 0$

$$P_{2}: 2x - y + 2z + \frac{\lambda}{2} = 0$$

$$P_{3}: 2x - y + 2z + \mu = 0$$
Given
$$\frac{1}{3} = \frac{\left|\frac{3 - \frac{\lambda}{2}}{\sqrt{9}}\right|}{\sqrt{9}} \Rightarrow \left|3 - \frac{\lambda}{2}\right| = 1$$

$$\lambda_{max} = 8$$
Also,
$$\frac{2}{3} = \frac{|\mu - 3|}{\sqrt{9}} \Rightarrow \mu_{max} = 5$$

$$(\lambda + \mu)_{max} = 13$$

(3) 1 (4) -7

Answer (4)

Sol.
$$\lim_{x \to 1} \frac{x^2 - ax + b}{x - 1} = 5$$

As limit is finite, 1 - a + b = 0

$$\Rightarrow \lim_{x \to 1} \frac{2x - a}{1} = 5 \qquad \left(\frac{0}{0} \text{ form}\right)$$

i.e., 2 - a = 5
or a = -3
 \therefore b = -4
a + b = -3 - 4 = -7

- 18. The number of real roots of the equation $5+|2^{x}-1|=2^{x}(2^{x}-2)$ is:
 - (1) 4
 (2) 2

 (3) 1
 (4) 3

Answer (3)

Sol. Let
$$2^{x} - 1 = t$$

 $5 + |t| = (t + 1) (t - 1)$
 $\Rightarrow |t| = t^{2} - 6$
For $t > 0, t^{2} - t - 6 = 0$
i.e., $t = 3$ or -2 (rejected)
For $t < 0, t^{2} + t - 6 = 0$
i.e., $t = -3$ or 2 (both rejected)
 $\therefore 2^{x} - 1 = 3$
 $\Rightarrow x = 2$

19. Lines are drawn parallel to the line 4x - 3y + 2= 0, at a distance $\frac{3}{5}$ from the origin. Then which one of the following points lies on any of these lines?

(1)
$$\left(\frac{1}{4}, -\frac{1}{3}\right)$$
 (2) $\left(\frac{1}{4}, \frac{1}{3}\right)$
(3) $\left(-\frac{1}{4}, \frac{2}{3}\right)$ (4) $\left(-\frac{1}{4}, -\frac{2}{3}\right)$

Answer (3)



Given $\frac{3}{5} = \left| \frac{\alpha}{5} \right|$

 $\Rightarrow \alpha = \pm 3$ Line is 4x - 3y + 3 = 0 or 4x - 3y - 3 = 0

Clearly
$$\left(-\frac{1}{4},\frac{2}{3}\right)$$
 satisfies $4x - 3y + 3 = 0$

20. Suppose that 20 pillars of the same height have been erected along the boundary of a circular stadium. If the top of each pillar has been connected by beams with the top of all its nonadjacent pillars, then the total number of beams is :

(1)	210	(2)	180
(3)	170	(4)	190

Answer (3)

Sol. Required number of beams = ${}^{20}C_2 - 20$

21. A spherical iron ball of radius 10 cm is coated with a layer of ice of uniform thickness that melts at a rate of $50 \text{ cm}^3/\text{min}$. When the thickness of the ice is 5 cm, then the rate at which the thickness (in cm/min) of the ice decreases, is :

(1)
$$\frac{5}{6\pi}$$
 (2) $\frac{1}{36\pi}$
(3) $\frac{1}{9\pi}$ (4) $\frac{1}{18\pi}$

Answer (4)

Sol.
$$\frac{dV_{ice}}{dt} = 50$$
$$V_{ice} = \frac{4}{3}\pi (10 + r)^3 - \frac{4}{3}\pi (10)^3$$
$$\frac{dV}{dt} = \frac{4}{3}\pi 3(10 + r)^2 \frac{dr}{dt}$$
$$= 4\pi (10 + r)^2 \frac{dr}{dt}$$
At r = 5, 50 = 4\pi (225) \frac{dr}{dt}
$$\frac{dr}{dt} = \frac{50}{4\pi (225)}$$
$$= \frac{1}{18\pi} \text{ cm/min}$$

one head is more than 99% is :

(1) 8	(2) 6
(3) 5	(4) 7

Answer (4)

Sol.
$$1 - \left(\frac{1}{2}\right)^n > \frac{99}{100}$$

 $\left(\frac{1}{2}\right)^n < \frac{1}{100}$
 $\therefore n \ge 7$

Minimum value is 7.

23. If 5x + 9 = 0 is the directrix of the hyperbola $16x^2 - 9y^2 = 144$, then its corresponding focus is :

(1)
$$\left(\frac{5}{3}, 0\right)$$
 (2) $\left(-\frac{5}{3}, 0\right)$
(3) (-5, 0) (4) (5, 0)

Answer (3)

i. e.
$$\frac{x^2}{9} - \frac{y^2}{16} = 1$$

$$S'_{1} = \frac{-9}{5}$$

a = 3, b = 4
$$e^{2} = 1 + \frac{16}{9} = \frac{25}{9}$$

S' = $\left(-3 \times \frac{5}{3}, 0\right) = (-5, 0)$

24. Let λ be a real number for which the system of linear equations

$$x + y + z = 6$$

 $4x + \lambda y - \lambda z = \lambda - 2$
 $3x + 2y - 4z = -5$

has infinitely many solutions. Then λ is a root of the quadratic equation :

(1)
$$\lambda^2 + 3\lambda - 4 = 0$$
 (2) $\lambda^2 - \lambda - 6 = 0$
(3) $\lambda^2 + \lambda - 6 = 0$ (4) $\lambda^2 - 3\lambda - 4 = 0$



$$\Delta = \begin{vmatrix} 1 & 1 & 1 & 1 \\ 3 & 2 & -4 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} 0 & 0 & 1 \\ 4 -\lambda & 2\lambda & -\lambda \\ 1 & 6 & -4 \end{vmatrix} = 0 \Rightarrow \lambda = 3$$

$$\Delta_{1} = \begin{vmatrix} 6 & 1 & 1 \\ \lambda -2 & \lambda & -\lambda \\ -5 & 2 & -4 \end{vmatrix} = 0 \quad \text{for } \lambda = 3$$

$$\Delta_{2} = \begin{vmatrix} 1 & 6 & 1 \\ 4 & \lambda -2 & -\lambda \\ 3 & -5 & -4 \end{vmatrix} = 0 \quad \text{for } \lambda = 3$$

$$\Delta_{3} = \begin{vmatrix} 1 & 1 & 6 \\ 4 & \lambda & \lambda -2 \\ 3 & 2 & -5 \end{vmatrix} = 0 \quad \text{for } \lambda = 3$$

$$\therefore \text{ For } \lambda = 3, \text{ infinitely many solutions is obtained.}$$
25. The integral $\int_{\frac{\pi}{6}}^{\frac{\pi}{5}} \sec^{\frac{2}{3}} x \csc^{\frac{4}{3}} x dx$ is equal to :
(1) $3^{\frac{7}{6}} - 3^{\frac{5}{6}}$
(2) $3^{\frac{5}{3}} - 3^{\frac{1}{3}}$
(3) $3^{\frac{5}{6}} - 3^{\frac{2}{3}}$
(4) $3^{\frac{4}{3}} - 3^{\frac{1}{3}}$
Answer (1)
Sol. $I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \sec^{\frac{2}{3}} x \csc^{\frac{4}{3}} x dx$

$$= \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{1 dx}{\cos^{\frac{2}{3}} x \sin^{\frac{4}{3}} x}{\frac{\pi}{6} \cos^{\frac{2}{3}} x \sin^{\frac{4}{3}} x}$$
Let $\tan x = t$

$$I = \int_{\frac{\pi}{3}}^{\frac{\pi}{3}} t^{-\frac{4}{3}} dt = \frac{3 [t^{\frac{1}{3}}]_{\frac{\sqrt{3}}{3}}}{-1}$$

x + y = 1 and $y = 4\sqrt{2x}$, then |c| is equal to .

(1)
$$\frac{1}{\sqrt{2}}$$

(2) $\frac{1}{2}$
(3) 2

(4) √<u>2</u>

Answer (4)

Sol. Tangent on $y^2 = 4\sqrt{2}x$ is $yt = x + \sqrt{2}t^2$

As it is tangent on circle also,

$$\left|\frac{\sqrt{2}t^2}{\sqrt{1+t^2}}\right| = 1$$

2t⁴ = 1 + t² i.e. t² = 1
Equation is $\pm y = x + \sqrt{2}$
Hence $|c| = \sqrt{2}$

27. The tangent and normal to the ellipse $3x^2 + 5y^2 = 32$ at the point P(2, 2) meet the x-axis at Q and R, respectively. Then the area (in sq. units) of the triangle PQR is :

(1)
$$\frac{16}{3}$$

(2) $\frac{14}{3}$
(3) $\frac{34}{15}$
(4) $\frac{68}{15}$

Answer (4)

Sol. For
$$\frac{3x^2}{32} + \frac{5y^2}{32} = 1$$

Tangent at P is
 $\frac{7}{32} + \frac{9}{32} = 1$
 $\frac{3(2)x}{32} + \frac{5(2)y}{32} = 1$



 $=-3(3^{-1/6}-3^{-1/6})$

 $=3(3^{\frac{1}{6}}-3^{-\frac{1}{6}})$

 $=3^{7/6}-3^{5/6}$

$$Q = \left(\frac{16}{3}, 0\right)$$
Normal at P is $\frac{32x}{3(2)} - \frac{32y}{5(2)} = \frac{32}{3} - \frac{32}{5}$

$$R = \left(\frac{4}{5}, 0\right)$$
area of $\triangle PQR = \frac{1}{2}$ (PQ) (PR) $= \frac{1}{2}\sqrt{\frac{136}{3}} \cdot \sqrt{\frac{136}{5}}$

$$= \frac{68}{15}$$
The distance of the point having position vector

- 28. The distance of the point having position vector $-\hat{i}+2\hat{j}+6\hat{k}$ from the straight line passing through the point (2, 3, -4) and parallel to the vector, $6\hat{i}+3\hat{j}-4\hat{k}$ is :
 - (1) 7 (2) $4\sqrt{3}$ (3) 6 (4) $2\sqrt{13}$

Answer (1)

Sol. Equation of I is
$$\frac{x-2}{6} = \frac{y-3}{3} = \frac{z+4}{-4}$$

P (-1, 2, 6)
 $(2, 3, -4)$ M < 6, 3, -4>
 $(6\lambda + 2, 3\lambda + 3, -4\lambda - 4)$
Let M ($6\lambda + 2, 3\lambda + 3, -4\lambda - 4$)
DR's of PM is < $6\lambda + 3, 3\lambda + 1, -4\lambda - 10 >$
 $\Rightarrow (6\lambda + 3)(6) + (3\lambda + 1)(3) + (-4\lambda - 10)(-4) =$
 $\Rightarrow \lambda = -1$
i.e. M = (-4, 0, 0)
 \therefore PM = $\sqrt{9+4+36} = 7$

- $x \le \frac{y}{2}$, then for all x, y, $4x^2 4xy \cos \alpha + y^2$ is equal to :
- **(1) 2 sin**²α
- (2) $4 \sin^2 \alpha 2x^2y^2$ (3) $4 \cos^2 \alpha + 2x^2y^2$
- (4) 4 sin²α

Answer (4)

Sol.
$$\cos^{-1}x - \cos^{-1}\frac{y}{2} = \alpha$$

$$\Rightarrow \cos^{-1}\left(\frac{xy}{2} + \sqrt{1 - x^2} \cdot \sqrt{1 - \frac{y^2}{4}}\right) = \alpha$$

$$\Rightarrow \frac{xy}{2} + \frac{\sqrt{1 - x^2}\sqrt{4 - y^2}}{2} = \cos\alpha$$

$$\Rightarrow xy + \sqrt{1 - x^2}\sqrt{4 - y^2} = 2\cos\alpha$$

$$(xy - 2\cos\alpha)^2 = (1 - x^2)(4 - y^2)$$

$$x^2y^2 + 4\cos^2\alpha - 4xy\cos\alpha = 4 - y^2 - 4x^2 + x^2y^2$$

$$4x^2 - 4xy\cos\alpha + y^2 = 4\sin^2\alpha$$

- 30. The negation of the Boolean expression ~ s \vee (~ r \wedge s) is equivalent to :
 - (1) s ∧ r
 (2) r
 (3) ~ s ∧ ~ r
 (4) s ∨ r

Answer (1)

Sol. $\sim s \lor (\sim r \land s)$ $\equiv (\sim s \lor \sim r) \land (\sim s \lor s)$ $\equiv (\sim s \lor \sim r) (\because (\sim s \lor s) \text{ is tautology})$ $\equiv \sim (s \land r)$ Hence its negation is $s \land r$

0



Answers & Solutions

for

JEE (MAIN)-2019 (Online) Phase-2

(Physics, Chemistry and Mathematics)

Time : 3 hrs.

M.M.: 360

Important Instructions :

- 1. The test is of **3 hours** duration.
- 2. The Test Booklet consists of **90** questions. The maximum marks are **360**.
- 3. There are *three* parts in the question paper A, B, C consisting of **Physics**, **Chemistry** and **Mathematics** having 30 questions in each part of equal weightage.
- 4. Each question is allotted 4 (four) marks for each correct response. ¼ (one-fourth) marks will be deducted for indicating incorrect response of each question. No deduction from the total score will be made if no response is indicated for an item in the answer sheet.
- 5. There is only one correct response for each question.



1. The value of numerical aperture of the objective lens of a microscope is 1.25. If light of wavelength 5000 Å is used, the minimum separation between two points, to be seen as distinct, will be :

(1)	0.24 μ m	(2)	0.38	μ m
(3)	0.48 μ m	(4)	0.12	μ m

Answer (1)

Sol.
$$\theta_{\min} = \frac{1.22\lambda}{D}$$

 $\frac{D}{2f} = 1.25$
 $d_{\min} = \frac{1.22\lambda f}{D} = \frac{1.22 \times 5000 \times 10^{-10}}{2.50}$
 $= 0.24 \ \mu m$

1.22λ

2. A circular disc of radius b has a hole of radius a at its centre (see figure). If the mass per unit

area of the disc varies as $\left(\frac{\sigma_0}{r}\right)$, then the radius of gyration of the disc about its axis passing through the centre is :



(1)
$$\frac{a+b}{2}$$
 (2) $\sqrt{\frac{a+b+ab}{2}}$
(3) $\frac{a+b}{3}$ (4) $\sqrt{\frac{a^2+b^2+ab}{3}}$

Answer (4)

Sol.
$$\Rightarrow \sigma = \frac{\sigma_0}{r} \qquad \therefore \frac{\sigma_0}{r} 2\pi r dr = dm$$

 $\Rightarrow m = \sigma_0 2\pi (b - a)$
 $I = \sigma_0 2\pi \int_a^b r^2 dr = \frac{2\pi \sigma_0}{3} (b^3 - a^3)$

$$\therefore \quad \mathbf{mk}^2 = \mathbf{I} \implies 2\pi\sigma_0(\mathbf{b} - \mathbf{a})\mathbf{k}^2 = \frac{2\pi\sigma_0}{3}(\mathbf{b}^3 - \mathbf{a}^3)$$
$$\implies \quad \mathbf{k}^2 = \frac{1}{3}(\mathbf{b}^2 + \mathbf{a}\mathbf{b} + \mathbf{a}^2)$$
$$\therefore \quad \mathbf{k} = \sqrt{\frac{1}{3}(\mathbf{b}^3 - \mathbf{a}^3)}(\mathbf{b}^3 - \mathbf{a}^3)$$

3. A shell is fired from a fixed artillery gun with an initial speed u such that it hits the target on the ground at a distance R from it. If t_1 and t_2 are the values of the time taken by it to hit the target in two possible ways, the product t_1t_2 is :

(1)
$$\frac{R}{2g}$$
 (2) $\frac{2R}{g}$
(3) $\frac{R}{g}$ (4) $\frac{R}{4g}$

Answer (2)

Sol. For same horizontal range.

$$\theta_{1} = \theta \qquad R = \frac{u^{2} \sin 2\theta}{g}$$

$$\theta_{2} = (90 - \theta)$$
so $t_{1} = \frac{2u \sin \theta}{g}$ and $t_{2} = \frac{2u \cos \theta}{g}$

$$\therefore \quad t_{1}t_{2} = \frac{u^{2} 4 \sin \theta \cos \theta}{g^{2}}$$

$$\Rightarrow \quad t_{1}t_{2} = \frac{2u^{2} \sin 2\theta}{g^{2}} = \frac{2R}{g}$$

4. The resistive network shown below is connected to a D.C. source of 16 V. The power consumed by the network is 4 Watt. The value of R is :





 $301. N_{eq} = 2N + N + 4N + N = 0N$

$$P = \frac{v^2}{R_{eq}} \Rightarrow \frac{16 \times 16}{8R} = 4$$
 watt

 $\frac{\mathbf{16}{\times}\mathbf{16}}{\mathbf{4}{\times}\mathbf{8}} \!=\! \mathbf{R} \,\Rightarrow\, \mathbf{R} \!=\! \mathbf{8}\,\Omega$

5. In a double slit experiment, when a thin film of thickness t having refractive index
$$\mu$$
 is introduced in front of one of the slits, the maximum at the centre of the fringe pattern shifts by one fringe width. The value of t is (λ is the wavelength of the light used) :

(1)
$$\frac{2\lambda}{(\mu-1)}$$

(2)
$$\frac{\lambda}{2(\mu-1)}$$

(3)
$$\frac{\lambda}{(2\mu-1)}$$

(4)
$$\frac{\lambda}{(\mu-1)}$$

Answer (4)

- Sol. $\mu t t = \lambda$ $\Rightarrow t(\mu - 1) = \lambda$ $\Rightarrow t = \frac{\lambda}{(\mu - 1)}$
- 6. A galvanometer of resistance 100 Ω has 50 divisions on its scale and has sensitivity of 20 μ A/division. It is to be converted to a voltmeter with three ranges, of 0-2 V, 0-10 V and 0-20 V. The appropriate circuit to do so is :

(1)
$$\begin{array}{|c|c|c|c|c|} \hline & & & & & & & & \\ \hline & & & & & \\ R_1 & & & & R_2 \\ \hline & & & & & \\ R_2 &= 8000 \ \Omega \\ R_3 &= 10000 \ \Omega \\ \hline & & & & \\ R_3 &= 10000 \ \Omega \\ \hline & & & & \\ R_1 & & & & \\ R_2 & & & \\ R_3 &= 1900 \ \Omega \\ \hline & & & & \\ R_3 &= 19900 \ \Omega \\ \hline & & & \\ 2 \ V & 10 \ V & 20 \ V \end{array}$$

Answer (1)

Sol. For
$$R_1$$

 $\therefore I_g = 10^{-3} A$
 $\therefore 10^{-3}(R_1 + 100) = 2 V \Rightarrow R_1 = 1900 \Omega$
For R_2
 $10^{-3}(R_1 + R_2 + 100) = 10 V$
 $\Rightarrow R_1 + R_2 + 100 = 10000$
 $\Rightarrow R_2 = 8000 \Omega$
For R_3
 $10^{-3}(R_1 + R_2 + R_3 + 100) = 20 V$
 $\Rightarrow R_1 + R_2 + R_3 + 100 = 20 \times 1000$
 $\Rightarrow R_3 = 10000 \Omega$

7. A point dipole $\vec{p} = -p_0 \hat{x}$ is kept at the origin. The potential and electric field due to this dipole on the y-axis at a distance d are, respectively : (Take V = 0 at infinity)

(1)
$$\frac{|\vec{p}|}{4\pi\epsilon_0 d^2}, \frac{\vec{p}}{4\pi\epsilon_0 d^3}$$

(2)
$$\frac{|\vec{p}|}{4\pi\epsilon_0 d^2}, \frac{-\vec{p}}{4\pi\epsilon_0 d^3}$$

(3)
$$0, \frac{-\vec{p}}{4\pi\epsilon_0 d^3}$$

(4)
$$0, \frac{\vec{p}}{4\pi\epsilon_0 d^3}$$

Answer (3)

Sol.
$$\vec{E} = K \frac{\vec{p}}{r^3} \sqrt{3\cos^2 \theta + 1}$$

 $\Rightarrow \theta = \pi/2 \ (0, d, 0)$
 $\therefore \vec{E} = \frac{-k\vec{p}}{d^3}$



is 45°, and 40 times per minute where the dip is 30°. If B_1 and B_2 are respectively the total magnetic field due to the earth at the two places, then the ratio B_1/B_2 is best given by :

(1)	2.2	(2) 0.7
(3)	3.6	(4) 1.8

Answer (2)

Sol. $T_1 = 2 \text{ sec.}, T_2 = 3/2$

For place (1), $B_{H_1} = B_1 \cos 45^\circ - \frac{B_1}{\sqrt{2}}$

For place (2),
$$B_{H_2} = B_2 \cos 30^\circ = \frac{B_2 \sqrt{3}}{2}$$

$$\therefore \quad \mathbf{T} = 2\pi \sqrt{\frac{\mathbf{I}}{\mathbf{MB}_{\mathbf{H}}}} \qquad \therefore \quad \frac{\mathbf{T}_{1}}{\mathbf{T}_{2}} = \sqrt{\frac{\mathbf{B}_{\mathbf{H}_{2}}}{\mathbf{B}_{\mathbf{H}_{1}}}}$$
$$\frac{4 \times 4}{9} = \frac{\mathbf{B}_{2}\sqrt{3} \times \sqrt{2}}{2 \mathbf{B}_{1}} \quad \Rightarrow \quad \frac{\mathbf{B}_{1}}{\mathbf{B}_{2}} = \frac{\sqrt{6} \times 9}{2 \times 16}$$
$$\Rightarrow \quad \frac{\mathbf{B}_{1}}{\mathbf{B}_{2}} = 0.68 \approx 0.7.$$

9. An electromagnetic wave is represented by the electric field

 $\vec{E} = E_0 \hat{n} \sin[\omega t + (6y - 8z)]$. Taking unit vectors in x, y and z directions to be \hat{i} , \hat{j} , \hat{k} , the direction of propogation \hat{s} , is:

(1)
$$\hat{\mathbf{s}} = \left(\frac{-3\hat{\mathbf{j}} + 4\hat{\mathbf{k}}}{5}\right)$$

(2) $\hat{\mathbf{s}} = \left(\frac{3\hat{\mathbf{i}} - 4\hat{\mathbf{j}}}{5}\right)$
(3) $\hat{\mathbf{s}} = \left(\frac{-4\hat{\mathbf{k}} + 3\hat{\mathbf{j}}}{5}\right)$
(4) $\hat{\mathbf{s}} = \left(\frac{4\hat{\mathbf{j}} - 3\hat{\mathbf{k}}}{5}\right)$

Answer (1)

Sol. $E = E_0 \sin(\omega t + 6y - 8z)$

$$\hat{\mathbf{s}} = \frac{\mathbf{8}\hat{\mathbf{k}} - \mathbf{6}\hat{\mathbf{j}}}{\mathbf{10}} = \left(\frac{\mathbf{4}\hat{\mathbf{k}} - \mathbf{3}\hat{\mathbf{j}}}{\mathbf{5}}\right)$$

change in the internal energy of the gas along the path ca is -180 J. The gas absorbs 250 J of heat along the path ab and 60 J along the path bc. The work done by the gas along the path abc is :



Answer (1)

- Sol. For the process (c a), $\Delta U_{ca} = -180 \text{ J}$ For process $(b - c) \rightarrow \text{Isochoric} (W_{bc} = 0)$ $\therefore \Delta U = 60 \text{ J}$ Heat absorbs along (a - b), $Q_{ab} = 250 \text{ J}$ Also $\therefore \Delta U_{cycle} = 0$ $\therefore \Delta U_{ab} = 120 \text{ J}$ So $W_{a \rightarrow b} = 130 \text{ J}$ Total work done from $(a \rightarrow b \rightarrow c)$ $= W_{ab} + W_{bc} = 130 \text{ J}$
- 11. At 40°C, a brass wire of 1 mm radius is hung from the ceiling. A small mass, M is hung from the free end of the wire. When the wire is cooled down from 40°C to 20°C it regains its original length of 0.2 m. The value of M is close to :

(Coefficient of linear expansion and Young's modulus of brass are 10^{-5} /°C and 10^{11} N/m², respectively ; g = 10 ms⁻²)

(1)	0.5	kg		(2)	0.9	kg
(a)						

(3) 1.5 kg (4) 9 kg

Answer (Bonus)

Sol.
$$\frac{MgI}{A \Delta I} = Y$$

 $\therefore \Delta I_{Mechanical} = \frac{MgI}{AY}$
 $\Delta I_{Thermal} = I \alpha \Delta T = I \alpha \times 20$
 $\frac{MgI}{AY} = 20 \alpha I$
 $M = \frac{20 \times 10^{-5} \times \pi \times 1 \times 10^{-6} \times 10^{11}}{10} = 6.28 \text{ kg}$


constant angular speed of 40 π rad s⁻¹ about its axis, perpendicular to its plane. If the magnetic field at its centre is 3.8 × 10⁻⁹ T, then the charge carried by the ring is close to ($\mu_0 = 4\pi \times 10^{-7}$ N/A²).

(1) 4 × 10 ⁻⁵ C	(2) 3 × 10⁻⁵ C
(3) 7 × 10 ⁻⁶ C	(4) 2 × 10 ^{−6} C

Answer (2)

Sol.
$$B = \frac{\mu_0 i}{2a} \text{ and } \frac{\omega q}{2\pi} = i$$

 $\therefore B = \frac{\mu_0}{2a} \cdot \frac{\omega q}{2\pi}$
 $B = \frac{10^{-7} \times 40}{0.1} \times q \times \pi \implies q = 3 \times 10^{-5} \text{ C}$

 An excited He⁺ ion emits two photons in succession, with wavelengths 108.5 nm and 30.4 nm, in making a transition to ground state. The quantum number n, corresponding to its initial excited state is (for photon of

wavelength
$$\lambda$$
, energy $E = \frac{1240 \text{ eV}}{\lambda(\text{in nm})}$):
(1) n = 5 (2) n = 7
(3) n = 4 (4) n = 6

Answer (1)

Sol.
$$\Rightarrow \Delta E_n = -\frac{E_0 Z^2}{n^2}$$

Let it start from n to m and from m to ground.

Then
$$13.6 \times 4 \left| 1 - \frac{1}{m^2} \right| = \frac{hc}{30.4 \text{ nm}}$$

$$\Rightarrow 1 - \frac{1}{m^2} = 0.7498 \Rightarrow 0.25 = \frac{1}{m^2}$$

$$\therefore \text{ m = 2, and now } 13.6 \times 4 \left(\frac{1}{4} - \frac{1}{n^2} \right) = \frac{hc}{108.5 \times 10^{-9}}$$

$$n \approx 5.$$

14. The figure shows a square loop L of side 5 cm which is connected to a network of resistances. The whole setup is moving towards right with a constant speed of 1 cm s⁻¹. At some instant, a part of L is in a uniform magnetic field of 1 T, perpendicular to the plane of the loop. If the resistance of L is 1.7 Ω , the current in the loop at that instant will be close to:



$$\therefore R_{eq} = \frac{4}{3}\Omega + 1.7 = 3\Omega$$

$$i = \frac{(BLV)}{R_{eq}} = \frac{(1)(5 \times 10^{-2}) \times 10^{-2}}{3}$$

$$= \frac{5}{3} \times 10^{-4} A \simeq 1.7 \times 10^{-4} A$$

$$= 170 \ \mu A$$

15. A uniform rod of length I is being rotated in a horizontal plane with a constant angular speed about an axis passing through one of its ends. If the tension generated in the rod due to rotation is T(x) at a distance x from the axis, then which of the following graphs depicts it most closely?



Answer (3)

Sol.

$$T_{x} = \frac{M}{L} (L - x) \left\{ x + \frac{L - x}{2} \right\} \omega^{2} = \frac{M \omega^{2}}{2L} (L^{2} - x^{2})$$



separated by a distance d. The space between the plates of the two capacitors, is filled with three dielectrics, of equal thickness and dielectric constants K_1 , K_2 and K_3 . The first capacitor is filled as shown in fig. I, and the second one is filled as shown in fig II.

If these two modified capacitors are charged by the same potential V, the ratio of the energy stored in the two, would be (E_1 refers to capacitor (I) and E_2 to capacitor (II)):



(1)
$$\frac{\mathsf{E}_{1}}{\mathsf{E}_{2}} = \frac{(\mathsf{K}_{1} + \mathsf{K}_{2} + \mathsf{K}_{3})(\mathsf{K}_{2}\mathsf{K}_{3} + \mathsf{K}_{3}\mathsf{K}_{1} + \mathsf{K}_{1}\mathsf{K}_{2})}{\mathsf{K}_{1}\mathsf{K}_{2}\mathsf{K}_{3}}$$

(2)
$$\frac{\mathsf{E}_{1}}{\mathsf{E}_{2}} = \frac{(\mathsf{K}_{1} + \mathsf{K}_{2} + \mathsf{K}_{3})(\mathsf{K}_{2}\mathsf{K}_{3} + \mathsf{K}_{3}\mathsf{K}_{1} + \mathsf{K}_{1}\mathsf{K}_{2})}{9\,\mathsf{K}_{1}\mathsf{K}_{2}\mathsf{K}_{3}}$$

(3)
$$\frac{\mathsf{E}_{1}}{\mathsf{E}_{2}} = \frac{9\,\mathsf{K}_{1}\mathsf{K}_{2}\mathsf{K}_{3}}{(\mathsf{K}_{1}+\mathsf{K}_{2}+\mathsf{K}_{3})\,(\mathsf{K}_{2}\mathsf{K}_{3}+\mathsf{K}_{3}\mathsf{K}_{1}+\mathsf{K}_{1}\mathsf{K}_{2})}$$

(4)
$$\frac{\mathsf{E}_{1}}{\mathsf{E}_{2}} = \frac{\mathsf{K}_{1}\mathsf{K}_{2}\mathsf{K}_{3}}{(\mathsf{K}_{1}+\mathsf{K}_{2}+\mathsf{K}_{3})(\mathsf{K}_{2}\mathsf{K}_{3}+\mathsf{K}_{3}\mathsf{K}_{1}+\mathsf{K}_{1}\mathsf{K}_{2})}$$

Answer (3)

Sol.
$$\frac{1}{C_{1}} = \frac{d}{3A\epsilon_{0}} \left(\frac{1}{K_{1}} + \frac{1}{K_{2}} + \frac{1}{K_{3}} \right)$$
$$C_{1} = \frac{3A\epsilon_{0} (K_{1}K_{2}K_{3})}{d (K_{1}K_{2} + K_{2}K_{3} + K_{3}K_{1})}$$
$$C_{2} = \frac{A\epsilon_{0}}{3d} (K_{1} + K_{2} + K_{3})$$
$$\frac{E_{1}}{E_{2}} = \frac{C_{1}}{C_{2}} = \frac{3K_{1}K_{2}K_{3}}{(K_{1}K_{2} + K_{2}K_{3} + K_{3}K_{1})} \times \frac{3}{(K_{1} + K_{2} + K_{3})}$$
$$\Rightarrow \quad \frac{E_{1}}{E_{2}} = \frac{9K_{1}K_{2}K_{3}}{(K_{1} + K_{2} + K_{3}) (K_{1}K_{2} + K_{2}K_{3} + K_{3}K_{1})}$$

in the figure. The work function of sodium, from the data plotted in the figure, will be:

(Given: Planck's constant)

(h) = 6.63×10^{-34} Js, electron

charge e = 1.6×10^{-19} C)



Answer (4)

Sol.
$$\phi = \frac{hc}{\lambda} = hv$$

 $\therefore \phi = h \times 4 \times 10^{14} \text{ Hz} = 1.654 \text{ eV}$
 $\Rightarrow \phi \approx 1.66 \text{ eV}$

18. The trajectory of a projectile near the surface of the earth is given as $y = 2x - 9x^2$. If it were launched at an angle θ_0 with speed v_0 then (g = 10 ms⁻²):

(1)
$$\theta_0 = \sin^{-1} \left(\frac{1}{\sqrt{5}} \right)$$
 and $v_0 = \frac{5}{3} \text{ms}^{-1}$
(2) $\theta_0 = \cos^{-1} \left(\frac{2}{\sqrt{5}} \right)$ and $v_0 = \frac{3}{5} \text{ms}^{-1}$

(3)
$$\theta_0 = \cos^{-1} \left(\frac{1}{\sqrt{5}} \right)$$
 and $v_0 = \frac{5}{3} \text{ms}^{-1}$
(4) $\theta_0 = \sin^{-1} \left(\frac{2}{\sqrt{5}} \right)$ and $v_0 = \frac{3}{5} \text{ms}^{-1}$

Answer (3)

Sol.
$$y = 2x - 9x^2$$

 $\Rightarrow v_0 = \frac{5}{3} m/s$

Comparing it with equation of trajectory

y = x tan
$$\theta$$
 - $\frac{g x^2}{24^2 \cos^2 \theta}$
∴ tan θ = 2
And $9 = \frac{10 \times 5}{2 v_0^2}$ 1



submarine (B) travelling at 27 km/hr. B sends a sonar signal of 500 Hz to detect A and receives a reflected sound of frequency v. The value of v is close to : (Speed of sound in water = 1500 ms⁻¹)

(1) 499 Hz	(2) 504 Hz
(3) 507 Hz	(4) 502 Hz

Answer (4)

Sol. f_1 (frequency received by A)

$$=v_0 \left[\frac{1500-5}{1500-7.5} \right]$$

f₂ [frequency received by B]

$$= v_0 \times \frac{1495}{1492.5} \times \frac{1507.5}{1505}$$

= 502 Hz.

20. A concave mirror has radius of curvature of 40 cm. It is at the bottom of a glass that has water filled up to 5 cm (see figure). If a small particle is floating on the surface of water, its image as seen, from directly above the glass, is at a distance d from the surface of water. The value of d is close to :

(Refractive index of water = 1.33)





Sol.
$$\frac{1}{V} + \frac{1}{U} = \frac{-1}{20}$$
$$\frac{1}{V} - \frac{1}{5} = \frac{-1}{20}$$
$$\frac{1}{V} = \frac{-1}{20} + \frac{1}{5}$$
$$\frac{1}{V} = \frac{3}{20}$$
$$d = \left(\frac{20}{3} + 5\right) \times 3/4$$
$$= \frac{35}{4}$$
$$d = 8.8 \text{ cm}$$

figure. The measured voltage is plotted as a function of the current, and the following graph is obtained :



If V_0 is almost zero, identify the correct statement :

- (1) The emf of the battery is 1.5 V and its internal resistance is 1.5 Ω
- (2) The emf of the battery is 1.5 V and the value of R is 1.5 Ω
- (3) The value of the resitance R is 1.5 Ω
- (4) The potential difference across the battery is 1.5 V when it sends a current of 1000 mA

Answer (1)

Sol.
$$V = \frac{ER}{R+r}$$

for $R = \infty$, $V = E = 1.5 V$
for $R = 0$, $I = E/r = 1$
 $r = 1.5 \Omega$

22. The truth table for the circuit given in the fig. is:





Sol. \Rightarrow y = $\overline{\mathbf{A} \cdot (\mathbf{A} + \mathbf{B})} = \overline{\mathbf{A}} + \overline{(\mathbf{A} + \mathbf{B})}$

$$\Rightarrow \mathbf{y} = \overline{\mathbf{A}} + \overline{\mathbf{A}} \cdot \overline{\mathbf{B}} = \overline{\mathbf{A}} (\mathbf{1} + \overline{\mathbf{B}})$$

 \Rightarrow y = \overline{A}

23. Shown in the figure is a shell made of a conductor. It has inner radius a and outer radius b, and carries charge Q. At its centre is a dipole \overline{P} as shown. In this case:



- (1) Surface charge density on the outer surface depends on $|\vec{\mathbf{p}}|$
- (2) Surface charge density on the inner

surface is uniform and equal to $\frac{(Q/2)}{4\pi a^2}$

- (3) Electric field outside the shell is the same as that of point charge at the centre of the shell
- (4) Surface charge density on the inner surface of the shell is zero everywhere

Answer (3)

- Sol. Since dipole is having zero net charge. So inside surface shall have non-zero non-uniform charge distribution. And net field outside the region would be same as that would have been for point charge at surface.
- 24. Which of the following combinations has the dimension of electrical resistance (ϵ_0 is the permittivity of vacuum and μ_0 is the permeability of vacuum)?

(1)
$$\sqrt{\frac{\varepsilon_0}{\mu_0}}$$
 (2) $\frac{\varepsilon_0}{\mu_0}$
(3) $\sqrt{\frac{\mu_0}{\varepsilon_0}}$ (4) $\frac{\mu_0}{\varepsilon_0}$

 $\mathbf{OOI}. \ \mathbf{AS} \ \mathbf{WC} \ \mathbf{WIOW} \ \mathbf{U} = \mathbf{WO}$

$$t = \frac{L}{R}$$
$$R^{2} \frac{C}{L} = 1$$
$$R = \sqrt{\frac{L}{C}} = \sqrt{\frac{\mu_{0}}{\epsilon_{0}}}$$

25. A progressive wave travelling along the positive x-direction is represented by $y(x,t) = A \sin(kx \cdot \omega t + \phi)$. Its snapshot at t = 0 is given in the figure.



For this wave, the phase ϕ is :

(1)
$$\pi$$
 (2) $-\frac{\pi}{2}$

(3)
$$\frac{\pi}{2}$$
 (4) 0

Answer (1)

Sol. y = A sin ($\omega t - kx + \phi$)

At t = 0 and x = 0 particle is at mean position and will proceed in positive y direction

- 26. A man (mass = 50 kg) and his son (mass = 20 kg) are standing on a frictionless surface facing each other. The man pushes his son so that he starts moving at a speed of 0.70 ms⁻¹ with respect to the man. The speed of the man with respect to the surface is :
 - (1) 0.20 ms^{-1} (2) 0.14 ms^{-1}

(3) 0.47 ms^{-1} (4) 0.28 ms^{-1}

Answer (1)

Sol. 50
$$V_1 = 20 V_2$$

 $V_1 + V_2 = 0.70$
 $V_1 = 0.20$



water at 50°C, finally no ice is left and the water is at 0°C. The value of latent heat of ice, in cal g^{-1} is :

(1)
$$\frac{50M_2}{M_1} - 5$$

(2) $\frac{5M_1}{M_2} - 50$
(3) $\frac{50M_2}{M_1}$
(4) $\frac{5M_2}{M_1} - 5$

Answer (1)

Sol. $M_1 \times 5 + M_1 L = M_2 50$

$$L = \frac{50M_2}{M_1} - 5$$

28. Two moles of helium gas is mixed with three moles of hydrogen molecules (taken to be rigid). What is the molar specific heat of mixture at constant volume?

(R = 8.3 J/mol K)

- (1) 19.7 J/mol K
- (2) 21.6 J/mol K
- (3) 15.7 J/mol K
- (4) 17.4 J/mol K

Answer (4)

Sol.
$$5C_V = 2 \times \frac{3R}{2} + 3 \times \frac{5R}{2}$$

 $C_V = \frac{21R}{10}$

29. A person of mass M is, sitting on a swing of length L and swinging with an angular amplitude θ_0 . If the person stands up when the swing passes through its lowest point, the work done by him, assuming that his centre of mass moves by a distance I (I<<L), is close to :

(1) Mgl

(2) Mgl
$$(1 + \theta_0^2)$$

$$(3) \operatorname{Mgl}\left(1+\frac{\theta_0^2}{2}\right)$$

(4) Mgl
$$(1 - \theta_0^2)$$

Sol. $W_{man} = Mg_{eff} I$

 $\boldsymbol{g_{eff}} = \boldsymbol{g(1 + \theta_0^2)}$

 $W_{man} = Mgl(1+\theta_0^2)$

30. The transfer characteristic curve of a transistor, having input and output resistance 100 Ω and 100 k Ω respectively, is shown in the figure. The Voltage and Power gain, are respectively :



(1)
$$5 \times 10^4$$
, 2.5 × 10⁶

- (2) 2.5×10^4 , 2.5×10^6
- (3) 5×10^4 , 5×10^6
- (4) 5 × 10⁴, 5 × 10⁵

Answer (1)

Sol.
$$\beta = \frac{\mathbf{I_c}}{\mathbf{I_b}}$$

$$=\frac{5\times10^{-3}}{100\times10^{-6}}$$

= 50

Voltage gain
$$=\beta \frac{R_0}{R_i} = 5 \times 10^4$$

Power gain = β (voltage gain)

 $= 250 \times 10^4 = 2.5 \times 10^6$





1. The major product of the following addition reaction is

H₃C-CH = CH₂
$$\xrightarrow{Cl_2/H_2O}$$

(1) H₃C CH₃
(2) CH₃-CH-CH₂
| |
CI OH
(3) H₃C \xrightarrow{O}
(4) H₃C-CH-CH₂
| |
OH CI

Answer (4)

Sol.
$$CH_3 - CH = CH_2 \xrightarrow[H_2O]{} CH_3 - CH - CH_2$$

 $\downarrow H_{2}O, -H^*$
 $CH_3 - CH - CH_2$
 $\downarrow H_{2}O, -H^*$
 $CH_3 - CH - CH_2$
 $\downarrow OH_2O$

- 2. But-2-ene on reaction with alkaline KMnO₄ at elevated temperature followed by acidification will give :
 - (1) 2 molecules of CH₃CHO
 - (2) 2 molecules of CH₃COOH
 - (3) CH₃-CH-CH-CH₃ | | OH OH
 - (4) One molecule of CH_3CHO and one molecule of CH_3COOH

Answer (2)

Sol. $CH_3 - CH \ge CH - CH_3 \xrightarrow{KMnO_4} 2CH_3COOH$

- 3. The correct sequence of thermal stability of the following carbonates is :
 - (1) $MgCO_3 < CaCO_3 < SrCO_3 < BaCO_3$
 - (2) $BaCO_3 < SrCO_3 < CaCO_3 < MgCO_3$
 - (3) $MgCO_3 < SrCO_3 < CaCO_3 < BaCO_3$
 - (4) $BaCO_3 < CaCO_3 < SrCO_3 < MgCO_3$

Answer (1)

Sol. Stability of alkaline earth metal carbonates increases down the group :

 $MgCO_3 < CaCO_3 < SrCO_3 < BaCO_3$

4. The major product(s) obtained in the following reaction is/are



- 5. Which of the following statements is not true about RNA?
 - (1) It usually does not replicate
 - (2) It is present in the nucleus of the cell
 - (3) It controls the synthesis of protein
 - (4) It has always double stranded $\alpha\text{-helix}$ structure

Answer (4)

Sol. RNA has a single helix structure.

DNA has a double helix structure.

6. The complex ion that will lose its crystal field stabilization energy upon oxidation of its metal to +3 state is :





$$\begin{array}{c} \mathsf{Co}^{2+}(\mathsf{d}^{7}) \longrightarrow \mathsf{Co}^{3+}(\mathsf{d}^{6}) \\ t_{2g}^{6} e_{g}^{1} & \xrightarrow{t_{2g}} t_{2g}^{6} e_{g}^{0} \end{array}$$
$$\begin{array}{c} \mathsf{Zn}^{2+}(\mathsf{d}^{10}) \longrightarrow \mathsf{Zn}^{2+}(\mathsf{d}^{9}) \\ t_{2g}^{6} e_{g}^{4} & \xrightarrow{t_{2g}} t_{2g}^{6} e_{g}^{3} \end{array}$$
$$\begin{array}{c} \mathsf{Fe}^{2+}(\mathsf{d}^{6}) \longrightarrow \mathsf{Fe}^{3+}(\mathsf{d}^{5}) \\ t_{2g}^{6} e_{g}^{0} & \xrightarrow{t_{2g}} t_{2g}^{6} e_{g}^{0} \end{array}$$

So, only Fe²⁺ will lose crystal field stabilisation upon oxidation to +3, others will gain crystal field stabilisation

7. An element has a face-centred cubic (fcc) structure with a cell edge of a. The distance between the centres of two nearest tetrahedral voids in the lattice is :

(1)
$$\frac{3}{2}a$$
 (2) $\frac{a}{2}$
(3) a (4) $\sqrt{2}a$

Answer (2)



body diagonal at a distance of $\frac{\sqrt{3}a}{4}$ from the corner. Together they form a smaller cube of edge length $\frac{a}{2}$.

8. Glucose and Galactose are having identical configuration in all the positions except position.

(1) C – 2	(2) C – 5
(3) C – 3	(4) C – 4

Answer (4)

Sol. Galactose and Glucose are C_4 epimers.

9. The metal that gives hydrogen gas upon treatment with both acid as well as base is :

(1) Zinc	(2) Magnesium
(3) Iron	(4) Mercury

Answer (1)

Sol. Zn + NaOH
$$\longrightarrow$$
 Na₂ZnO₂ + H₂
Zn + H₂SO₄ \longrightarrow ZnSO₄ + H₂



Answer (1)

Sol. EWG attached to benzene ring will reduce the basic strength and increase pK_b while EDG decreases pK_b .

Correct order of pK_b

11. The electrons are more likely to be found :



- (1) In the region a and c
- (2) Only in the region c
- (3) Only in the region a
- (4) In the region a and b

Answer (1)



both 'a' and 'c'.

12. Complete removal of both the axial ligands (along the z-axis) from an octahedral complex leads to which of the following splitting patterns? (relative orbital energies not on scale).





- Sol. The field becomes square planar and the order of energy is $d_{x^2-y^2} > d_{xy} > d_{z^2} > d_{zx} = d_{yz}$.
- 13. An example of a disproportionation reaction is:
 - (1) $2MnO_4^- + 10I^- + 16H^+ \rightarrow 2Mn^{2+} + 5I_2 + 8H_2O$
 - (2) $2CuBr \rightarrow CuBr_2$ + Cu
 - (3) $2KMnO_4 \rightarrow K_2MnO_4 + MnO_2 + O_2$
 - (4) $2NaBr + Cl_2 \rightarrow 2NaCl + Br_2$

Answer (2)

 $\begin{array}{ccc} \text{Sol. } CuBr & \longrightarrow & Cu + CuBr_2 \\ & & & Cu^{\circ} & & Cu^{2+} \end{array}$

It is an example of disproportionation reaction.



15. An organic compound 'A' is oxidized with Na_2O_2 followed by boiling with HNO_3 . The resultant solution is then treated with ammonium molybdate to yield a yellow precipitate

Based on above observation, the element present in the given compound is :

- (1) Fluorine
- (2) Nitrogen
- (3) Phosphorus
- (4) Sulphur

Answer (3)

Sol. Phosphorus is detected in the form of canary yellow ppt on reaction with ammonium molybdate.



(1)
$$(SiO_4)^{4-}$$

(3) SiO₂

(4) (SiO₃)²⁻

Answer (1)

- Sol. These are examples of silicates, the basic unit being SiO_4^{4-} in each of them.
- 17. The mole fraction of a solvent in aqueous solution of a solute is 0.8. The molality (in mol kg^{-1}) of the aqueous solution is :
 - (1) 13.88×10^{-2} (2) 13.88×10^{-3} (3) 13.88 (4) 13.88×10^{-1}

Answer (3)

Sol. Let, total 1 moles be present

n_{solute} = 0.2

$$n_{solvent} = 0.8 \Rightarrow g_{solvent} = 0.8 \times 18$$

$$m = \frac{0.2 \times 1000}{0.8 \times 18}$$

$$=\frac{1000}{4\times18}\approx13.88$$

18. The group number, number of valence electrons, and valency of an element with atomic number 15, respectively, are :

(1) 13, 5 and 5 (2) 13, 6 and	(1)	15, 5 and 3	(2) 15, 6 and
---------------------------------	-----	-------------	---------------

(3) 16, 5 and 2 (4) 16, 6 and 3

Answer (1)

Sol. Phosphorus has atomic number equal to 15. Its group number is 15, it has 5 valence electrons and valency equal to 3.

Al(OH)₃ = 2.4×10^{-24} : (1) 3×10^{-19} (2) 12×10^{-21} (3) 12×10^{-23} (4) 3×10^{-22}

Answer (4)

Sol. Al(OH)₃
$$\rightleftharpoons$$
 Al³⁺ + $\underset{s}{\overset{3OH^-}{,}}$, K_{sp} = 2.4×10⁻²⁴

s(0.2)³ = 2.4 × 10⁻²⁴

$$s = \frac{24 \times 10^{-25}}{8 \times 10^{-3}} = 3 \times 10^{-22} \, \frac{\text{mol}}{\text{L}}$$

- 20. An ideal gas is allowed to expand from 1 L to 10 L against a constant external pressure of 1 bar. The work done in kJ is :
 - (1) -9.0 (2) -0.9

Answer (2)

- 21. The idea of froth floatation method came from a person X and this method is related to the process Y of ores. X and Y, respectively, are :
 - (1) Fisher woman and concentration
 - (2) Washer woman and concentration
 - (3) Washer man and reduction
 - (4) Fisher man and reduction

Answer (2)

Sol. Froth floatation is a method of concentration and it was discovered by a washer women.



- (1) PVC (2) Buna-N
- (3) Bakelite (4) Nylon 6

Answer (3)

- Sol. Bakelite is an example of thermosetting polymer.
- 23. Peptization is a :
 - (1) Process of converting a colloidal solution into precipitate
 - (2) Process of converting precipitate into colloidal solution
 - (3) Process of converting soluble particles to form colloidal solution
 - (4) Process of bringing colloidal molecule into solution

Answer (2)

- Sol. Peptisation is the process of converting a precipitate into a colloidal sol by shaking it with dispersion medium in the presence of small amount of electrolyte.
- 24. The correct set of species responsible for the photochemical smog is :
 - (1) CO_2 , NO_2 , SO_2 and hydrocarbons
 - (2) N_2 , O_2 , O_3 and hydrocarbons
 - (3) NO, NO₂, O_3 and hydrocarbons
 - (4) N_2 , NO_2 and hydrocarbons

Answer (3)

- Sol. Photochemical smog contains oxides of nitrogen, ozone and hydrocarbons.
- 25. Enthalpy of sublimation of iodine is 24 cal g⁻¹ at 200°C. If specific heat of $I_2(s)$ and $I_2(vap)$ are 0.055 and 0.031 cal g⁻¹K⁻¹ respectively, then enthalpy of sublimation of iodine at 250°C in cal g⁻¹ is :

(1) 11.4	(2) 2.85
----------	----------

(3) 5.7 (4) 22.8

Answer (4)

$$\therefore \quad (\Delta H)_{250} = (\Delta H)_{200} + (0.031 - 0.055) \ 50$$
$$= 24 - 50 \times 0.024$$
$$= 22.8$$

26. The major products of the following reaction are :



 $Co^{3+} + e^- \rightarrow Co^{2+}$; $E^\circ = +1.81 V$

 $Pb^{4+} + 2e^- \rightarrow Pb^{2+}$; E° = +1.67 V

 $Ce^{4+} + e^- \rightarrow Ce^{3+}$; E° = +1.61 V

 $\rm Bi^{3+}$ + 3e^- \rightarrow Bi ; E° = +0.20 V

Oxidizing power of the species will increase in the order :

- (1) $Co^{3+} < Ce^{4+} < Bi^{3+} < Pb^{4+}$
- (2) $Co^{3+} < Pb^{4+} < Ce^{4+} < Bi^{3+}$
- (3) $Ce^{4+} < Pb^{4+} < Bi^{3+} < Co^{3+}$
- (4) $Bi^{3+} < Ce^{4+} < Pb^{4+} < Co^{3+}$

Answer (4)

Sol. Greater the reduction potential, greater is the oxidising power.

So, Co³⁺ > Pb⁴⁺ > Ce⁴⁺ > Bi³⁺

- 28. The correct statement among the following is :
 - (1) $(SiH_3)_3N$ is planar and less basic than $(CH_3)_3N$
 - (2) $(SiH_3)_3N$ is pyramidal and more basic than $(CH_3)_3N$
 - (3) (SiH₃)₃N is pyramidal and less basic than (CH₃)₃N
 - (4) $(SiH_3)_3N$ is planar and more basic than $(CH_3)_3N$

Answer (1)



Trisilylamine is planar, due to backbonding of lone pairs of nitrogen into vacant d-orbitals of Si. In trimethylamine, there is no such delocalisation and hence it is more basic.

29. In the following reaction; $xA \rightarrow yB$

$$\log_{10}\left[-\frac{d[A]}{dt}\right] = \log_{10}\left[\frac{d[B]}{dt}\right] + 0.3010$$

'A' and 'B' respectively can be :

- (2) N_2O_4 and NO_2
- (3) n-Butane and Iso-butane
- (4) C_2H_2 and C_6H_6

Answer (1)

Sol. $xA \rightarrow yB$

$$\therefore \quad \frac{-dA}{xdt} = \frac{1}{y}\frac{dB}{dt}$$

$$\frac{-dA}{dt} = \frac{dB}{dt} \times \frac{x}{y}$$

$$\log\left[\frac{-dA}{dt}\right] = \log\left[\frac{dB}{dt}\right] + \log\left(\frac{x}{y}\right)$$

$$\frac{x}{y} = 2$$

The reaction is of type 2A \rightarrow B.

30. 5 moles of AB_2 weigh 125×10^{-3} kg and 10 moles of A_2B_2 weigh 300×10^{-3} kg. The molar mass of $A(M_A)$ and molar mass of $B(M_B)$ in kg mol⁻¹ are :

(1)
$$M_A = 25 \times 10^{-3}$$
 and $M_B = 50 \times 10^{-3}$

- (2) $M_{A} = 50 \times 10^{-3}$ and $M_{B} = 25 \times 10^{-3}$
- (3) $M_A = 5 \times 10^{-3}$ and $M_B = 10 \times 10^{-3}$
- (4) $M_A = 10 \times 10^{-3}$ and $M_B = 5 \times 10^{-3}$

Answer (3)

Sol. 5 mol AB₂ weighs 125 g

 \therefore AB₂ = 25 g/mol

10 mol
$$A_2B_2$$
 weighs 300 g

- \therefore A₂B₂ = 30 g/mol
- ... Molar mass of A = 5

Molar mass of B = 10



If A is a symmetric matrix and B is a skew-1. $=\frac{\alpha}{1-\alpha}+\frac{\beta}{1-\beta}$ symmetric matrix such that $A+B = \begin{vmatrix} 2 & 3 \\ 5 & -1 \end{vmatrix}$, $=\frac{\alpha+\beta-2\alpha\beta}{1-(\alpha+\beta)+\alpha\beta}$ then AB is equal to : $=\frac{\frac{25}{375}+\frac{4}{375}}{1-\frac{25}{375}-\frac{2}{375}}=\frac{29}{375-25-2}$ $\begin{array}{c|c} 1 & 4 & -2 \\ 1 & -4 \end{array}$ (2) $\begin{array}{c} 4 & -2 \\ -1 & -4 \end{array}$ (3) $\begin{bmatrix} -4 & -2 \\ -1 & 4 \end{bmatrix}$ (4) $\begin{bmatrix} -4 & 2 \\ 1 & 4 \end{bmatrix}$ $=\frac{29}{348}=\frac{1}{12}$ Answer (2) If $\mathbf{B} = \begin{bmatrix} 5 & 2\alpha & 1 \\ 0 & 2 & 1 \\ \alpha & 3 & -1 \end{bmatrix}$ is the inverse of a 3 × 3 Sol. Let $A = \begin{bmatrix} a & c \\ c & b \end{bmatrix}$ and $B = \begin{bmatrix} 0 & d \\ -d & 0 \end{bmatrix}$ 3. $\Rightarrow \mathbf{A} + \mathbf{B} = \begin{bmatrix} \mathbf{a} & \mathbf{c} + \mathbf{d} \\ \mathbf{c} - \mathbf{d} & \mathbf{b} \end{bmatrix} = \begin{bmatrix} \mathbf{2} & \mathbf{3} \\ \mathbf{5} & -\mathbf{1} \end{bmatrix}$ matrix A, then the sum of all value of α for \Rightarrow a = 2, b = -1, c - d = 5, c + d = 3 which det (A) + 1 = 0, is : (1) -1 (2) 2 \Rightarrow a = 2, b = -1, c = 4, d = -1 (4) 1 (3) 0 $\Rightarrow AB = \begin{bmatrix} 2 & 4 \\ 4 & -1 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ Answer (4) Sol. As $B = A^{-1}$ $= \begin{bmatrix} 4 & -2 \\ -1 & -4 \end{bmatrix}$ $|\mathbf{B}| = \frac{1}{|\mathbf{A}|}$ 2. If α and β are the roots of the equation Now $|\mathbf{B}| = \begin{vmatrix} 5 & 2\alpha & 1 \\ 0 & 2 & 1 \\ \alpha & 3 & -1 \end{vmatrix} = 2\alpha^2 - 2\alpha - 25$ $375x^2 - 25x - 2 = 0$, then $\lim_{n \to \infty} \sum_{r=1}^{n} \alpha^r + \lim_{n \to \infty} \sum_{r=1}^{n} \beta^r$ is equal to : Given |A| + 1 = 0 (1) $\frac{21}{346}$ $\frac{1}{2\alpha^2-2\alpha-25}$ +1=0 (2) $\frac{7}{116}$ $\Rightarrow \frac{2\alpha^2 - 2\alpha - 24}{2\alpha^2 - 2\alpha - 25} = 0$ 29 (3) 358 $\alpha = 4. -3$ Sum of values = 1 (4) $\frac{1}{12}$ 4. For $x \in R$, let [x] denote the greatest integer \leq x, then the sum of the series Answer (4) $\begin{bmatrix} -\frac{1}{3} \end{bmatrix} + \begin{bmatrix} -\frac{1}{3} - \frac{1}{100} \end{bmatrix} + \begin{bmatrix} -\frac{1}{3} - \frac{2}{100} \end{bmatrix} + \dots + \begin{bmatrix} -\frac{1}{3} - \frac{99}{100} \end{bmatrix}$ **Sol.** $375x^2 - 25x - 2 = 0$ $\alpha + \beta = \frac{25}{375}, \alpha\beta = \frac{-2}{375}$ is: (1) = 135(2) - 153

Answer (3)



 $\lim_{n \to \infty} \sum_{r=1}^{n} \left(\alpha^{r} + \beta^{r} \right) = \left(\alpha + \alpha^{2} + \alpha^{3} + \dots \infty \right) + \left(\beta + \beta^{2} + \beta^{3} + \dots \infty \right)$

As
$$[x] + [-x] = -1$$
 ($x \notin z$)
Required value
 $= -100 - \left\{ \left[\frac{1}{3} \right] + \left[\frac{1}{3} + \frac{1}{100} \right] + \dots \left[\frac{1}{3} + \frac{99}{100} \right] \right\}$
 $= -100 - \left[\frac{100}{3} \right]$
 $= -133$

If m is the minimum value of k for which the 5. function $f(x) = x\sqrt{kx - x^2}$ is increasing in the interval [0, 3] and M is the maximum value of f in [0, 3] when k = m, then the ordered pair (m, M) is equal to :

(1) $(4, 3\sqrt{2})$ (2)	(3, 3√3)
--------------------------	----------

(4) (4, 3√3) (3) $(5, 3\sqrt{6})$

Answer (4)

Sol.
$$f(x) = x\sqrt{kx - x^2} = \sqrt{kx^3 - x^4}$$

 $f'(x) = \frac{(3kx^2 - 4x^3)}{2\sqrt{kx^3 - x^4}} \ge 0 \text{ for } x \in [0, 3]$
 $\Rightarrow 3k - 4x \ge 0$
 $3k \ge 4x$
 $3k \ge 4x \text{ for } x \in [0, 3]$
Hence $k \ge 4$
i.e., $m = 4$
For $k = 4$,
 $\Rightarrow f(x) = x\sqrt{4x - x^2}$
For max. value, $f'(x) = 0$
 $\Rightarrow x = 3$
i.e., $y = 3\sqrt{3}$
Hence $M = 3\sqrt{3}$
6. If the line $\frac{x-2}{3} = \frac{y+1}{2} = \frac{z-1}{-1}$ intersects the plane $2x + 3y - z + 13 = 0$ at a point P and the plane $3x + y + 4z = 16$ at a point Q, then PQ is equal to:
(1) $2\sqrt{14}$ (2) 14
(3) $2\sqrt{7}$ (4) $\sqrt{14}$
Answer (1)

 $Q(0\mu + 2, 2\mu - 1, -\mu + 1)$ As P lies on 2x + 3y - z + 13 = 0 $6\lambda + 4 + 6\lambda - 3 + \lambda - 1 + 13 = 0$ \Rightarrow 13 λ = -13 $\Rightarrow \lambda = -1$ ∴ P(-1, -3, 2) Q lies on 3x + y + 4z = 16 $9\mu + 6 + 2\mu - 1 - 4\mu + 4 = 16$ \Rightarrow 7 μ = 7 $\Rightarrow \mu = 1$ Q is (5, 1, 0)

 $PQ = \sqrt{36 + 16 + 4} = \sqrt{56} = 2\sqrt{14}$

If three of the six vertices of a regular hexagon 7. are chosen at random, then the probability that the triangle formed with these chosen vertices is equilateral is:

(1)
$$\frac{3}{20}$$
 (2) $\frac{1}{5}$
(3) $\frac{3}{10}$ (4) $\frac{1}{10}$

Answer (4)

Sol. Only two equilateral triangles are possible i.e. $\triangle AEC$ and $\triangle BDF$.



Hence, required probability

$$=\frac{2}{{}^{6}C_{3}}=\frac{1}{10}$$

If the angle of intersection at a point where the 8. two circles with radii 5 cm and 12 cm intersect is 90°, then the length (in cm) of their common chord is:

(1)
$$\frac{120}{13}$$
 (2) $\frac{13}{2}$
(3) $\frac{13}{5}$ (4) $\frac{60}{13}$

Answer (1)

is





Length of common chord (PQ) = $\frac{120}{13}$

9. If the volume of parallelopiped formed by the vectors $\hat{\mathbf{i}} + \lambda \hat{\mathbf{j}} + \hat{\mathbf{k}}, \hat{\mathbf{j}} + \lambda \hat{\mathbf{k}}$ and $\lambda \hat{\mathbf{i}} + \hat{\mathbf{k}}$ is minimum, then λ is equal to :

(1)
$$\frac{1}{\sqrt{3}}$$
 (2) $-\sqrt{3}$
(3) $\sqrt{3}$ (4) $-\frac{1}{\sqrt{3}}$

Answer (1*) Vector are coplanar for $\lambda = \lambda_1$ where $\lambda_1^3 - \lambda_1 + 1 = 0 \Rightarrow$ volume is minimum when $\lambda = \lambda_1$.

Sol.
$$V = \begin{vmatrix} 1 & \lambda & 1 \\ 0 & 1 & \lambda \\ \lambda & 0 & 1 \end{vmatrix}$$
$$= |1(1) + \lambda(\lambda^{2}) + 1(-\lambda)|$$
$$= |\lambda^{3} - \lambda + 1|$$
Let $f(x) = x^{3} - x + 1$
f'(x) = $3x^{2} - 1$
For maxima/minima, f'(x) = 0
 $x = \pm \frac{1}{\sqrt{3}}$
f''(x) = $6x$
 \therefore f'' $\left(\frac{1}{\sqrt{3}}\right) > 0$
 $x = \frac{1}{\sqrt{3}}$ is point of local minima



When $\lambda = \lambda_1$, volume of parallelopiped is zero (vectors are coplanar)

10. The integral $\int \frac{2x^3 - 1}{x^4 + x} dx$ is equal to :

(Here C is a constant of integration)

(1)
$$\log_{e} \left| \frac{x^{3} + 1}{x} \right| + C$$
 (2) $\frac{1}{2} \log_{e} \frac{\left(x^{3} + 1 \right)^{2}}{\left| x^{3} \right|} + C$
(3) $\frac{1}{2} \log_{e} \frac{\left| x^{3} + 1 \right|}{x^{2}} + C$ (4) $\log_{e} \frac{\left| x^{3} + 1 \right|}{x^{2}} + C$

Answer (1)

Sol.
$$I = \int \frac{(2x^3 - 1)dx}{x^4 + x} = \int \frac{(2x - x^{-2})dx}{x^2 + x^{-1}}$$

Put $x^2 + x^{-1} = t$
 $(2x - x^{-2})dx = dt$
 $I = \int \frac{dt}{t} = \ln|t| + c$
 $= \ln|x^2 + x^{-1}| + c$
 $= \ln|\frac{x^3 + 1}{x}| + c$

11. If the normal to the ellipse $3x^2 + 4y^2 = 12$ at a point P on it is parallel to the line, 2x + y = 4 and the tangent to the ellipse at P passes through Q(4, 4) then PQ is equal to :

(1)
$$\frac{\sqrt{61}}{2}$$
 (2) $\frac{5\sqrt{5}}{2}$
(3) $\frac{\sqrt{157}}{2}$ (4) $\frac{\sqrt{221}}{2}$
Answer (2)



$$3x^2 + 4y^2 = 12 \qquad \Rightarrow \quad \frac{x^2}{2^2} + \frac{y^2}{(\sqrt{3})^2} = 1$$

Let point $P(2\cos\theta, \sqrt{3}\sin\theta)$

 \Rightarrow Equation of tangent at P is

$$\frac{x}{2}\cos\theta + \frac{y}{\sqrt{3}}\sin\theta = 1$$

$$\Rightarrow m_{T} = -\frac{\sqrt{3}}{2}\cot\theta = \frac{1}{2}$$

$$\tan\theta = -\sqrt{3} \quad \Rightarrow \theta = \pi - \frac{\pi}{3} \quad \text{or } \theta = 2\pi - \frac{\pi}{3}$$
If $\theta = \frac{2\pi}{3}$, then $P\left(-1, \frac{3}{2}\right)$ and $PQ = \frac{5\sqrt{5}}{2}$
If $\theta = \frac{5\pi}{3}$, then tangent does not pass through $Q(4, 4)$

- 12. Let S_n denote the sum of the first n terms of an A.P. If $S_4 = 16$ and $S_6 = -48$, then S_{10} is equal to :
 - (1) -260(2) -380(3) -320(4) -410
- Answer (3)
- Sol. $S_4 = 16$, $S_6 = -48$ 2(2a + 3d) = 16 $\Rightarrow 2a + 3d = 8$ Also 3[2a + 5d] = -48 $\Rightarrow 2a + 5d = -16$ 2d = -24 $d = -12 \Rightarrow a = 22$ $S_{10} = 5(44 + 9(-12))$ = -320
- 13. A 2 m ladder leans against a vertical wall. If the top of the ladder begins to slide down the wall at the rate 25 cm/sec., then the rate (in cm/sec.) at which the bottom of the ladder slides away from the wall on the horizontal ground when the top of the ladder is 1 m above the ground is :

(3)
$$\frac{25}{\sqrt{3}}$$
 (4) 25

Answer (3)

Sol. Given
$$\frac{dy}{dt} = -25$$
 at $y = 1$
 $x^2 + y^2 = 4$
When $y = 1$, $x = \sqrt{3}$
 $2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$
 $\Rightarrow x \frac{dx}{dt} + y \frac{dy}{dt} = 0$
 $\Rightarrow \sqrt{3} \frac{dx}{dt} + (-25) = 0$
 $\Rightarrow \frac{dx}{dt} = \frac{25}{\sqrt{3}}$ cm/s

- 14. The number of ways of choosing 10 objects out of 31 objects of which 10 are identical and the remaining 21 are distinct, is :
 - (1) $2^{20} + 1$ (2) 2^{21} (3) $2^{20} - 1$ (4) 2^{20}

Answer (4)

Sol. Number of ways of selecting 10 objects

= (10I, 0D) or (9I, 1D) or (8I, 1D) or ... (0I, 10D)

where D signifies distinct object and I indicates identical object

$$= 1 + {}^{21}C_1 + {}^{21}C_2 + \dots + {}^{21}C_{10}$$
$$= \frac{2^{21}}{2} = 2^{20}$$

15. Let P be the point of intersection of the common tangents to the parabola $y^2 = 12x$ and the hyperbola $8x^2 - y^2 = 8$. If S and S' denote the foci of the hyperbola where S lies on the positive x-axis then P divides SS' in a ratio :

(1) 13 : 11	(2) 14 : 13
(3) 5:4	(4) 2:1

Answer (3)



$$\mathbf{m} = \mathbf{e}^{\mathbf{y}} \frac{\mathbf{d}}{\mathbf{dy}}$$
Equation of tangent $\frac{x^2}{1} - \frac{y^2}{8} = 1$ is $\mathbf{y} = \mathbf{m} \times \sqrt{\mathbf{m}^2 - 8}$
for common tangent,

$$\frac{3}{\mathbf{m}} = \pm \sqrt{\mathbf{m}^2 - 8} \qquad \Rightarrow \frac{9}{\mathbf{m}^2} = \mathbf{m}^2 - 8$$
Put $\mathbf{m}^2 = \mathbf{t}$
 $\mathbf{t}^2 - 8\mathbf{t} - 9 = 0 \qquad \Rightarrow \mathbf{t}^2 - 9\mathbf{t} + \mathbf{t} - 9 = 0$
 $\Rightarrow (\mathbf{t} + 1) (\mathbf{t} - 9) = 0$
 $\therefore \mathbf{t} = \mathbf{m}^2 \ge 0 \qquad \Rightarrow \mathbf{t} = \mathbf{m}^2 = 9$
 $\Rightarrow \mathbf{m} = \pm 3$
 $\Rightarrow \text{ Equation of tangent is $\mathbf{y} = 3\mathbf{x} + 1$
or $\mathbf{y} = -3\mathbf{x} - 1$
Intersection point $P\left(-\frac{1}{3}, 0\right)$
 $8 = 1(\mathbf{e}^2 - 1) \qquad \Rightarrow \mathbf{e} = 3$
foci $(\pm 3, 0) \qquad \Rightarrow \frac{\mathbf{S}'}{(-3, 0)} \qquad (-\frac{1}{3}, 0)\mathbf{p}^{-(3, 0)}$
 $8 = 1(\mathbf{e}^2 - 1) \qquad \Rightarrow \mathbf{e} = 3$
foci $(\pm 3, 0) \qquad \Rightarrow \frac{\mathbf{S}'}{(-3, 0)} \qquad (-\frac{1}{3}, 0)\mathbf{p}^{-(3, 0)}$
 $8 = 1(\mathbf{e}^2 - 1) \qquad \Rightarrow \mathbf{e} = 3$
foci $(\pm 3, 0) \qquad \Rightarrow \frac{\mathbf{S}'}{(-3, 0)} \qquad (-\frac{1}{3}, 0)\mathbf{p}^{-(3, 0)}$
 $8 = 1(\mathbf{e}^2 - 1) \qquad \Rightarrow \mathbf{e} = 3$
foci $(\pm 3, 0) \qquad \Rightarrow \frac{\mathbf{S}'}{(-3, 0)} \qquad (-\frac{1}{3}, 0)\mathbf{p}^{-(3, 0)}$
 $8 = 1(\mathbf{e}^2 - 1) \qquad \Rightarrow \mathbf{e} = 3$
foci $(\pm 3, 0) \qquad \Rightarrow \frac{\mathbf{S}'}{(-3, 0)} \qquad (-\frac{1}{3}, 0)\mathbf{p}^{-(3, 0)}$
 $\mathbf{x}_1 + \mathbf{x}_2$
 $\mathbf{x} = 0$ is equal to :
(1) $\left(\frac{1}{\mathbf{e}}, -\frac{1}{\mathbf{e}^2}\right)$
(2) $\left(-\frac{1}{\mathbf{e}}, -\frac{1}{\mathbf{e}^2}\right)$
(3) $\left(-\frac{1}{\mathbf{e}}, \frac{1}{\mathbf{e}^2}\right)$
(4) $\left(\frac{1}{\mathbf{e}}, \frac{1}{\mathbf{e}^2}\right)$
Answer (3)
Sol. $(\mathbf{1} - \mathbf{x})$
Sol. $(1 - \mathbf{x})$
Sol. $(1 - \mathbf{x})$
 $\mathbf{S} = \mathbf{e} \Rightarrow \mathbf{y} = 1$
Differentiate (i) w.r. to \mathbf{x}
 $\mathbf{e}^{\mathbf{y}} \frac{\mathbf{d}}{\mathbf{dx}} + \mathbf{x} \frac{\mathbf{dy}}{\mathbf{dx}} + \mathbf{y} = 0$
Put $\mathbf{y} = 1$ in (i)
 $\mathbf{e} \frac{\mathbf{dy}}{\mathbf{dx}} + 0 + 1 = 0 \Rightarrow \frac{\mathbf{dy}}{\mathbf{dx}} = -\frac{1}{\mathbf{e}}$
 $\mathbf{x} = \mathbf{x} = \mathbf{x}$
 $\mathbf{x} = \mathbf{x} = \mathbf{x}$
 $\mathbf{x} = \mathbf{x} = \mathbf{x}$
 $\mathbf{x} = \mathbf{x}$
 $\mathbf{x$$

$$e^{y} \frac{d}{dx^{2}} + \frac{dy}{dx} \cdot e^{y} \frac{dy}{dx} + x \frac{d}{dx^{2}} + \frac{dy}{dx} + \frac{dy}{dx} = 0 \quad ...(iii)$$
Put $y = 1, x = 0, \frac{dy}{dx} = -\frac{1}{e}$

$$e \frac{d^{2}y}{dx^{2}} + \frac{1}{e} - \frac{2}{e} = 0 \Rightarrow \frac{d^{2}y}{dx^{2}} = \frac{1}{e^{2}}$$

$$\Rightarrow \left(\frac{dy}{dx}, \frac{d^{2}y}{dx^{2}}\right) = \left(-\frac{1}{e}, \frac{1}{e^{2}}\right)$$
7. If the data $x_{1}, x_{2}, ..., x_{10}$ is such that the mean of first four of these is 11, the mean of the remaining six is 16 and the sum of squares of all of these is 2,000; then the standard deviation of this data is :
(1) $2\sqrt{2}$ (2) 4
(3) 2 (4) $\sqrt{2}$
mswer (3)
pl. $\frac{x_{1} + x_{2} + x_{3} + x_{4}}{4} = 11 \text{ and } x_{1} + x_{2} + x_{3} + x_{4} = 44$
 $\frac{x_{5} + x_{6} + ... + x_{10}}{6} = 16 \Rightarrow x_{5} + x_{6} + ... + x_{10} = 96$
 $x_{1}^{2} + x_{2}^{2} + ... + x_{10}^{2} = 2000$
 $\sigma^{2} = \frac{\sum x_{1}^{2}}{N} - (\overline{x})^{2}$
 $= \frac{2000}{10} - \left(\frac{140}{10}\right)^{2} = 4$

18. The coefficient of
$$x^{18}$$
 in the product $(1 + x)$
 $(1 - x)^{10}(1 + x + x^2)^9$ is :

(1)

= 2

Sol.
$$(1-x)^{10} (1+x+x^2)^9 (1+x)$$

$$= (1-x^3)^9 (1-x^2)$$

$$= (1-x^3)^9 - x^2 (1-x^3)^9$$

$$\Rightarrow \text{Coefficient of } x^{18} \text{ in } (1-x^3)^9 - \text{coeff. of } x^{16} \text{ in } (1-x^3)^9.$$

$$= {}^9C_6 = \frac{9!}{6!3!} = \frac{7 \times 8 \times 9}{6} = 84$$



16.

Sol.

statements p, q, r are respectively :

(1) F, T, T	(2) T, T, F
(3) T, F, F	(4) T, F, T

Answer (2)

- Sol. $\mathbf{P} \rightarrow (\mathbf{\sim q \lor r})$ is a fallacy
 - \Rightarrow P is True and ~q \lor r is False
 - \Rightarrow P is True and ~q is False and r is False
 - \Rightarrow Truth values of p, q, r are
 - T, T, F respectively.
- 20. Let a random variable X have a binomial distribution with mean 8 and variance 4.

If P(X \leq 2) = $\frac{k}{2^{16}}$, then k is equal to :
(1) 121	(2) 1
(3) 17	(4) 137

Answer (4)

Sol.
$$\mu = 8$$
, $\sigma^2 = 4$
 $\Rightarrow \mu = np = 8$, $\sigma^2 = npq = 4$, $p + q = 1$
 $\Rightarrow q = \frac{1}{2}$, $p = \frac{1}{2}$, $n = 16$
 $P(X \le 2) = \frac{k}{2^{16}}$
 ${}^{16}C_0 \left(\frac{1}{2}\right)^{16} + {}^{16}C_1 \left(\frac{1}{2}\right)^{16} + {}^{16}C_2 \left(\frac{1}{2}\right)^{16} = \frac{k}{2^{16}}$
 $\Rightarrow k = (1 + 16 + 120) = 137$

21. Let $\vec{a} = 3\hat{i} + 2\hat{j} + 2\hat{k}$ and $\vec{b} = \hat{i} + 2\hat{j} - 2\hat{k}$ be two vectors. If a vector perpendicular to both the vectors $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$ has the magnitude 12 then one such vector is :

(1)	$4\left(-2\hat{i}-2\hat{j}+\hat{k}\right)$	(2)	$4\left(2\hat{i}+2\hat{j}-\hat{k}\right)$
(3)	$4 \Big(2 \hat{i} + 2 \hat{j} + \hat{k} \Big)$	(4)	$4 \Big(2 \hat{i} - 2 \hat{j} - \hat{k} \Big)$

Answer (4)

Sol. Let vector be
$$\lambda \left[\left(\vec{a} + \vec{b} \right) \times \left(\vec{a} - \vec{b} \right) \right]$$

$$\vec{a} + \vec{b} = 4\hat{i} + 4\hat{j}$$

$$\vec{a} - \vec{b} = 2\hat{i} + 4\hat{k}$$

$$= \lambda \Big[16\hat{i} - 16\hat{j} - 8\hat{k} \Big]$$
$$= 8\lambda \Big[2\hat{i} - 2\hat{j} - \hat{k} \Big]$$
$$\Rightarrow 12 = 8 |\lambda| \sqrt{4 + 4 + 1}$$
$$|\lambda| = \frac{1}{2}$$

Hence required vector is $\pm 4(2\hat{i}-2\hat{j}-\hat{k})$

22. Consider the differential equation, $y^2 dx + \left(x - \frac{1}{y}\right) dy = 0$. If value of y is 1 when

x = 1, then the value of x for which y = 2, is :

(1)
$$\frac{5}{2} + \frac{1}{\sqrt{e}}$$
 (2) $\frac{3}{2} - \sqrt{e}$
(3) $\frac{3}{2} - \frac{1}{\sqrt{e}}$ (4) $\frac{1}{2} + \frac{1}{\sqrt{e}}$

Answer (3)

Sol.
$$y^2 dx + \left(x - \frac{1}{y}\right) dy = 0$$

 $\frac{dx}{dy} + \left(\frac{1}{y^2}\right) x = \frac{1}{y^3}$
I.F. $= e^{\int \frac{1}{y^2} dy} = e^{-\frac{1}{y}}$
its solution is
 $x \cdot e^{-\frac{1}{y}} = \int e^{-\frac{1}{y}} \frac{1}{y^3} dy + c$
put $-\frac{1}{y} = t \Rightarrow \frac{1}{y^2} dy = dt$
 $\Rightarrow x \cdot e^{-\frac{1}{y}} = -\int te^t dt + c = -te^t + e^t + c$
 $x \cdot e^{-\frac{1}{y}} = e^{-\frac{1}{y}} \left(\frac{1}{y} + 1\right) + c$ passes through (1, 1)
 $\Rightarrow 1 = 2 + ce \Rightarrow c = -\frac{1}{e}$
 $\Rightarrow x = \left(1 + \frac{1}{y}\right) - \frac{1}{e}e^{\frac{1}{y}}$ passes through (k, 2)
 $\Rightarrow k = \frac{3}{2} - \frac{1}{\sqrt{e}}$



function such that f(2) = 6 and $f'(2) = \frac{1}{48}$. If $\int_{6}^{f(x)} 4t^{3}dt = (x-2)g(x), \text{ then } \lim_{x \to 2} g(x) \text{ is equal}$ to: (1) 18 (2) 36 (3) 24 (4) 12 Answer (1) Sol. $\int_{0}^{f(x)} 4t^{3}dt = (x-2)g(x)$

of.
$$\int_{6}^{9} 4t \, dt = (x - 2)g(x)$$

$$4(f(x))^{3} \cdot f'(x) = g'(x)(x - 2) + g(x)$$
put x = 2,
$$\frac{4(6)^{3} \cdot 1}{48} = g(2)$$

$$\lim_{x\to 2} g(x) = 18$$

24. The equation $y = \sin x \sin(x + 2) - \sin^2(x + 1)$ represents a straight line lying in :

- (1) Third and fourth quadrants only
- (2) First, third and fourth quadrants
- (3) First, second and fourth quadrants
- (4) Second and third quadrants only

Answer (1)

Sol. $y = sinx \cdot sin(x + 2) - sin^2(x + 1)$

$$= \frac{1}{2}\cos(-2) - \frac{\cos(2x+2)}{2} - \left[\frac{1 - \cos(2x+2)}{2}\right]$$
$$= \frac{(\cos 2) - 1}{2} = -\sin^2 1$$



Graph of y lies in III and IV Quadrant

a - b is equal to :

(1)
$$-\frac{2}{3}$$
 (2) 6
(3) $\frac{10}{3}$ (4) $\frac{8}{3}$

Answer (2)

Sol.
$$y^2 = 4x$$

 $x + y = 1$
 $y^2 = 4(1 - y)$
 $y^2 + 4y - 4 = 0$
 $(y + 2)^2 = 8$

 y
 $y^2 = 4x$
 $P(3 - 2\sqrt{2}, 2\sqrt{2} - 2)$
 $(1, 0)$
 $x + y = 1$

required area

 $y + 2 = \pm 2\sqrt{2}$

$$= \int_{0}^{3-2\sqrt{2}} 2\sqrt{x} \, dx + \frac{1}{2} \times (2\sqrt{2} - 2) \times (2\sqrt{2} - 2)$$

$$= \left[2 \times \frac{2}{3} x^{3/2} \right]_{0}^{3-2\sqrt{2}} + \frac{1}{2} (8 + 4 - 8\sqrt{2})$$

$$= \frac{4}{3} \times (3 - 2\sqrt{2}) \sqrt{3 - 2\sqrt{2}} + 6 - 4\sqrt{2}$$

$$= \frac{4}{3} (3 - 2\sqrt{2}) (\sqrt{2} - 1) + 6 - 4\sqrt{2}$$

$$= \frac{4}{3} (3\sqrt{2} - 3 - 4 + 2\sqrt{2}) + 6 - 4\sqrt{2}$$

$$= \left(6 - \frac{28}{3} \right) + \left(\frac{20}{3} - 4 \right) \sqrt{2}$$

$$= -\frac{10}{3} + \frac{8}{3} \sqrt{2}$$

$$\Rightarrow a - b = \frac{10}{3} + \frac{8}{3} = 6$$

26. The value of $\sin^{-1}\left(\frac{12}{13}\right) - \sin^{-1}\left(\frac{3}{5}\right)$ is equal to :

(1)
$$\frac{\pi}{2} - \sin^{-1}\left(\frac{56}{65}\right)$$
 (2) $\pi - \sin^{-1}\left(\frac{63}{65}\right)$
(3) $\pi - \cos^{-1}\left(\frac{33}{65}\right)$ (4) $\frac{\pi}{2} - \cos^{-1}\left(\frac{9}{65}\right)$
Answer (1)

Sol.
$$-\sin^{-1}\left(\frac{3}{5}\right) + \sin^{-1}\left(\frac{12}{13}\right) = -\sin^{-1}\left(\frac{3}{5} \times \frac{5}{13} - \frac{12}{13} \times \frac{4}{5}\right)$$

(:: xy ≥ 0 and x² + y² ≤ 1)



$$= \sin^{-1}\left(\frac{33}{65}\right)$$

$$= \cos^{-1}\left(\frac{56}{65}\right)$$

$$= \frac{\pi}{2} - \sin^{-1}\left(\frac{56}{65}\right)$$
27. If $\int_{0}^{\frac{\pi}{2}} \frac{\cot x}{\cot x + \csc x} dx = m(\pi + n)$, then $m \cdot n$ is equal to :
(1) $\frac{1}{2}$ (2) 1
(3) -1 (4) $-\frac{1}{2}$
Answer (3)
Sol. $\int_{0}^{\frac{\pi}{2}} \frac{\cot x}{2} dx}{1 + \cos x} = \int_{0}^{\frac{\pi}{2}} \left(1 - \frac{1}{1 + \cos x}\right) dx$

$$= \left[x\right]_{0}^{\frac{\pi}{2}} - \int_{0}^{\frac{\pi}{2}} \frac{1}{2\cos^{2}\frac{x}{2}} dx$$

$$= \frac{\pi}{2} - \left[\tan \frac{x}{2}\right]_{0}^{\frac{\pi}{2}} \sec^{2}\frac{x}{2} dx$$

$$= \frac{\pi}{2} - \left[\tan \frac{x}{2}\right]_{0}^{\frac{\pi}{2}} \sec^{2}\frac{x}{2} dx$$

$$= \frac{\pi}{2} - \left[1\right] = \left(\frac{\pi}{2} - 1\right)$$

$$m = \frac{1}{2}, n = -2$$

$$\Rightarrow mn = -1$$
28. For $x \in (0, \frac{3}{2})$, let $f(x) = \sqrt{x}, g(x) = \tan x$ and
 $h(x) = \frac{1 - x^{2}}{1 + x^{2}}$. If $\phi(x) = ((hof)og)(x)$, then $\phi\left(\frac{\pi}{3}\right)$ is equal to :
(1) $\tan \frac{5\pi}{12}$ (2) $\tan \frac{\pi}{12}$
(3) $\tan \frac{11\pi}{12}$ (4) $\tan \frac{7\pi}{12}$

Sol.
$$\phi\left(\frac{\pi}{3}\right) = h\left(f\left(g\left(\frac{\pi}{3}\right)\right)\right)$$

 $= h\left(f\left(\sqrt{3}\right)\right) = h(3^{\frac{1}{4}})$
 $= \frac{1-\sqrt{3}}{1+\sqrt{3}} = -\frac{1}{2}(1+3-2\sqrt{3}) = \sqrt{3}-2 = -(-\sqrt{3}+2)$
 $= -\tan 15^{\circ} = \tan(180^{\circ}-15^{\circ}) = \tan\left(\pi - \frac{\pi}{12}\right)$
 $= \tan\frac{11\pi}{12}$
29. The equation $|z-i| = |z-1|, i = \sqrt{-1}$, represents :
(1) The line through the origin with slope -1
(2) A circle of radius $\frac{1}{2}$
(3) A circle of radius 1
(4) The line through the origin with slope 1
Answer (4)
Sol. $|z-1| = |z-i|$
Let $z = x + iy$
 $(x-1)^2 + y^2 = x^2 + (y-1)^2$
 $1-2x = 1-2y$

Locus is straight line with slope 1

30. The number of solutions of the equation

1 + sin⁴x = cos²3x,
$$x \in \left[-\frac{5\pi}{2}, \frac{5\pi}{2}\right]$$
 is :

(1) 3	(2) 5
(3) 4	(4) 7

Answer (2)

 \Rightarrow x - y = 0

Sol. 1 + sin⁴x = cos²3x
L.H.S = 1 + sin⁴x, R.H.S = cos²3x
L.H.S
$$\ge$$
 1 R.H.S \le 1
 \Rightarrow L.H.S. = R.H.S. = 1
sin⁴x = 0 and cos² 3x = 1
sinx = 0 and (4cos²x - 3)²cos²x = 1
 \Rightarrow sinx = 0 and cos²x = 1
 \Rightarrow x = 0, ± π , ± 2π
 \Rightarrow Total number of solutions is 5

 \Rightarrow Total number of solutions is 5



Answers & Solutions

for

JEE (MAIN)-2019 (Online) Phase-2

(Physics, Chemistry and Mathematics)

Time : 3 hrs.

M.M.: 360

Important Instructions :

- 1. The test is of **3 hours** duration.
- 2. The Test Booklet consists of **90** questions. The maximum marks are **360**.
- 3. There are *three* parts in the question paper A, B, C consisting of **Physics**, **Chemistry** and **Mathematics** having 30 questions in each part of equal weightage.
- 4. Each question is allotted 4 (four) marks for each correct response. ¼ (one-fourth) marks will be deducted for indicating incorrect response of each question. No deduction from the total score will be made if no response is indicated for an item in the answer sheet.
- 5. There is only one correct response for each question.



- A uniform cylindrical rod of length L and radius r, is made from a material whose Young's modulus of Elasticity equals Y. When this rod is heated by temperature T and simultaneously subjected to a net longitudinal compressional force F, its length remains unchanged. The coefficient of volume expansion, of the material of the rod, is (nearly) equal to:
 - (1) 9F/(π r²YT)
 - (2) $3F/(\pi r^2 YT)$
 - (3) F/(3πr²YT)
 - (4) 6F/(πr²YT)

Sol.
$$Y = \frac{FL}{A \mid \Delta L \mid} \Rightarrow \Delta L = \frac{FL}{AY} = L \alpha T$$

 $\alpha = \frac{F}{AYT}$
 $\gamma = 3\alpha = \frac{3F}{\pi r^2 YT}$

2. A block of mass 5 kg is (i) pushed in case (A) and (ii) pulled in case (B), by a force F = 20 N, making an angle of 30° with the horizontal, as shown in the figures. The coefficient of friction between the block and floor is μ = 0.2. The difference between the accelerations of the block, in case (B) and case (A) will be :

 $(g = 10 \text{ ms}^{-2})$





For B N = 40 N

$$F = 8 \text{ N} \Rightarrow \frac{20\sqrt{3}}{2} - 8 = 5a_{\text{B}}$$
$$a_{\text{B}} = \frac{17.3 - 8}{5} = \frac{9.3}{5} = 1.86 \text{ m/s}^2$$

$$a_{B} - a_{A} = 0.8 \text{ m/s}^{2}$$

- 3. The electron in a hydrogen atom first jumps from the third excited state to the second excited state and subsequently to the first excited state. The ratio of the respective wavelengths, λ_1/λ_2 , of the photons emitted in this process is
 - (1) 7/5
 - (2) 27/5
 - (3) 9/7
 - (4) 20/7

Answer (4)

Sol.
$$\frac{1}{\lambda_1} = R\left[\frac{1}{9} - \frac{1}{16}\right] = R\frac{7}{144} \qquad \dots(i)$$
$$\frac{1}{\lambda_2} = R\left[\frac{1}{4} - \frac{1}{9}\right] = R\frac{5}{36} \qquad \dots(ii)$$
$$\implies \frac{\lambda_1}{\lambda_2} = \frac{\lambda_1}{\lambda_2} = \frac{5 \times 144}{36 \times 7} = \frac{20}{7}$$

4. A moving coil galvanometer, having a resistance G, produces full scale deflection when a current I_g flows through it. This galvanometer can be converted into (i) an ammeter of range 0 to $I_0(I_0 > I_g)$ by connecting a shunt resistance R_A to it and (ii) into a voltmeter of range 0 to $V(V = GI_0)$ by connecting a series resistance R_V to it. Then,



$$R_{V} = R_{V} = \left(I_{0} - I_{g}\right)$$

$$(2) \quad R_{A}R_{V} = G^{2}\left(\frac{I_{g}}{I_{0} - I_{g}}\right) \text{ and } \frac{R_{A}}{R_{V}} = \left(\frac{I_{0} - I_{g}}{I_{g}}\right)^{2}$$

$$(3) \quad R_{A}R_{V} = G^{2} \text{ and } \frac{R_{A}}{R_{V}} = \frac{I_{g}}{(I_{0} - I_{g})}$$

$$(4) \quad R_{A}R_{V} = G^{2}\left(\frac{(I_{0} - I_{g})}{I_{g}}\right) \text{ and } \frac{R_{A}}{R_{V}} = \left(\frac{I_{g}}{(I_{0} - I_{g})}\right)^{2}$$
Answer (1)
Sol. $V = I_{g}(R_{V} + G) = GI_{0} \qquad \dots(1)$

$$(I_{0} - I_{g})R_{A} = I_{g}G \qquad \dots(2)$$

Sol. V = I_g(R_v + G) = GI₀
(I₀ - I_g)R_A = I_gG
From (1), R_v =
$$\frac{G(I_0 - I_g)}{I_g}$$

From (2), R_A = $\frac{I_gG}{I_0 - I_g}$
 \Rightarrow R_AR_v = G²
 $\frac{R_A}{R_v} = \left(\frac{I_g}{I_0 - I_g}\right)^2$

5. A spring whose unstretched length is I has a force constant k. The spring is cut into two pieces of unstretched lengths ${\rm I_1}$ and ${\rm I_2}$ where, $I_1 = nI_2$ and n is an integer. The ratio k_1/k_2 of the corresponding force constants, k_1 and k_2 will be

(1)
$$n^2$$
 (2) $\frac{1}{n^2}$
(3) n (4) $\frac{1}{n}$

1

Answer (4)

Sol.
$$l_1 = nl_2$$

 $\therefore k \propto \frac{1}{l}$
 $\therefore \frac{k_1}{k_2} = \frac{l_2}{l_1} = \frac{1}{n}$

One kg of water, at 20°C, is heated in an 6. electric kettle whose heating element has a mean (temperature averaged) resistance of 20 Ω . The rms voltage in the mains is 200 V. Ignoring heat loss from the kettle, time taken for water to evaporate fully, is close to

[Specific heat of water = 4200 J/(kg °C),

(0) 22 minutes

Answer (3)

Sol.
$$\triangle Q = 1 \times 4200 \times 80 + 2260 \times 10^3$$
 J

$$\Delta Q = I_{rms} V_{rms} t = 200 \times \frac{200}{20} t = 2000t$$
$$\Rightarrow t = \frac{2596}{2} s \approx 21.6 \text{ minutes} \approx 22 \text{ minutes}$$

7. An electron, moving along the x-axis with an initial energy of 100 eV, enters a region of magnetic field $\vec{B} = (1.5 \times 10^{-3} \text{ T}) \hat{k}$ at S (See figure). The field extends between x = 0 and x = 2 cm. The electron is detected at the point Q on a screen placed 8 cm away from the point S. The distance d between P and Q (on the screen) is

(electron's charge = 1.6×10^{-19} C, mass of electron = 9.1×10^{-31} kg)



Answer (2)

Sol. Radius of path,
$$R = \frac{mv}{eB} = \frac{\sqrt{2m KE}}{eB}$$

= $\frac{\sqrt{2 \times 9.1 \times 10^{-31} \times 100e}}{e \times 1.5 \times 10^{-3}} = \frac{\sqrt{2 \times 9.1 \times 10^{-29}}}{\sqrt{1.6 \times 10^{-19} \times 1.5 \times 10^{-3}}} m$
 $\frac{Q}{\sqrt{1.6 \times 10^{-19} \times 1.5 \times 10^{-3}}}$
= $\frac{3.37 \times 10^{-5}}{1.5 \times 10^{-3}} \times 100 \text{ cm} = 2.25 \text{ cm}$

$$\sin\theta=\frac{2}{2.25}=\frac{8}{9}$$



- = 1.22 + 11.64 = 12.86 cm
- 8. A Carnot engine has an efficiency of $\frac{1}{6}$. When

the temperature of the sink is reduced by 62°C, its efficiency is doubled. The temperatures of the source and the sink are, respectively,

(1) 99°C, 37°C
(2) 37°C, 99°C
(3) 124°C, 62°C
(4) 62°C, 124°C

Answer (1)

- Sol. $\frac{1}{6} = 1 \frac{T_C}{T_H} \implies \frac{T_C}{T_H} = \frac{5}{6} \qquad ...(i)$ $\frac{1}{3} = 1 - \frac{(T_C - 62)}{T_H} \implies \frac{T_C - 62}{T_H} = \frac{2}{3} \qquad ...(ii)$ $\frac{T_C - 62}{T_C} = \frac{2 \times 6}{3 \times 5} = \frac{4}{5}$ $\Rightarrow T_C = 310 \text{ K} = 37^{\circ}\text{C}$ $T_H = 372 \text{ K} = 99^{\circ}\text{C}$
- 9. A small speaker delivers 2 W of audio output. At what distance from the speaker will one detect 120 dB intensity sound?

[Given reference intensity of sound as 10^{-12} W/m²]

(1) 30 cm	(2)	40	cm
-----------	-----	----	----

(3) 10 cm (4) 20 cm

Answer (2)

Sol.
$$120 = 10 \log_{10} \frac{1}{10^{-12}}$$

 $\Rightarrow \frac{1}{10^{-12}} = 10^{12} \Rightarrow I = 1 W/m^2$
 $\frac{2}{4\pi r^2} = 1 \Rightarrow r = \sqrt{\frac{2}{4\pi}} m = 0.399 m$
 $= 40 \text{ cm}$

10. A system of three polarizers P_1 , P_2 , P_3 is set up such that the pass axis of P_3 is crossed with respect to that of P_1 . The pass axis of P_2 is inclined at 60°C to the pass axis of P_3 . When a beam of unpolarized light of intensity I_0 is incident on P_1 , the intensity of light transmitted by the three polarizers is I. The ratio (I_0/I) equals (nearly) :

(1) 1.80	(2) 5.33
(3) 10.67	(4) 16.00

$$\frac{I_0}{I_0} \xrightarrow{P_1} \frac{I_0}{2} \xrightarrow{P_2} \xrightarrow{P_3} \xrightarrow{P_3} \xrightarrow{I_0} \xrightarrow{I_0}$$

11. A solid sphere, of radius R acquires a terminal velocity v_1 when falling (due to gravity) through a viscous fluid having a coefficient of viscosity η . The sphere is broken into 27 identical solid spheres. If each of these spheres acquires a terminal velocity, v_2 , when falling through the

same fluid, the ratio
$$\left(\frac{\mathbf{v_1}}{\mathbf{v_2}} \right)$$
 equals :

(1)
$$\frac{1}{27}$$
 (2) 9
(3) $\frac{1}{9}$ (4) 27

Answer (2)

Sol.
$$R^3 = 27r^3$$

 $\Rightarrow R = 3r$ $\frac{V_1}{V_2} = \left(\frac{R}{r}\right)^2 = 9$

12. Figure shows a DC voltage regulator circuit, with a Zener diode of breakdown voltage = 6 V. If the unregulated input voltage varies between 10 V to 16 V, then what is the maximum Zener current?





$$I_{S} = \frac{10}{2 \times 10^{3}} = 5 \text{ mA}$$
$$I_{L} = \frac{6}{4 \times 10^{3}} = 1.5 \text{ mA}$$
$$I_{Z(max)} = I_{S} - I_{L} = 3.5 \text{ mA}$$

13. A smooth wire of length $2\pi r$ is bent into a circle and kept in a vertical plane. A bead can slide smoothly on the wire. When the circle is rotating with angular speed ω about the vertical diameter AB, as shown in figure, the bead is at rest with respect to the circular ring at position P as shown. Then the value of ω^2 is equal to:







a liquid of refractive index $\mu_1(\mu_1 < \mu_2)$. A ray is incident on the face AB at an angle θ (shown in the figure). Total internal reflection takes place at point E on the face BC.



Then θ must satisfy:

(1)
$$\theta > \sin^{-1}\frac{\mu_1}{\mu_2}$$
 (2) $\theta < \sin^{-1}\frac{\mu_1}{\mu_2}$
(3) $\theta > \sin^{-1}\sqrt{\frac{\mu_2^2}{\mu_1^2} - 1}$ (4) $\theta < \sin^{-1}\sqrt{\frac{\mu_2^2}{\mu_1^2} - 1}$

Answer (4)

 \Rightarrow



$$\mu_1 \sin\theta = \mu_2 \sin r_2 = \mu_2 \sin (90^\circ - r_1)$$

$$\sin\theta = \frac{\mu_2}{\mu_1} \sin (90^\circ - r_1) = \frac{\mu_2}{\mu_1} \cos r_1$$
$$\theta < \sin^{-1} \left[\frac{\mu_2}{\mu_1} \frac{\sqrt{\mu_2^2 - \mu_1^2}}{\mu_2} \right]$$

$$\Rightarrow \quad \theta < \sin^{-1} \left[\sqrt{\frac{\mu_2^2}{\mu_1^2}} - \mathbf{1} \right]$$

15. A particle is moving with speed $v = b\sqrt{x}$ along positive x-axis. Calculate the speed of the particle at time t = τ (assume that the particle is at origin at t = 0).

2

(1)
$$b^2 \tau$$
 (2) $\frac{b^2 \tau}{4}$

(3)
$$\frac{b^2 \tau}{2}$$
 (4) $\frac{b^2 \tau}{\sqrt{2}}$

Answer (3)



$$\Rightarrow \int_{0}^{x} \frac{dx}{\sqrt{x}} = \int_{0}^{\tau} b dt$$

$$\Rightarrow 2\sqrt{x} = b\tau \qquad \dots (ii)$$

$$\Rightarrow v = b \cdot \frac{b\tau}{2} = \frac{b^{2}\tau}{2}$$

16. Two sources of sound S_1 and S_2 produce sound waves of same frequency 660 Hz. A listener is moving from source S_1 towards S_2 with a constant speed u m/s and he hears 10 beats/s. The velocity of sound is 330 m/s. Then, u equals:

(1) 10.0 m/s	(2) 5.5 m/s
(3) 2.5 m/s	(4) 15.0 m/s

Answer (3)

Sol.
$$\begin{array}{c} 0 \text{ bs.} \\ \mathbf{S}_1 & \mathbf{v} & \mathbf{S}_2 \end{array} \\ \mathbf{n}_1 = \frac{\mathbf{v} - \mathbf{u}}{\mathbf{v}} \times \mathbf{n}_0 \\ \mathbf{n}_2 = \frac{\mathbf{v} + \mathbf{v}}{\mathbf{v}} \times \mathbf{n}_0 \\ \Rightarrow & \mathbf{n}_2 - \mathbf{n}_1 = \frac{2\mathbf{u} \times \mathbf{n}_0}{\mathbf{v}} \\ \Rightarrow & \mathbf{10} = \frac{2 \times \mathbf{u} \times 660}{330} \\ \Rightarrow & \mathbf{u} = 2.5 \text{ m/s} \end{array}$$

17. Consider the LR circuit shown in the figure. If the switch S is closed at t = 0 then the amount of charge that passes through the battery



$$\Rightarrow \int dq = \frac{E}{R} \int \left(1 - e^{-\frac{t}{\tau}} \right) dt$$
$$\Rightarrow Q = \frac{E}{R} \times \left[t + e^{-\frac{t}{\tau}} \times \tau \right]_{0}^{t}$$
$$= \frac{E}{R} \times \left[\frac{L}{R} + \frac{L}{R} \cdot e^{-1} - \frac{L}{R} \right]$$
$$\Rightarrow Q = \frac{EL}{eR^{2}} = \frac{EL}{27R^{2}}$$

18. The ratio of the weights of a body on the Earth's surface to that on the surface of a

planet is 9 : 4. The mass of the planet is $\frac{1}{9}$ th of that of the Earth. If 'R' is the radius of the Earth, what is the radius of the planet ? (Take the planets to have the same mass density)

(1) $\frac{R}{4}$	(2) R / 3
(3) R / <u>9</u>	(4) R / 2

Sol.
$$\frac{g_e}{g_p} = \frac{9}{4}$$

 $\Rightarrow \frac{M_e \times R_p^2}{R_e^2 \times M_p} = \frac{9}{4}$
 $\Rightarrow 9 \times \left(\frac{R_p}{R_e}\right)^2 = \frac{9}{4}$
 $\Rightarrow R_p = \frac{R_e}{2}$
 $\therefore R_p = \frac{R}{2}$

19. In an amplitude modulator circuit, the carrier wave is given by,

C(t) = 4 sin (20000 π t) while modulating signal is given by, m(t) = 2 sin (2000 π t). The values of modulation index and lower side band frequency are :

- (1) 0.5 and 10 kHz
- (2) 0.3 and 9 kHz
- (3) 0.4 and 10 kHz
- (4) 0.5 and 9 kHz

Answer (4)



$$f_c = \frac{\omega_c}{2\pi} = \frac{20000\pi}{2\pi} = 10000 \text{ Hz}$$

 $f_m = \frac{\omega_m}{2\pi} = \frac{2000\pi}{2\pi} = 1000 \text{ Hz}$
∴ Lower band side = (10000 – 1000) Hz

= 9 kHz

20. Find the magnetic field at point P due to a straight line segment AB of length 6 cm carrying a current of 5 A. (See figure)

$$(\mu_0 = 4\pi \times 10^{-7} \text{ N-A}^{-2})$$

Sol.

$$5 \text{ cm}$$

$$4 \text{ cm}$$

$$\theta_2 (\cdot)$$

$$\theta_3 (\cdot)$$

$$\theta_4 (\cdot$$

- 21. Consider an electron in a hydrogen atom, revolving in its second excited state (having radius 4.65 Å). The de-Broglie wavelength of this electron is :
 - (1) 3.5 Å
 - (2) 12.9 Å
 - (3) 9.7 Å
 - (4) 6.6 Å



22. Let a total charge 2 Q be distributed in a sphere of radius R, with the charge density given by
$$\rho(\mathbf{r}) = \mathbf{kr}$$
, where r is the distance from the centre. Two charges A and B, of –Q each, are placed on diametrically opposite points, at equal distance, a, from the centre. If A and B do not experience any force, then :

(1)
$$a = \frac{3R}{2^{\frac{1}{4}}}$$
 (2) $a = R/\sqrt{3}$
(3) $a = 2^{-1/4}R$ (4) $a = 8^{-1/4}R$

Answer (4)

Sol.
$$E \times 4\pi a^2 = \frac{q_{in}}{\varepsilon_0}$$

 $q_{in} = \int_0^a (kr) \times 4\pi r^2 dr$
 $= 4\pi k \cdot \left(\frac{a^4}{4}\right)$
 $\therefore E \times 4\pi a^2 = \frac{4\pi k a^4}{4\varepsilon_0} \Rightarrow E = \frac{ka^2}{4\varepsilon_0}$
 $\therefore \frac{ka^2}{4\varepsilon_0} \times Q = \frac{Q^2}{4\pi\varepsilon_0 \times 4a^2}$
 $\frac{2Q}{\pi R^4} \cdot \frac{a^2Q}{4\varepsilon_0} = \frac{Q^2}{4\pi\varepsilon_0 \times 4a^2} \Rightarrow 8a^4 = R^4$
 $\Rightarrow 8^{1/4} \cdot a = R$
 $\Rightarrow a = \frac{R}{8^{1/4}}$
As, $2Q = \frac{4\pi k R^2}{4} \Rightarrow k = \frac{2Q}{\pi R^4}$



triangle of side 1 m (as shown in the figure). The (x, y) coordinates of the centre of mass will be :



Answer (4)



24. Half lives of two radioactive nuclei A and B are 10 minutes and 20 minutes, respectively. If, initially a sample has equal number of nuclei, then after 60 minutes, the ratio of decayed numbers of nuclei A and B will be :

(1) 9:8	(2) 3:8
(3) 8:1	(4) 1:8

Answer (1)

Sol.
$$N_A = \frac{N_0}{2^{\left(\frac{60}{20}\right)}} = \frac{N_0}{64}$$

 $N_B = \frac{N_0}{2^{\left(\frac{60}{20}\right)}} = \frac{N_0}{8}$

- $\frac{1}{N_0 \frac{N_0}{8}}$
- $=\frac{63\times8}{64\times7}=\frac{9}{8}$ 25. A tuning fork of frequency 480 Hz is used in an experiment for measuring speed of sound (v) in air by resonance tube method. Resonance is observed to occur at two successive lengths of the air column,
 - $I_1 = 30$ cm and $I_2 = 70$ cm. Then, v is equal to
 - (1) 332 ms^{-1} (2) 384 ms^{-1}
 - (3) 379 ms⁻¹ (4) 338 ms⁻¹

Answer (2)

Sol. $I_1 = 30 \text{ cm}, I_2 = 70 \text{ cm}$

$$\therefore \quad \frac{\lambda}{2} = (I_2 - I_1) = 40 \text{ cm}$$

$$\Rightarrow \quad \lambda = 80 \text{ cm}$$

$$\therefore \quad \mathbf{U} = v\lambda = 480 \times (0.8) \text{ m/s}$$

26. The number density of molecules of a gas depends on their distance r from the origin as,

 $n(r) = n_0 e^{-\alpha r^4}$. Then the total number of molecules is proportional to

(1) $n_0 \alpha^{-3/4}$ (2) $\sqrt{n_0} \alpha^{1/4}$ (3) $n_0 \alpha^{1/4}$ (4) $n_0 \alpha^{-3}$

Answer (1)

Sol.
$$n = n_0 e^{-\alpha r^4}$$

$$\Rightarrow \int dN = \int n_0 e^{-\alpha r^4} \times 4\pi r^2 dr$$

$$\Rightarrow N = 4\pi n_0 \cdot \int_0^{\infty} r^2 e^{-\alpha r^4} dr$$
Put $\sqrt{\alpha} r^2 = t$
 $2\sqrt{\alpha} r dr = dt$
 $N = \frac{4\pi n_0}{2\sqrt{\alpha}} \int_0^{\infty} \frac{t^2 e^{-t^2}}{\alpha^{\frac{1}{4}}} dt$
 $= \frac{4\pi n_0}{2\alpha^{\frac{3}{4}}} \int t^{\frac{1}{2}} e^{-t^2} dt$
 $N \propto n_0 \alpha^{-\frac{3}{4}}$





28. Two particles are projected from the same point with the same speed u such that they have the same range R, but different maximum heights, h_1 and h_2 . Which of the following is correct?

(1)
$$R^2 = 4 h_1 h_2$$
 (2) $R^2 = 16 h_1 h_2$
(3) $R^2 = 2 h_1 h_2$ (4) $R^2 = h_1 h_2$

Answer (2)

Sol. At complementary angles, ranges are equal.

$$\therefore \quad h_1 = \frac{u^2 \sin^2 \theta}{2g}, h_2 = \frac{u^2 \cos^2 \theta}{2g}$$
$$\therefore \quad h_1 \times h_2 = \left(\frac{2u^2 \sin \theta \cos \theta}{g}\right)^2 \times \left(\frac{1}{16}\right)$$
$$\Rightarrow \quad 16h_1h_2 = R^2$$

What would be the heat energy absorbed by the gas, in this process?

(1) 30 J	(2) 35 J
(3) 25 J	(4) 40 J

Answer (2)

Sol. W = nR∆T = 10 J

$$\Delta \mathbf{Q} = (\Delta \mathbf{Q})_{\mathbf{P}} = \mathbf{n}\mathbf{C}_{\mathbf{P}}\Delta \mathbf{T}$$
$$\therefore \quad \Delta \mathbf{Q} = \mathbf{n} \times \frac{7}{2}\mathbf{R} \times \Delta \mathbf{T}$$
$$= \frac{7}{2} \times (10)$$
$$= 35 \text{ J}$$

- 30. A plane electromagnetic wave having a frequency v = 23.9 GHz propagates along the positive z-direction in free space. The peak value of the electric field is 60 V/m. Which among the following is the acceptable magnetic field component in the electromagnetic wave?
 - (1) $\vec{=} 0 \sin(0.5 \times 10^3 \text{ x} + 1.5 \times 10^{11} \text{ t}) \hat{\text{k}}$

(2)
$$\vec{B} = 2 \times 10^7 \sin(0.5 \times 10^3 z + 1.5 \times 10^{11} t)\hat{i}$$

(3) $\vec{B} = 2 \times 10^{-7} \sin(0.5 \times 10^3 z - 1.5 \times 10^{11} t)\hat{i}$

(4)
$$\vec{B} = 2 \times 10^{-7} \sin(1.5 \times 10^2 x + 0.5 \times 10^{11} t) \hat{j}$$

Answer (3)

Sol.
$$B_0 = \frac{E_0}{c} = \frac{60}{3 \times 10^8} = 2 \times 10^{-7} T$$

 $v = 23.9 \times 10^9 Hz$
 $\therefore \quad \omega = 2\pi v = 2 \times 3.142 \times 23.9 \times 10^9$
 $= 1.5 \times 10^{11} S^{-1}$
 $\therefore \quad c = \frac{\omega}{k} \implies \qquad k = \frac{\omega}{c} = \frac{1.5 \times 10^{11}}{3 \times 10^8}$
 $= 0.5 \times 10^3$
 $\therefore \quad \vec{B} = 2 \times 10^{-7} \sin(0.5 \times 10^3 z - 1.5 \times 10^{11} t)\hat{i}$

PART-B : CHEMISTRY

1. The INCORRECT statement is :

- (1) Lithium is least reactive with water among the alkali metals.
- (2) $LiNO_3$ decomposes on heating to give $LiNO_2$ and O_2 .
- (3) Lithium is the strongest reducing agent among the alkali metals.
- (4) LiCl crystallises from aqueous solution as LiCl.2H₂O.

Answer (2)

Sol.
$$4LiNO_3 \xrightarrow{\Delta} 2Li_2O + 4NO_2 + O_2$$

- 2. The pair that has similar atomic radii is :
 - (1) Mo and W (2) Ti and Hf
 - (3) Sc and Ni (4) Mn and Re

Answer (1)



Due to lanthanoid contraction, radius of Mo and W are almost same.

3. The C–C bond length is maximum in :

(1) graphite	(2) C ₆₀
(3) diamond	(4) C ₇₀

Answer (3)

Sol. Carbon-carbon bond length is maximum in diamond

Species	C – C bond length
Diamond	154 pm
Graphite	141.5 pm
C ₆₀	138.3 pm and 143.5 pm
	(double bond) (single bond)

- 4. The decreasing order of electrical conductivity of the following aqueous solutions is:
 - 0.1 M Formic acid (A),
 - 0.1 M Acetic acid (B),
 - 0.1 M Benzoic acid (C),

(1) A > B > C	(2) A > C > B
---------------	---------------

(3)
$$C > B > A$$
 (4) $C > A > B$

Answer (2)

Sol. HCOOH CH_3COOH C_6H_5COOH (A) (B) (C)

Order of acidic strength

 $\begin{array}{cc} \mbox{HCOOH} > \mbox{C}_6\mbox{H}_5\mbox{COOH} > \mbox{CH}_3\mbox{COOH} \\ \mbox{pK}_a & 3.8 & 4.2 & 4.8 \end{array}$

More the acidic strength more will be the dissociation of acid into ions and more will be conductivity.

... Order of conductivity:

$$\begin{array}{rll} \mbox{HCOOH} > \mbox{C}_6\mbox{H}_5\mbox{COOH} > \mbox{CH}_3\mbox{COOH} \\ \Rightarrow & \mbox{A} & > & \mbox{C} & > & \mbox{B} \end{array}$$

- 5. The compound used in the treatment of lead poisoning is :
 - (1) EDTA (2) desferrioxime B
 - (3) Cis-platin (4) D-penicillamine

Answer (1)

Sol. EDTA is used in the treatment of lead poisoning.

options :

Assertion (A) : Vinyl halides do not undergo nucleophilic substitution easily.

Reason (R) : Even though the intermediate carbocation is stabilized by loosely held π -electrons, the cleavage is difficult because of strong bonding.

- (1) Both (A) and (R) are correct statements and (R) is the correct explanation of (A).
- (2) Both (A) and (R) are correct statements but (R) is not the correct explanation of (A).
- (3) Both (A) and (R) are wrong statements.
- (4) (A) is a correct statement but (R) is a wrong statement.

Answer (4)

Sol.
$$CH_2 = CH - CI CH_2 = C - H \text{ or } H - C - C - H$$

Fig. (1) Also,

 $\dot{cH_2} = CH \stackrel{\checkmark}{-} \dot{ci} \iff \overset{\Theta}{CH_2} - CH = \dot{ci}$

Fig. (2)

Due to partial double bond character of C – Halogen bond, Halogen leaves with great difficulty, if at all it does. Hence, vinyl halides do not undergo nucleophilic substitution easily. Assertion is correct.

Fig. (3)

Intermediate carbocation is not stabilised by loosely held π -electrons because empty orbital (see Fig. (3)), being at 90°, cannot overlap with p-orbitals of π bond. Reason is wrong.

7. Benzene diazonium chloride on reaction with aniline in the presence of dilute hydrochloric acid gives :



Answer (3)



Sol.
$$H^*$$
 + H^*

$$\sim$$
 N = N \sim NH₂

p-aminoazobenzene

- 8. 25 g of an unknown hydrocarbon upon burning produces 88 g of CO_2 and 9 g of H_2O . This unknown hydrocarbon contains :
 - (1) 22 g of carbon and 3 g of hydrogen
 - (2) 24 g of carbon and 1 g of hydrogen
 - (3) 20 g of carbon and 5 g of hydrogen
 - (4) 18 g of carbon and 7 g of hydrogen

Answer (2)

Sol. $C_x H_y + \left(x + \frac{y}{4}\right) O_2 \longrightarrow x CO_2 + \frac{y}{2} H_2 O_2$ (excess) 88 gm 25 g 9 g 2 moles $\frac{1}{2}$ moles \Rightarrow x = 2 \Rightarrow y = 1 \therefore x = 2 and y = 1 Hydrocarbon : $(C_2H)_n$ 2 mol carbon contains 24 g 1 mol hydrogen contains 1 g 9. In which one of the following equilibria, $K_{P} \neq K_{C}$? (1) $2HI(g) \rightleftharpoons H_2(g) + I_2(g)$ (2) $2NO(g) \rightleftharpoons N_2(g) + O_2(g)$ (3) $NO_2(g) + SO_2(g) \rightleftharpoons NO(g) + SO_3(g)$

(4) $2C(s) + O_2(g) \rightleftharpoons 2CO(g)$

Answer (4)

- Sol. :: $K_P = K_C \cdot (RT)^{\Delta n_g}$:. If $\Delta n_g \neq 0$ then $K_P \neq K_C$ $2C(s) + O_2(g) \implies 2CO(g)$ $\Delta n_g = +1$
 - \Rightarrow K_P = K_C · (RT)¹

- (1) They can scatter light.
- (2) The range of diameters of colloidal particles is between 1 and 1000 nm.
- (3) The osmotic pressure of a colloidal solution is of higher order than the true solution at the same concentration.
- (4) They are larger than small molecules and have high molar mass.

Answer (3)

Sol. The osmotic pressure of a colloidal solution is of lower order than that of true solution at the same concentration due to association of solute molecule till they acquire colloidal dimensions.

 π = iCRT

i is less in colloidal solution than true solution

- 11. The primary pollutant that leads to photochemical smog is
 - (1) Nitrogen oxides (2) Sulphur dioxide
 - (3) Ozone (4) Acrolein

Answer (1)

Sol. In photochemical smog :

Primary pollutants \Rightarrow NO₂, hydrocarbons Secondary pollutants \Rightarrow Ozone, acrolein

- 12. The INCORRECT match in the following is
 - (1) $\Delta G^{\circ} = 0, K = 1$ (2) $\Delta G^{\circ} < 0, K < 1$
 - (3) $\Delta G^{\circ} > 0, K < 1$ (4) $\Delta G^{\circ} < 0, K > 1$

Answer (2)

Sol. $\Delta G^\circ = -RT \ln K$

: If K > 1 then
$$\Delta G^{\circ} < 0$$

If K < 1 then
$$\triangle G^{\circ} > 0$$

If K = 1 then $\triangle G^\circ = 0$

13. NO_2 required for a reaction is produced by the decomposition of N_2O_5 in CCI_4 as per the equation,

 $2N_2O_5(g) \rightarrow 4 \ NO_2(g) \textbf{+} O_2(g).$

The initial concentration of N_2O_5 is 3.00 mol L⁻¹ and it is 2.75 mol L⁻¹ after 30 minutes. The rate of formation of NO₂ is

(1) 1.667 × 10^{-2} mol L⁻¹ min⁻¹

- (2) 4.167 × 10⁻³ mol L⁻¹ min⁻¹
- (3) 8.333 × 10⁻³ mol L⁻¹ min⁻¹
- (4) 2.083 × 10⁻³ mol L⁻¹ min⁻¹

Answer (1)



rate =
$$\frac{-1}{2} \frac{d[N_2O_5]}{dt} = \frac{1}{4} \frac{d[NO_2]}{dt} = \frac{d[O_2]}{dt}$$

Since, instant of finding rate of formation of NO_2 is not mentioned, hence

$$\therefore \quad \frac{-\Delta[N_2O_5]}{\Delta t} = -\frac{(2.75 - 3)}{30} = \frac{0.25}{30} \,\mathrm{M \,min^{-1}}$$
$$\therefore \quad \frac{\Delta[NO_2]}{\Delta t} = 2 \times \frac{-\Delta[N_2O_5]}{\Delta t}$$
$$= 2 \times \frac{0.25}{30}$$
$$= 1.67 \times 10^{-2} \,\mathrm{M \,min^{-1}}$$

14. The correct name of the following polymer is



(1) Polyisobutane (2) Polyisoprene

(3) Polyisobutylene (4) Polytert-butylene

Answer (3)



15. A solution is prepared by dissolving 0.6 g of urea (molar mass = 60 g mol⁻¹) and 1.8 g of glucose (molar mass = 180 g mol⁻¹) in 100 mL of water at 27°C. The osmotic pressure of the solution is

(R = 0.08206 L atm K⁻¹ mol⁻¹)

(1) 1.64 atm	(2) 2.46 atm
(3) 8.2 atm	(4) 4.92 atm

Answer (4)

Sol. Osmotic pressure $(\pi) = CRT$

Solute : urea and glucose

$$\therefore \pi = (C_1 + C_2) RT$$

= $\left(\frac{0.6}{60 \times 0.1} + \frac{1.8}{180 \times 0.1}\right) \times 0.0821 \times 300$
= $0.2 \times 0.0821 \times 300$

= 4.926 atm

centered cubic structure are, respectively

Answer (4)

Sol. No. of atoms in

Simple cubic unit cell
$$=\frac{1}{8} \times 8 = 1$$

B.C.C. unit cell $=\frac{1}{8} \times 8 + 1 \times 1 = 2$

FCC unit cell =
$$\frac{1}{8} \times 8 + \frac{1}{2} \times 6 = 4$$

- 17. In comparison to boron, berylium has
 - (1) Greater nuclear charge and lesser first ionisation enthalpy.
 - (2) Greater nuclear charge and greater first ionisation enthalpy.
 - (3) Lesser nuclear charge and greater first ionisation enthalpy.
 - (4) Lesser nuclear charge and lesser first ionisation enthalpy.

Answer (3)

Sol. Nuclear charge B > Be

Ionisation energy Be > B

(due to ns² outer electronic configuration)
Be =
$$1s^2 2s^2$$
 (more stable)
B = $1s^2 2s^2 2p^1$

18. The IUPAC name for the following compound is



- (1) 3-methyl-4-(1-methylprop-2-ynyl)-1-heptene
- (2) 3-methyl-4-(3-methylprop-1-enyl)-1-heptyne
- (3) 3,5-dimethyl-4-propylhept-1-en-6-yne
- (4) 3,5-dimethyl-4-propylhept-6-en-1-yne Answer (3)



a buffer solution of pH = 12 is:

(1)
$$1.84 \times 10^{-9}$$
 M (2) 6.23×10^{-11} M
(3) $\frac{2.49}{1.84} \times 10^{-9}$ M (4) 2.49×10^{-10} M

Answer (4)

Sol. $Cd(OH)_2 \rightleftharpoons Cd^{2+} + 2QH^{-}$

At equilibrium, $K_{sp} = S(2S)^2$ = 4S³

$$\Rightarrow$$
 K_{sp} = 4 × (1.84 × 10⁻⁵)³

Solubility in buffer solution having pH = 12 $[OH^{-}] = 10^{-2}$

Cd(OH)₂ ⇒ Cd²⁺ + 2OH⁻_{25'+10⁻²=10⁻²}
∴ K_{sp} = 4 × (1.84 × 10⁻⁵)³ = S'(10⁻²)²
⇒ S' =
$$\frac{2.49 \times 10^{-15}}{10^{-4}}$$
 = 2.49×10⁻¹⁰ M

- 20. The temporary hardness of a water sample is due to compound X. Boiling this sample converts X to compound Y. X and Y, respectively are:
 - (1) $Ca(HCO_3)_2$ and $Ca(OH)_2$
 - (2) $Mg(HCO_3)_2$ and $Mg(OH)_2$
 - (3) $Mg(HCO_3)_2$ and $MgCO_3$
 - (4) $Ca(HCO_3)_2$ and CaO

Answer (2)

Sol. Temporary hardness is caused by bicarbonates of calcium and magnesium

 $\underset{(X)}{\text{Mg}(\text{HCO}_3)_2(\text{aq})} \xrightarrow{\text{Boiling}} \text{Mg}(\text{OH})_2 \downarrow + 2\text{CO}_2 \uparrow$

21. Heating of 2-chloro-1-phenylbutane with EtOK/ EtOH gives X as the major product. Reaction of X with Hg(OAc)₂/H₂O followed by NaBH₄ gives Y as the major product. Y is:





22. What will be the major product when m-cresol is reacted with propargyl bromide $(HC = C-CH_2Br)$ in presence of K_2CO_3 in acetone ?



Answer (3)



Meta cresol is neutralised by K_2CO_3 . The phenoxide ion attacks at the C-atom carrying Br-atom to give ether following S_N^2 mechanism.

- 23. Among the following, the energy of 2s orbital is lowest in:
 - (1) Li (2) K
 - (3) H (4) Na

Answer (2)

- Sol. As the value of Z (atomic number) increases, energy of orbitals decreases (becomes more –ve value)
 - ... order of energy of 2s orbital is

H > Li > Na > K



gaseous product. MnO_2 reacts with NaCl and concentrated H_2SO_4 to give a pungent gas Z. X, Y and Z respectively are:

(1) K₂MnO₄, KMnO₄ and Cl₂

- (2) K_3MnO_4 , K_2MnO_4 and Cl_2
- (3) K_2MnO_4 , $KMnO_4$ and SO_2
- (4) $KMnO_4$, K_2MnO_4 and Cl_2

Answer (4)

Sol.
$$2KMnO_4 \xrightarrow{513K} K_2MnO_4 + MnO_2 + O_2$$

(X) (Y) (Y) (Y) (X)

 $\underset{\text{Conc.}}{\text{MnO}_2} \text{+} 4\text{NaCl} \text{+} 4\text{H}_2\text{SO}_4 \rightarrow \text{MnCl}_2 \text{+} 4\text{NaHSO}_4 \text{+}$

pur

- 25. The correct statement is
 - (1) Pig iron is obtained from cast iron.
 - (2) Leaching of bauxite using concentrated NaOH solution gives sodium aluminate and sodium silicate.
 - (3) The blistered appearance of copper during the metallurgical process is due to the evolution of CO₂.
 - (4) The Hall-Heroult process is used for the production of aluminium and iron.

Answer (2)

Sol. During metallurgy of aluminium, when bauxite (powdered ore) is treated with NaOH (conc.), sodium aluminate (Na[Al(OH)₄]) is formed and impurity SiO₂ dissolves by forming sodium silicate (Na₂SiO₃)

 $AI_2O_3(s)$ + 2NaOH(aq) + $3H_2O(I) \rightarrow$

2Na[Al(OH)₄](aq)

- 26. Which of the given statements is INCORRECT about glycogen?
 - (1) It is present in some yeast and fungi.
 - (2) It is a straight chain polymer similar to amylose.
 - (3) It is present in animal cells.
 - (4) Only α -linkages are present in the molecule.

Answer (2)

Sol. Structure of glycogen is similar to amylopectin glycogen

- - is found in yeast and fungi
- 27. Consider the following reactions :

$$A \xrightarrow{Ag_2O} ppt$$

$$A \xrightarrow{Ag_2'H^+} B \xrightarrow{NaBH_4} C \xrightarrow{ZnCl_2} Conc. HCl} \xrightarrow{Turbidity} Within 5 minutes$$

$$A' \text{ is}$$

$$(1) CH_3 - C = CH$$

$$(2) CH_3 - C = C - CH_3$$

$$(3) CH_2 = CH_2$$

$$(4) CH = CH$$
Answer (1)
Sol. $CH_3 - C \equiv CH$

$$\downarrow HgSO_4 + H_2SO_4$$

$$OH$$

$$CH_3 - C \equiv CH$$

$$\downarrow HgSO_4 + H_2SO_4$$

$$OH$$

$$CH_3 - C = CH \xrightarrow{Lautomerise} CH_3 - \overrightarrow{C} - CH_3$$

$$\downarrow NaBH_4$$

$$Ppt. (in 5 min) \underbrace{ZnCl_2 + HCl}_{(Lucas reagent)} CH_3 - \overrightarrow{CH} - CH_3$$

28. The coordination numbers of Co and Al in

 $[Co(CI)(en)_2]CI$ and $K_3[AI(C_2O_4)_3]$, respectively, are (en = ethane-1, 2-diamine)

- (1) 3 and 3
- (2) 5 and 3
- (3) 5 and 6
- (4) 6 and 6

Answer (3)

Sol. [Co(en)₂Cl]Cl

Cl⁻ - monodentate ligand

en - bidentate ligand

 \therefore Co-ordination No. of Co = (2 × 2) + 1 = 5

 $K_3[AI(C_2O_4)_3]$

C₂O₄²⁻- bidentate ligand

 \therefore Co-ordination No. of AI = 2 × 3 = 6



(1)
$$CH_2 = CH - CI$$
 (2) $CHCI_3$
(3) CCI_4 (4) $(CH_3)_3CCI$
Answer (4)
Sol. Carbocation is formed on reaction with Ag⁺
 $R - CI + Ag^+ \longrightarrow R^+ + AgCI \downarrow_{ppt.}$
The more the stability of R^+ , the more $R - CI$ is
likely to give precipitate
 $CH_3 - CH_3 + CH_3$ is most stable carbocation
 $CH_3 - CH_3 + CH_3$ is most stable carbocation
 $CH_3 = CH, CHCI_2, CCI_3$
 $CH_3 = CH, CHCI_2, CCI_3$
 $CH_3 = CH, CHCI_2, CCI_3$

PART-C : MATHEMATICS

- 1. An ellipse, with foci at (0, 2) and (0, -2) and minor axis of length 4, passes through which of the following points?
 - (1) $(\sqrt{2}, 2)$
 - (2) $(2, 2\sqrt{2})$
 - (3) (1, 2√2)
 - **(4)** (2, √2)

Answer (1)

Sol. Let the equation of ellipse : $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

- $\therefore be = 2 \text{ and } a = 2 \quad (Given)$ $\therefore a^2 = b^2(1 - e^2)$ $\Rightarrow 4 = b^2 - 4$
- \Rightarrow b = $2\sqrt{2}$

Equation of ellipse will be $\frac{x^2}{4} + \frac{y^2}{8} = 1$

Only $(\sqrt{2}, 2)$ satisfies this equation.

2. For an initial screening of an admission test, a candidate is given fifty problems to solve. If the probability that the candidate can solve any

problem is $\frac{4}{5}$, then the probability that he is unable to solve less than two problems is:

(1)
$$\frac{316}{25} \left(\frac{4}{5}\right)^{48}$$

(2) $\frac{54}{5} \left(\frac{4}{5}\right)^{49}$
(3) $\frac{201}{5} \left(\frac{1}{5}\right)^{49}$
(4) $\frac{164}{25} \left(\frac{1}{5}\right)^{48}$

Answer (2)

р

Sol. Let p is the probability that candidate can solve a problem.

$$=\frac{4}{5}$$
; $q=\frac{1}{5}$ (: p+q=1)



$$= p^{49} [p + 50q]$$

$$= \left(\frac{4}{5}\right)^{49} \cdot \left(\frac{4}{5} + \frac{50}{5}\right)$$

$$= \frac{54}{5} \cdot \left(\frac{4}{5}\right)^{49}$$
3. The derivative of $\tan^{-1}\left(\frac{\sin x - \cos x}{\sin x + \cos x}\right)$, with respect to $\frac{x}{2}$, where $\left(x \in \left(0, \frac{\pi}{2}\right)\right)$ is :
(1) $\frac{1}{2}$
(2) $\frac{2}{3}$
(3) 2
(4) 1
Answer (3)
Sol. $f(x) = \tan^{-1}\left(\frac{\tan x - 1}{\tan x + 1}\right) = -\tan^{-1}\left(\tan\left(\frac{\pi}{4} - x\right)\right)$
 $\therefore \quad \frac{\pi}{4} - x \in \left(-\frac{\pi}{4}, \frac{\pi}{4}\right)$
So, $f(x) = -\left(\frac{\pi}{4} - x\right) = x - \frac{\pi}{4}$
Let $y = \frac{x}{2}$
 $\frac{d}{dy}f(x) = \frac{\frac{d}{dx}f(x)}{\frac{d}{dx}y} = \frac{1}{\frac{1}{2}} = 2$

The tangents to the curve $y = (x - 2)^2 - 1$ at its 4. points of intersection with the line x - y = 3, intersect at the point :

(1) $\left(\frac{5}{2}, 1\right)$	$(2)\left(-\frac{5}{2},1\right)$
$(3) \left(\frac{5}{2}, -1\right)$	$(4) \left(-\frac{5}{2},-1\right)$

Answer (3)

Sol. Equation of chord is T = 0

 $\frac{1}{dx}$

$$\Rightarrow \frac{1}{2}(y+k) = (x-2)(h-2)-1$$
$$\Rightarrow \frac{y+k}{2} = xh-2x-2h+3$$



5. If the area (in sq. units) bounded by the parabola $y^2 = 4\lambda x$ and the line $y = \lambda x$, $\lambda > 0$, is

$$\frac{1}{9}$$
, then λ is equal to :

(3)
$$4\sqrt{3}$$
 (4) $2\sqrt{6}$

Answer (2)

Sol.
$$y^2 = 4\lambda x$$
 and $y = \lambda x$
On solving ; $(\lambda x)^2 = 4\lambda x$
 $x = 0, \frac{4}{\lambda}$

Required area = $\int_{0}^{\frac{4}{\lambda}} (2\sqrt{\lambda x} - \lambda x) dx$

$$=\frac{2\sqrt{\lambda}\cdot\mathbf{x}^{3/2}}{3/2}-\frac{\lambda\mathbf{x}^{2}}{2}\Big|_{0}^{4/2}$$
$$=\frac{32}{3\lambda}-\frac{8}{\lambda}=\frac{8}{3\lambda}=\frac{1}{9}$$
$$\lambda=24$$

A group of students comprises of 5 boys and n 6. girls. If the number of ways, in which a team of 3 students can randomly be selected from this group such that there is at least one boy and at least one girl in each team, is 1750, then n is equal to :

(1) 24 (2) 27

(3) 25 (4) 28

Answer (3)

=

Sol. Number of ways of selecting three persons such that there is atleast one boy and atleast one girl in the selected persons

$$= {}^{n+5}C_3 - {}^{n}C_3 - {}^{5}C_3 = 1750$$

$$\Rightarrow \frac{(n+5)(n+4)(n+3)}{6} - \frac{n(n-1)(n-2)}{6} = 1760$$

$$\Rightarrow n^2 + 3n - 700 = 0$$

$$\Rightarrow n = -28 \text{ (rejected) or } n = 25$$

 \Rightarrow n = – 28 (rejected) or n = 25



this A.P. is :

- (1) 150
- (2) 280
- (3) 200
- (4) 120

Answer (3)

Sol. Let the common difference is 'd'.

$$a_{1} + a_{7} + a_{16} = 40$$

$$\Rightarrow a_{1} + a_{1} + 6d + a_{1} + 15d = 40$$

$$\Rightarrow 3a_{1} + 21d = 40$$

$$\Rightarrow a_{1} + 7d = \frac{40}{3}$$

$$S_{15} = \frac{15}{2} [2a_{1} + 14d]$$

$$= 15 (a_{1} + 7d)$$

$$= 15 \left(\frac{40}{3}\right)$$

$$= 200$$

- 8. The angle of elevation of the top of a vertical tower standing on a horizontal plane is observed to be 45° from a point A on the plane. Let B be the point 30 m vertically above the point A. If the angle of elevation of the top of the tower from B be 30°, then the distance (in m) of the foot of the tower from the point A is :
 - (1) $15(3+\sqrt{3})$ (2) $15(1+\sqrt{3})$

(3) $15(3-\sqrt{3})$ (4) $15(5-\sqrt{3})$

Answer (1)

Sol. Let the height of the tower be h.

Refer to diagram ; $\tan 45^\circ = \frac{h}{d} = 1$ h = d ...(i) $\tan 30^\circ = \frac{h-30}{d}$ 30 m $A5^\circ$ P $\sqrt{3}(h-30) = d$...(ii) $\sqrt{3}d = d + 30\sqrt{3}$ $d = \frac{30\sqrt{3}}{\sqrt{2}-1} = 15\sqrt{3}(\sqrt{3}+1) = 15(3+\sqrt{3})$ [cotθ]x + y = 0

- (1) Has a unique solution if $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$ and have infinitely many solutions if $\theta \in \left(\pi, \frac{7\pi}{6}\right)$.
- (2) Has a unique solution if $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right) \cup \left(\pi, \frac{7\pi}{6}\right).$
- (3) Have infinitely many solutions if $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$ and has a unique solution if $\theta \in \left(\pi, \frac{7\pi}{6}\right)$.
- (4) Have infinitely many solutions if $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right) \cup \left(\pi, \frac{7\pi}{6}\right).$

Answer (3)

Sol. There are two cases.

Case 1:
$$\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$$

So ; $[\sin\theta] = 0, [-\cos\theta] = 0, [\cot\theta] = -1$ The system of equations will be ;

 $0 \cdot x + 0 \cdot y = 0 \quad \text{and} \quad -x + y = 0$

(Infinitely many solutions)

Case 2:
$$\theta \in \left(\pi, \frac{7\pi}{6}\right)$$

So ; $[\sin\theta] = -1, [-\cos\theta] = 0,$

The system of equations will be ;

 $-x + 0 \cdot y = 0$ and $[\cot \theta] x + y = 0$

Clearly x = 0 and y = 0 (unique solution) 10. A value of $\theta \in (0, \pi/3)$, for which

$$\begin{vmatrix} 1+\cos^2\theta & \sin^2\theta & 4\cos6\theta\\ \cos^2\theta & 1+\sin^2\theta & 4\cos6\theta\\ \cos^2\theta & \sin^2\theta & 1+4\cos6\theta \end{vmatrix} = 0, \text{ is:}$$

$$(1) \quad \frac{\pi}{18}$$

$$(2) \quad \frac{7\pi}{36}$$

$$(3) \quad \frac{7\pi}{24}$$

$$(4) \quad \frac{\pi}{9}$$
Answer (4)


$$\begin{vmatrix} 2 & \sin^2 \theta & 4\cos 6\theta \\ 2 & 1+\sin^2 \theta & 4\cos 6\theta \\ 1 & \sin^2 \theta & 1+4\cos 6\theta \\ R_1 \rightarrow R_1 - R_2, R_2 \rightarrow R_2 - R_3 \\ \begin{vmatrix} 0 & -1 & 0 \\ 1 & 1 & -1 \\ 1 & \sin^2 \theta & (1+4\cos 6\theta) \\ \Rightarrow & 2+4\cos 6\theta = 0 \\ & \cos 6\theta = \frac{-1}{2} \\ \because & 6\theta \in (0, 2\pi) \\ \text{So, } 6\theta = \frac{2\pi}{3} \text{ or } \frac{4\pi}{3} \\ \Rightarrow & \theta = \frac{\pi}{9} \text{ or } \frac{2\pi}{9} \end{vmatrix}$$

11. The Boolean expression $\sim (p \ \Rightarrow \ (\sim q))$ is equivalent to

(1) (~p) ⇒ q	(2) p ∨ q
(3) p ∧ q	(4) $q \Rightarrow \sim p$

Answer (3)

- Sol. $\sim (p \Rightarrow (\sim q))$ {: $p \Rightarrow q$ is same as $\sim p \lor q$ } $\equiv \sim ((\sim p) \lor (\sim q))$ $\equiv p \land q$
- 12. The equation of a common tangent to the curves, $y^2 = 16x$ and xy = -4, is :
 - (1) x + y + 4 = 0 (2) 2x y + 2 = 0
 - (3) x 2y + 16 = 0 (4) x y + 4 = 0

Answer (4)

Sol. $y^2 = 16x$ and xy = -4

Equation of tangent to the given parabola;

$$y = mx + \frac{4}{m}$$

If this is common tangent, then

$$x\left(mx + \frac{4}{m}\right) + 4 = 0$$

$$\Rightarrow mx^{2} + \frac{4}{m}x + 4 = 0$$

$$D = 0$$

$$\frac{16}{m^{2}} = 16m$$

$$\Rightarrow m^{3} = 1 \qquad \Rightarrow m = 1$$

Equation of common tangent is y = x + 4

(where c is a constant of integration)

(1) $y^2 + 2x^3 + cx^2 = 0$ (2) $y^2 - 2x^2 + cx^3 = 0$ (3) $y^2 - 2x^3 + cx^2 = 0$ (4) $y^2 + 2x^2 + cx^3 = 0$ Answer (1) Sol. $y^2 dx - xy dy = x^3 dx$ $\Rightarrow \frac{(y dx - x dy)y}{x^2} = x dx$ y d(y) = x dx

$$\Rightarrow -yd\left(\frac{y}{x}\right) = xdx$$
$$\Rightarrow -\frac{y}{x}\cdot d\left(\frac{y}{x}\right) = dx$$
$$\Rightarrow -\frac{1}{2}\left(\frac{y}{x}\right)^2 = x + c_1$$
$$\Rightarrow 2x^3 + cx^2 + y^2 = 0$$

- 14. A plane which bisects the angle between the two given planes 2x y + 2z 4 = 0 and x + 2y + 2z 2 = 0, passes through the point :
 - (1) (1, -4, 1) (2) (2, -4, 1)(3) (1, 4, -1) (4) (2, 4, 1)

Answer (2)

Sol. Equation of angle bisectors;

$$\frac{x+2y+2z-2}{3} = \pm \frac{2x-y+2z-4}{3}$$

$$\Rightarrow x-3y-2=0 \quad \text{or} \quad 3x+y+4z-6=0$$

Only (2, -4, 1) lies on the second plane

15. Let S be the set of all $\alpha \in R$ such that the equation, $\cos 2x + \alpha \sin x = 2\alpha - 7$ has a solution. Then S is equal to :

Answer (3)

Sol. cos2x +
$$\alpha$$
sinx = 2 α - 7

$$1 - 2\sin^{2}x + \alpha \sin x = 2\alpha - 7$$

$$\Rightarrow 2\sin^{2}x - \alpha \sin x + (2\alpha - 8) = 0$$

$$\Rightarrow \sin x = \frac{\alpha \pm \sqrt{\alpha^{2} - 8(2\alpha - 8)}}{4}$$

$$\Rightarrow \sin x = \frac{\alpha \pm (\alpha - 8)}{4}$$

$$\Rightarrow \sin x = 2 \text{ (rejected)} \text{ or } \sin x = \frac{\alpha - 4}{2}$$

$$\therefore \text{ Equation has solution, then } \frac{\alpha - 4}{2} \in [1, 1]$$

$$\Rightarrow \alpha \in [2, 6]$$



 $\vec{a} = \alpha i + j + 3k$, $b = 2i + j - \alpha k$ and

 $\vec{c} = \alpha \hat{i} - 2\hat{j} + 3\hat{k}$. Then the set

S = { α : \vec{a} , \vec{b} and \vec{c} are coplanar}

- (1) contains exactly two numbers only one of which is positive
- (2) is singleton
- (3) contains exactly two positive numbers
- (4) is empty

Answer (4)

Sol. If $\vec{a}, \vec{b}, \vec{c}$ are coplanar, then

 $\left[\vec{a},\vec{b},\vec{c}\right]=0$ $\Rightarrow \begin{vmatrix} \alpha & \mathbf{1} & \mathbf{3} \\ \mathbf{2} & \mathbf{1} & -\alpha \\ \alpha & -\mathbf{2} & \mathbf{3} \end{vmatrix} = \mathbf{0}$ $\Rightarrow \alpha^2 + 6 = 0$

No value of ' α ' exist

Set S is an empty set.

17. The length of the perpendicular drawn from the point (2, 1, 4) to the plane containing the lines $\vec{z} = (\hat{i} + \hat{j}) + \gamma (\hat{i} + \hat{j}) + \hat{j}$ _

$$\mathbf{r} = (\mathbf{i} + \mathbf{j}) + \lambda(\mathbf{i} + 2\mathbf{j} - \mathbf{k}) \text{ and}$$

$$\vec{\mathbf{r}} = (\hat{\mathbf{i}} + \hat{\mathbf{j}}) + \mu(-\hat{\mathbf{i}} + \hat{\mathbf{j}} - 2\hat{\mathbf{k}}) \text{ is :}$$

(1) $\frac{1}{3}$
(2) 3
(3) $\frac{1}{\sqrt{3}}$

Answer (4)

Sol. Equation of plane containing two given lines;

$$\begin{vmatrix} x - 1 & y - 1 & z \\ 1 & 2 & -1 \\ -1 & 1 & -2 \end{vmatrix} = 0$$

$$\Rightarrow x - y - z = 0$$

The length of perpendicular from (2, 1, 4) to

this plane = $\left|\frac{2-1-4}{\sqrt{1^2+1^2+1^2}}\right| = \sqrt{3}$

$$\int \frac{\tan x + \tan \alpha}{\tan x - \tan \alpha} dx =$$

A(x) $\cos 2\alpha$ + B(x) $\sin 2\alpha$ + C, where C is a constant of integration, then the functions A(x) and B(x) are respectively :

(1) $\mathbf{x} - \alpha$ and $\log_{\mathbf{x}} |\cos(\mathbf{x} - \alpha)|$

(2) $\mathbf{x} + \alpha$ and $\log_{\alpha} |\sin(\mathbf{x} - \alpha)|$

(3) $\mathbf{x} - \alpha$ and $\log_{\mathbf{a}} |\sin(\mathbf{x} - \alpha)|$

(4) $\mathbf{x} + \alpha$ and $\log_{e} |\sin(\mathbf{x} + \alpha)|$

Answer (3)

Sol.
$$\int \frac{\tan x + \tan \alpha}{\tan x - \tan \alpha} dx$$
$$= \int \frac{\sin(x + \alpha)}{\sin(x - \alpha)} dx \qquad \text{let } x - \alpha = t$$
$$\Rightarrow dx = dt$$

$$= \int \frac{\sin(t+2\alpha)}{\sin t} dt$$

$$= [\cos 2\alpha + \sin 2\alpha \cdot \cot t] dt$$

$$= \cos 2\alpha \cdot t + \sin 2\alpha \cdot \ln |\sin t| + c$$

$$= (x - \alpha) \cdot \cos 2\alpha + \sin 2\alpha \cdot \ln |\sin(x - \alpha)| + c$$

19. If ²⁰C₁ + (2²) ²⁰C₂ + (3²) ²⁰C₃ + + (20²) ²⁰C₂₀

$$= A(2^{\beta}), \text{ then the ordered pair (A, \beta) is equal to :}$$

(1) (420, 19)
(2) (380, 19)
(3) (420, 18)
(4) (380, 18)
Answer (3)
Sol. ²⁰C₁ + 2² · ²⁰C₂ + 3² · ²⁰C₃ + ... + 20² · ²⁰C₂₀

$$= \sum_{r=1}^{20} r^{2} \cdot ^{20}C_{r}$$

$$= 20 \sum_{r=1}^{20} r \cdot ^{19}C_{r-1}$$

$$= 20 \left[\sum_{r=1}^{20} (r - 1)^{19}C_{r-1} + \sum_{r=1}^{20} {}^{19}C_{r-1} \right]$$

$$= 20 \left[19 \sum_{r=2}^{20} {}^{18}C_{r-2} + 2^{19} \right]$$

= 20[19 \cdot 2^{18} + 2^{19}]
= 20 \times 21 \times 2^{18}
= 420 \times 2^{18}
So; A = 420 and \beta = 18



•

and (2, 3). Then the centroid of this triangle is :

(1)	$\left(\frac{1}{3},2\right)$	(2)	$\left(\frac{1}{3},\frac{5}{3}\right)$
(3)	$\left(\frac{1}{3},1\right)$	(4)	$\left(1,\frac{7}{3}\right)$

Answer (1)

Sol. Co-ordinates of vertex B and C are B(-3, 0) and C(3, 4)



21. If α , β and γ are three consecutive terms of a non-constant G.P. such that the equations $\alpha x^2 + 2\beta x + \gamma = 0$ and $x^2 + x - 1 = 0$ have a common root, then $\alpha(\beta + \gamma)$ is equal to :

(1)	0	(2)	αγ
(3)	βγ	(4)	αβ

Answer (3)

Sol. $\beta^2 = \alpha \gamma$ so roots of the equation $\alpha x^2 + 2\beta x + \gamma = 0$

are $\frac{-2\beta \pm 2\sqrt{\beta^2 - \alpha\gamma}}{2\alpha} = -\frac{\beta}{\alpha}$ This root satisfy the equation $x^2 + x - 1 = 0$ $\beta^2 - \alpha\beta - \alpha^2 = 0$ $\Rightarrow \alpha\gamma - \alpha\beta - \alpha^2 = 0$ $\Rightarrow \alpha + \beta = \gamma$ Now, $\alpha(\beta + \gamma) = \alpha\beta + \alpha\gamma = \alpha\beta + \beta^2$ $= (\alpha + \beta)\beta = \beta\gamma$ 22. $\lim_{x \to 0} \frac{x + 2\sin x}{\sqrt{x^2 + 2\sin x + 1} - \sqrt{\sin^2 x - x + 1}}$ is : (1) 3 (2) 6 (3) 1 (4) 2 Answer (4) $\sqrt{x} + 2 \sin x + 1 - \sqrt{\sin x} - x + 1$

$$= \lim_{x \to 0} \frac{(x+2\sin x) \left[\sqrt{x^2+2\sin x+1}+\sqrt{\sin^2 x-x+1}\right]}{(x^2-\sin^2 x)+(x+2\sin x)}$$
$$= \lim_{x \to 0} \frac{\left[1+2\left(\frac{\sin x}{x}\right)\right] \left[\sqrt{x^2+2\sin x+1}+\sqrt{\sin^2 x-x+1}\right]}{\left(x-\frac{\sin^2 x}{x}\right)+\left(1+2\left(\frac{\sin x}{x}\right)\right)}$$
$$= \frac{3 \times 2}{3} = 2$$

23. A straight line L at a distance of 4 units from the origin makes positive intercepts on the coordinate axes and the perpendicular from the origin to this line makes an angle of 60° with the line x + y = 0. Then an equation of the line L is :

1)
$$(\sqrt{3} + 1)x + (\sqrt{3} - 1)y = 8\sqrt{2}$$

(2)
$$(\sqrt{3} - 1)x + (\sqrt{3} + 1)y = 8\sqrt{2}$$

(3)
$$\sqrt{3}x + y = 8$$

(4)
$$x + \sqrt{3}y = 8$$

Answer (1) or (2)

Sol. If perpendicular makes an angle of 60° with the line x + y = 0.

Then the perpendicular makes an angle of 15° or 75° with x-axis. So the equation of line will be



 $x\cos 75^\circ + y\sin 75^\circ = 4$ or $x\cos 15^\circ + y\sin 15^\circ = 4$

$$(\sqrt{3}-1)x + (\sqrt{3}+1)y = 8\sqrt{2} \text{ or}$$

 $(\sqrt{3}+1)x + (\sqrt{3}-1)y = 8\sqrt{2}$

By rotating the normal towards the line x + y = 0 in anticlockwise sense we get the answer (2).



$$\left(\frac{1}{60} - \frac{x^8}{81}\right) \cdot \left(2x^2 - \frac{3}{x^2}\right)^6 \text{ is equal to :}$$
(1) -108
(2) -36
(3) -72
(4) 36

Answer (2)

Sol.
$$\left(\frac{1}{60} - \frac{x^8}{81}\right) \left(2x^2 - \frac{3}{x^2}\right)^6$$

$$= \frac{1}{60} \left(2x^2 - \frac{3}{x^2}\right)^6 - \frac{x^8}{81} \left(2x^2 - \frac{3}{x^2}\right)^6$$
Coefficient of x° in $\frac{1}{60} \left(2x^2 - \frac{3}{x^2}\right)^6 - \frac{1}{81}$
coefficient of x⁻⁸ in $\left(2x^2 - \frac{3}{x^2}\right)^6$

$$= \frac{-1}{60} {}^6C_3 (2)^3 (3)^3 + \frac{1}{81} {}^6C_5 (2) (3)^5$$

$$= -72 + 36$$

$$= -36$$

25. A circle touching the x-axis at (3, 0) and making an intercept of length 8 on the y-axis passes through the point :

(1) (2, 3)	(2) (1, 5)
(3) (3, 5)	(4) (3, 10)

Answer (4)

Sol. Let centre of circle is A and circle cuts the y axis at B and C. Let mid point of chord BC is P.



 $\mathbf{AB} = \sqrt{\mathbf{PA}^2 + \mathbf{PB}^2}$

= 5 = radius of circle

Equation of circle is : $(x - 3)^2 + (y - 5)^2 = 5^2$

Only (3, 10) satisfies this equation.

Although there will be another circle satisfying the same conditions that will lie below the xaxis having equation $(x - 3)^2 + (y - 5)^2 = 5^2$ two dice), wins Rs. 12 when the throw results in the sum of 9, and loses Rs. 6 for any other outcome on the throw. Then the expected gain/ loss (in Rs.) of the person is :

(1)
$$\frac{1}{2} \log s$$
 (2) 2 gain
(3) $\frac{1}{2} gain$ (4) $\frac{1}{4} \log s$

Answer (1)

Sol. Let X be the random variable which denotes the Rs gained by the person.

$$P(X = 15) = \frac{6}{36} = \frac{1}{6}$$

$$\begin{cases}
Total cases = 36 \\
favourable cases are \\
(1,1), (2,2), (3,3), (4,4), \\
(5,5), (6,6)
\end{cases}$$

$$P(X = 12) = \frac{4}{36} = \frac{1}{9}$$

$$\begin{cases}
Favourable cases are \\
(6,3), (5,4), (4,5), (3,6)
\end{cases}$$

$$P(X = -6) = \frac{26}{36} = \frac{13}{18}$$

$$\boxed{X \quad 15 \quad 12 \quad -6}$$

		. –	-	
P (X)	1	1	13	
	6	9	18	
X · P (X)	5	4	-13	
	2	3	3	
-				

$$\mathsf{E}(\mathsf{X}) = \sum \mathsf{X} \cdot \mathsf{P}(\mathsf{X}) = \frac{5}{2} + \frac{4}{3} - \frac{13}{3} = -\frac{1}{2}$$

- 27. Let A, B and C be sets such that $\phi \neq A \cap B \subseteq C$. Then which of the following statements is not true?
 - (1) $B \cap C \neq \phi$
 - (2) (C ∪ A) ∩ (C ∪ B) = C
 - (3) If $(A C) \subseteq B$, then $A \subseteq B$
 - (4) If $(A B) \subseteq C$, then $A \subseteq C$

Answer (3)

- Sol. :: $A \cap B \subseteq C$ and $A \cap B \neq \phi$
 - (1) $\mathbf{B} \cap \mathbf{C} \neq \phi$ is correct
 - (2) $(C \cup A) \cap (C \cup B) = C \cup (A \cap B) = C$ (correct) (becasue $A \cap B \subseteq C$)
 - (3) If A = C then A C = ϕ

Clearly $\varphi \subseteq B$ but $A \subseteq B$ is not always true.

(4) \because A – B \subseteq C and A \cap B \subseteq C so A \subseteq C (correct)



$$\int_{\alpha}^{\alpha+1} \frac{dx}{(x+\alpha)(x+\alpha+1)} = \log_{e}\left(\frac{9}{8}\right) \text{ is :}$$
(1) $\frac{1}{2}$
(2) -2
(3) $-\frac{1}{2}$
(4) 2
Answer (2)
Sol. $\int_{\alpha}^{\alpha+1} \frac{dx}{(x+\alpha)(x+\alpha+1)} = \int_{\alpha}^{\alpha+1} \left[\frac{1}{x+\alpha} - \frac{1}{x+\alpha+1}\right] dx$

$$= \ln \left(\frac{\mathbf{x} + \alpha}{\mathbf{x} + \alpha + 1} \right) \Big|_{\alpha}^{\alpha + 1}$$
$$= \ln \left(\frac{2\alpha + 1}{2\alpha + 2} \cdot \frac{2\alpha + 1}{2\alpha} \right) = \ln \frac{9}{8}$$
So, $\frac{(2\alpha + 1)^2}{\alpha(\alpha + 1)} = \frac{9}{2}$
$$\Rightarrow 8\alpha^2 + 8\alpha + 2 = 9\alpha^2 + 9\alpha$$
$$\Rightarrow \alpha^2 + \alpha - 2 = 0$$
$$\Rightarrow \alpha = 1, -2$$

29. Let $z \in C$ with Im(z) = 10 and it satisfies $\frac{2z - n}{2z + n} = 2i - 1$ for some natural number n. Then : (1) n = 20 and Re(z) = 10(2) n = 20 and Re(z) = -10(3) n = 40 and Re(z) = -10

(4)
$$n = 40$$
 and $Re(z) = 10$

Answer (3)

Sol. Let z = x + 10i 2z - n = (2i - 1) (2z + n) (2x - n) + 20i = (2i - 1) ((2x + n) + 20i)Comparing real and imaginary part -(2x + n) - 40 = 2x - n and 20 = 4x + 2n - 20 $\Rightarrow 4x = -40$ 40 = -40 + 2n $\Rightarrow x = -10$ n = 40 $\Rightarrow Re(z) = -10$ minimum value at β , then



